

[Research]

Floristic study on the algae of Siahdarvishan River in Guilan Province, North Iran

M. B. Faghir ^{*1}, S. Shafii ²

1- Dept. of Biology, Faculty of science, University of Guilan, Rasht, Iran.

2- Dept. of Fisheries Sciences, Faculty of Natural Resources, University of Guilan, Someh Sara, Iran.

* Corresponding author's E-mail: marziehbeygomfaghir@gmail.com

(Received: Sept. 19.2011, Accepted: Feb.06.2012)

ABSTRACT

In the present study the freshwater algae of Siahdarvishan River were sampled on a monthly basis from December 2007 through November 2008. Three sampling stations (Gaght roudkhan, Shahkezar dame and Talab laleh) were chosen in the river. Some physical and chemical parameters (water temperature, total hardness, nitrate, nitrite, phosphate and pH) of water samples were measured. A total of 84 species were identified including: Chlorophyta, 36 species; Cyanophyta, 24 species; Bacillariophyta, 20 species; Rhodophyta, 2 species; Euglenophyta and Xanthophyta each with 1 species. Among the families, Oscillatoriaceae with 14 species showed the highest species richness. Taxonomic keys to genera and species are presented to facilitate the identification of algal flora of the region.

Keywords: Floristic study, algae, Siahdarvishan River.

INTRODUCTION

Algae play a vital role in all aquatic ecosystems by providing the food and energy base for all organisms living in lakes, ponds, streams and rivers (Addy and Green, 1966).

They have significant impact on determining water pollution and cleaning waste water (Çolak and Kaya, 1988). A floristic study on the fresh water algal flora reveals the species composition and taxonomic diversity of biological communities in an ecosystem (Pfiester *et al.*, 1980; Oguni *et al.*, 1987; Fumanti *et al.*, 1995; Kolayli *et al.*, 1998; Sahin and Akar, 2004; El-Awamri *et al.*, 2007; Škaloud, 2009; Andrejic *et al.*, 2012). In addition, it reflects the seasonal variations (Sen and Sonmez, 2006; Kim *et al.*, 2008; Ezekiel *et al.*, 2011), evolutionary processes, ecological functions and stability of aquatic ecosystems (Komulaynen 2009). The most outstanding algal flora studies in Iran are those of Löffler, (1961); Hirano (1973); Moghaddam, (1976); Compere, (1981), Nejadstari *et al.*, (2005); Zarei-Darki, (2009); Noroozi *et al.*, (2009). The present study

deals with species composition, diversity and distribution of algal flora in Siahdarvishan River.

MATERIALS AND METHODS

The study area

Siahdarvishan River is one of the important rivers entering Anzali wetland, situated at 37° 16' North latitude and 49° 21' East longitude (Abbassii *et al.* 2006) (Fig.1). In order to provide effective sampling, three sampling stations were chosen, (Gaght roudkhan, Shahkezar dame and Talab laleh). Siahdarvishan river is not seasonal and the minimum water flow is seen during summer.

Sampling procedure

The samples were collected on a monthly basis from December 2007 through November 2008, between 10 Am and 13.30 Pm. Phytoplankton samples were collected using a phytoplankton net of 5-10 µm mesh. Algal mat samples were collected using a small flat shovel; immediately

were placed in sterile plastic bags and were stored in the dark until they were transferred to the laboratory. Filamentous algae were collected by forceps from the different zones (e.g. old branches, basins, pools, stones etc). Water samples were taken in 0.5 liter. Polyethylene bottles and fixed immediately with 4% Formalin. Algal flora identification were carried out using available literatures (Anderson, 1966; Gomont, 1972; Kiyammehr, 1994; Maosen, 1983; Prescott, 1954; Smith, 1920; 1924; West, 1904; Starmach, 1972) and the samples were photographed by an Olympus Bx51 microscope. Water temperature was measured in situ using a mercury thermometer and total hardness (German degree d°), nitrate, nitrite, phosphate and pH of water samples were measured using Aquaquant® test kits based on standard methods (Adams, 2000; Apha, 1986).

RESULTS

Physicochemical factor

Physicochemical parameters in the three sampling stations are presented in Table 1. The pH value varies from 7.5 to 8.0. The minimum (6 d°) and maximum total hardness (7-8 d°) values were respectively recorded from first to third stations. Nitrate (0 mg/l), nitrite (2 mg/l) (Table 1) and Phosphate concentrations (ranged from 0.033 to 0.1 mg/l, not shown in Table 1) were almost identical in three stations. The average of air and surface water temperatures in the three stations were measured. The lowest average of air (10-27 C°) and water (3-15 C°) temperatures were

recorded in station I, while the highest average of air (12-31 C°) and water temperatures (8-28 C°) were reported from the station III (Table 1).

Taxonomical remarks

In this study a total of 84 species of fresh water algae, belonging to 34 families and 51 genera were identified (Fig. 2). Chlorophyta with 20 genera/ 36 species was the most abundant phylum followed by Cyanophyta (12 genera/ 24 species); Bacillariophyta (13 genera/ 12 species); Rhodophyta (2 genera/ 2 species), Euglenophyta and Xanthophyta (each with 1 genus/ 1 species). While Oscillatoriaceae was the most abundant family with 14 species, then Hydrodictyceae, Coelastraceae, Scenedesmaceae (5 species), Cladophoraceae Zygnemataceae, Ulothrichaceae, Cymbellaceae, Pleurosigmataceae, Naviculaceae (3 species), Nostocaceae, Coscinodiscaceae, Melosiraceae, Tabellariaceae, Cocconeidaceae (2 species) and Chamaesiphonaceae, Chaetopeltidiaceae, Chaetophoraceae, Oedogoniaceae, Botryococcaceae, Characiaceae, Microsporaceae, Chlorococcaceae, Diatomaceae, Fragilariaceae, Catenulaceae, Bacillariaceae, Diploneidaceae, Euglenaceae, Vaucheriaceae, Porphyridiaceae and Helminthocladia each with 1 species respectively were present in the studied stations (Fig. 2).




Fig.1. Location of study area

Table 1. Range of physicochemical parameters recorded in the three stations; average of air temperatures (Av. A. Temp, spring (S), summer (Su), autumn(A), winter(W), average of air temperature (Av. A. temp.), average of water temperature (Av. W. temp.), pH, total hardness (German degree d°) (T. H.), nitrate(NO₃) and nitrite (NO₂).

Stations	Av. A. Temp.				Av. W. temp.				PH	T. H d°	NO ₂ mg/l	NO ₃ mg/l
	C°				C°							
	S	Su	A	W	S	Su	A	W				
1. Gahtrudkhan	18	27	20	10	10	15	12	3	7.5-8	6	2	0
2. Shahkhezh Dame	20	30	21	12	16	25	14	7.5	7.5-8	6-8	2	0
3. Talab laleh	22	31	23	12	17	28	14	8	7.7-8	7-8	2	0




Fig 2. Number of species among the families

DISCUSSION

The taxonomic composition of freshwater algae collected from different stations is presented in Table 2. Out of 84 identified species, 12 were common in the three stations, while 20 species were recorded only from the second (Table 3), 5 species from the first (Table 4) and 4 species from the third stations (Table 5). The second station (Shahkhezh Dame) with 68 species showed the highest species richness, comparing to the first (34 species) and the third stations (24 species) (Fig. 3). This shows that algal population

(especially in Shahkhezh Dame station) is greatly influenced by water conditions and physico-chemical parameters. Increase in number of species were also significant during summer in the station I and summer - autumn months in the stations II and III (Table 4 - 5). Based on Devika *et al.* (2006) increase in algal population especially during summer is related to physical parameters (water temperature and transparency) rather than chemical conditions (which are almost identical in the three stations). On the other hand, low transparency (in the

station III), and heavy flood (in the station I) resulted the decline of algal density and physico-chemical parameters. (Rajagopal, 2010; Pundhir and Rana, 2002). According to several previous studies (Wu and Chou, 1999; Richardson *et al.*, 2000; Izaguirre, 2001 ; Ersanli and Gönülol, 2003; Tyor and

Chawla, 2012) the seasonal variations of phytoplankton are related to different environmental factors especially temperature (water and air temperatures) that regulate the growth and distribution of these organisms (Simon and Hildrev, 1998 Thebault and Rabouille, 2003).




Fig 3. Number of species and seasonal variation in the studied area

Cyanophyta(key to genera)

- 1- Cells solitary or small colony never contain more than a few cells.....2
- Cell arranged in to colony or filament4
- 2- Cell spherical or hemispherical.....3
- Cells erect, subcylindrical clubs, straight or curved;epiphytic;.....1. *Chamaesiphon*
- 3-Cell enclosed within transparent layered sheath.....2. *Chroococcus*
- Cell without a gelatinous sheath, not spherical or hemispherical-cells solitary or in pairs.....3.*Synechocystis*
- 4-Thalus colony5
- Thallus filamentous.....8
- 5- Cells globose, or hemispherical.....6- Cells elongate cylinders, forming large colonies; have no individual sheaths4. *Aphanothece*
- Cells forming irregularly globular or oval colonies7

- 7-Cells very numerous and crowded within the colonial mucilage.....6. *Microcystis*
- Cells distributed throughout the colonial mucilage.....7. *Aphanocapsa*
- 8-heterocysts present.9
- heterocysts absent.....10
- 9- Trichomes not parallel, or if so, forming indefinitely shaped flakes, composed of bead-like or barrel-shaped cell ..8.. *Anabaena*
- Trichomes parallel, forming a free-floating flake or bundle.....9. *Aphanizomenon*
- 10- Trichomes with sheath.....11
- Trichomes without a sheath; solitary, not lying in parallel bundles, sometimes tapered slightly at the anterior end, or with the apical cell swollen (capitate)10. *Oscillatoria*
- 11- Trichomes with a thin sheath; blue or black-green in color and slimy or slippery to touch.....11. *Phormidium*
- Sheath colorless or yellowish, extending beyond the end of the trichome..12. *Lyngbya*

Oscillatoria (key to species)

- 1- Trichomes straight, or slightly curved with calyptra.....2
 -Trichomes straight or slightly curved, without calyptra.....5
 2- Trichomes not constricted at the junction.. 3
 - Trichomes constricted at the junction.....4
 3-Trichome straight, slightly tapered toward both ends, 3.5-5 µm wide; with granules along the cross wall, with conical shape calyptra*O. agardhii* (Gomont).
 -Trichomes straight, 2-5.5 µm broad; granulated; terminal cell broad..... *O. prolifica* N.
 4- Trichomes straight, bent or some time twisted in apical region; cells 6-8 µm in width with granules along the cross wall, terminal cell curve.....*O. anguina* (Bory).
 - Trichomes, 2.5-5 µm broad....*O. amoema* N.5-
 Trichomes straight or slightly curved, not constricted at the junction, cells 3.5-5 µm in width with granules along the cross wall, ...*O. granulata* Gardner.
 -Trichomes straight or slightly curved, constricted at the junction.....6
 6-Cells 4-6 µm in width with granules along the cross wall.....*O. formosa* Bory.
 -Cell 2.5-8 µm in in length, with granules along the cross wall,*O. tenuis* Ag.

Phormidium (key to species)

- Trichomes straight or slightly curved 4.5-12 µm in width, not constricted d at the junction, with granules along the cross wall; terminal cell not attenuated or rounded, without calyptra*P. retzii* N.
 -Trichomes straight or slightly curved, 4-7 µm in width, not constricted at the junction; with granules along the cross wall, with diffuent sheath; apical cells not attenuated and often calyptrate.....*P. ambiguum* Gom.

Lyngbya (key to species)

- 1- Trichomes 4- 5.5 µm in width, straight or curved; not constricted at the junction, cells without granules along the cross wall.....
*L. aerugineo-coerulea* (Kutz.) Gom.
 -Trichomes more than 7 µm in diameter, straight.....2
 2- Trichomes 7- 10 µm in diameter; not constricted at the junction and not attenuated; cells with or without granules along the cross walls.....*L. martensiana* Menegh .
 - Trichomes 14- 19.5 µm in width, not constricted at the junction, cells with or without granules along the cross walls*L. hieormymusii* Lemm.

Cholorophyta (key to genera)

- 1-Thallus not filamentous, solitary or a colony of 2 or more cells.....2
 -Thallus a filament with or without branches11
 2- Thallus composed of solitary cell; bearing an unbranched hair with a basal sheath 1. *Chaetosphaeridium*
 -Thallus with solitary cells; without hair.....3
 3-Solitary cells, not adjoined to form colonies, not constricted or constricted in the midregion.....4
 - Solitary cells adjoined to form colonies..... 6
 4-Solitary cells constricted in the midregion, margin without spines..... 2. *Cosmarium*
 - Solitary cells, not constricted, crescent or sickle shaped, with pointed or narrowly rounded and tapering apices.....5
 5- Cell with 2 axial chloroplasts bearing longitudinal ridges, a chloroplast in either horn of the cell; pyrenoids conspicuous, usually in an axial row.....3. *Closterium*
 -Cell with one chloroplasts or with parietal chloroplasts not arranged as above, cells only slightly crescent-shaped, attached by a slender stipe to other algae ...4. *Characium*
 6-Colonies with definite patterns, definitely arranged and definite in number or forming, nets and plates.....7
 -Colonies with indefinite patterns, irregularly shaped; compounded by interconnecting strands of tough mucilage between clusters of cells.....5. *Botryococcus*

- 7- Cells cylindrical, 1 attached by 2 others at the end walls, repeatedly to form a network.6. *Hydrodidtyon*
 -Cells arranged otherwise, definite in number8
- 8- Colonies not bearing long or short needle-like spines, cell forming somewhat angular or oval plates9
 -Colonies bearing long or short spines10
- 9-Cells forming oval colony, trapezoidal, outer free walls deeply incised.....7. *Pediastrum*
 -Cells forming quadrate plates, the outer walls entire (not incised).....8. *Crucigenia*
- 10- Cells arranged side by side in one or two alternating rows; spines short, arising from the poles of the cells only.....9. *Scenedesmus*
 -Cells forming oval colony, bearing long needle-like spines10. *Golenkinia*
- 11- Filament without branches.....12
 - Filament with branches.....17
- 12- Chloroplasts extend only part way around the cell, filament not showing a basal-distal differentiation.....11. *Hormidium*
 - Chloroplasts are not about one-half the cell in length.....13
- 13- Chloroplast a parietal band or ring which encircles the cell or nearly so; apical cell not tapering.....12. *Ulothrix*
 - Chloroplast otherwise.....14
- 14- Chloroplasts spiraled, conjugation by the formation of tubes from one or both cells, either between cells of two different filaments (scalariform conjugation), or between adjacent cells in the same filament (lateral conjugation.....13. *Spirogyra*
 - Chloroplasts network, perforated, padded sheet, branched, beaded ribbon shape15
- 15- Chloroplast a perforated, padded sheet, branched, beaded ribbon; cell wall forms H-shaped pieces.....14. *Microspora*
 - Chloroplast a parietal network, axial plate or band shape16
- 16- Chloroplast an axial plate or band shape15. *Mougeotia*
 -Chloroplast a parietal network; cells cylindrical, with at least 1 ring-like scar at the anterior end just below the cross wall 16. *Oedogonium*

- 17- Multicellular hairs resulting from the apical tapering of branches, filaments of cells in 1 series throughout.....17. *Stigeoclonium*
 -Ends of branches not tapering to form hairs...18
- 18- Branching habit has been reduced; cells cylindrical.....18. *Rhizoclonium*
 - The habit of branching is variable according to habitat, branching open and spreading; cells cylindrical.....19. *Cladophora*

Cosmarium (key to species)

- Semicells ovate-pyramidal from a broad, flat base, basal angles rounded, sides convex, apex rather narrowly truncate or subtruncate, apical angles rounded. Chloroplasts axile, with two pyrenoids in each semicell..... *C. botrytis* Menegh.
 - Semicells compressed or rounded when seen from the top or side, not with radiating arms, margin of cell without spines.....
*C. panamenes* Presc.

Closterium (key to species)

- 1-Cells of 247-1000 μm long, 46.5-120 μm broad.....*C. lunula* (Mull).
 -Cells less than 1000 μm long,2
- 2-Inner margin tumid, cells of 221-370 μm long; inner margin tumid.....
*C. moniliferum* Ehr.
 -Inner margin not tumid; cells of 250-790 μm long, elongated, attenuated curved, rarely straight, with a single chloroplast in each semicell.....*C. acerosum* (Schr.).

Pediastrum (key to species)

- 1- Colony of upto 16 cells with two projections on margin2
 -Colony of more than 16 cells.....3
- 2-Colony 12-16 cells, 40 μm in diameter, projections side by side in front view of cell *P. boryanum* (Menegh).
 -Colony of 12-14 cells, 8-32 μm in diameter; cells covered with short, linear, concentrically arranged granules; 12-22 μm broad; 13-24 μm long..... *P. duplex* var. *rolundatum* Lucks.
- 3-Colony of 8-16 or 32-64 cells, with openings between cells. Cell wall smooth; cells 32 μm in diameter*P. biradiatum* Meyen.
 -Colony of 16-32 cells, 80-150 μm in diameter, oval to circular, with 8 interior and 4-6 straight sides; marginal cells with two lobes; cell wall covered with a

network of very fine ridges; cells 10-15 μm diameter .*P. sculptatum* G. M. Smith.

Scenedesmus (key to species)

- 1-Walls smooth, lateral ridges, terminal teeth or spines.....2
 -Wall with terminal teeth, or spines; 8-42 μm in diameter and 3-15 long, oblong-elliptical, arranged in one row or alternate.*S. quadricauda* Breb.
 2- Cells ovoid to cylindrical, with rounded ends; colony of 4 cells, 5-8 μm in diameter and 3-10 long, arranged in one row*S. bijuga* (TurP.)
 - Cells acicular to broadly elliptic.....3
 3- Colony with erect cells(2-4); with pointed ends; 5-30 μm in diameter and 2.5-10 long,*S. obliquus* (TurP.)
 -Colony with erect or unerect cells.....4
 4- Median cells erect, terminal cells lunate; 5-25 μm in diameter and 3-9 long, oblong-elliptical, arranged in one row.....*S. dimorphus* (TarP.)
 - Colony of 4-8 crescent- shaped cells; 5-20 μm in diameter and 2.5-11.....*S. incrassatulus* var. *mononae* G. M. Smith.

Spirogyra (key to species)

- Cells cylindrical, 40-85 μm in diameter, 115-200 μm long.....*S. rhizobrachalis* Jao.
 -Cells of 30-35 μm in diameter, 50-200 μm long; chloroplast single making up to 5.5 turns; end wall plane.....*S. borgeana* Transeau.

Cladophora (key to species)

- Attached or free floating; branch obliquely and rarely lateral; terminal cell 25-50 μm in diameter.....*C. glumerata* (L.) utz .
 -Free floating, lying loosely about the substratum; branch pseudochotomously, growing mainly intercalary; terminal cell 16-28 μm in diameter*C. fracta* (Muller ex Vahi)Kutz.

Bacillariophyta (key to genera)

- 1- Raphe or pseudoraphe absent ;valves with radial ornamentation 2
 - Raphe or pseudoraphe present; valves with transverse / longitudinal ornamentation4
 2- Frustules attached to form long chain or filament.....1. *Melosira*

- Frustules solitary but may form short chains.....3
 3-Valve margin with costae but central area not sharply distinct from margin; valves with radial rows of punctae in multiseriate rows, each row gradually or abruptly becoming uniseriate toward the center; marginal spines always present*Stephanodiscus*
 -Valve margin with costae and ornamentation in central area different from margin3. *Cyclotella*
 4-Raphe present on at least one valve; very short or rudimentary8
 - Raphe absent, pseudoraphe present on both valves..... 5
 5- Frustules with thick longitudinal septae, parallel to the valve face.....4. *Tabellaria*
 -Frustules without septae.....6
 6- Valves with thickened costae, symmetrical about the transapical plane.....5. *Diatoma*
 -Valves without thickened costae, symmetrical about the transapical plane.....7
 7-Frustules forming long ribbon-like chains, rarely solitary.....6. *Fragilaria*
 - Frustules solitary, or forming stellate colonies; striae often appearing as costae7. *Synedra*
 8- Raphe evident on at lease one valve9
 - Raphe not evident, concealed in a keel; frustules usually solitary or forming irregular ribbonlike chains or stellate colonies.....8. *Nitzschia*
 9- Valves not similarly ornamented, raphe on one valve, opposing valve with pseudoraphe, or with rudimentary raphe near valve10
 --Valves with similar ornamentation, raphe on both valves.....11
 10-Valves elliptical, never linear.9.*Cocconeis*
 -Valves elliptical-lanceolat.....10. *Encyonema*
 11- Valves elliptical.....12
 -Valves not elliptical, asymmetrical about the transapical plane, with completely developed raphe, opposing valve with rudimentary raphe near valve pole11. *Rhoicosphenia*
 12-Valves symmetrical about the transapical plane, asymmetrical about the apical plane.....13

- Valves asymmetrical about both the apical and apical plane14
- 13- Valve faces not parallel, both valve faces can be seen in girdle view.....12. *Amphora*
- Valve faces parallel.....13. *Cymbella*
- 14-Valves and raphe sigmoid.....14
- Valves and raphe not sigmoid, central area not extending to the margins of the valve, striae present along lateral margins of the central area.....14. *Navicula*
- 15 -Valves with transverse and longitudinal striae.....15. *Gyrosigma*
- Valves with transverse and oblique striae 16. *Pleurosigma*

***Melosira* (key to species)**

- Valves length less than 20 μm ;.....2
- Valves 9-13 μm long, no visible ornamentation.....*M. varians* C. A. Ag
- Valves 5-18 μm long.....*M. undulata* (Eh r.).

***Tabellaria* (key to species)**

- Cells 12-50 μm in diameter, 5-16 μm wide*T. flocculosa* (Roth) Kutz.
- Cells 30-140 long, 3-9 μm wide; intercalary bands per cell.....*T. fenestrata* (Lyngb.) Kuetz.

***Cymbella* (key to species)**

- Cells 20-70 μm long, 7-17 μm wide, ventral median striae ending in an isolated prominent puncta..... *C. affinis* Kützing.
- Cells 10-40 μm long, 2-12 μm wide.....*C. ventricosa* Kutz.

***Navicula* (key to species)**

- 1- cells upto 28 μm long, 5-8 μm wide; may be found in chains *N. confervacea* (Ehr.)
- Cell over 28 μm long.....2
- 2-Cell 20-40 μm long, 5-7 μm wide*N. cryptocephala* (Kutz).
- Cells 36-80 μm long, 6-10 μm wide ...*N. gracilis* (Ehr.).

Table 2. The taxonomic composition of taxa found in the studied are

Phylum	Family	Genus	Species	
Cyanophyta	1. Chamaesiphonaceae	1. <i>Chamaesiphon</i>	1. <i>C. incrustans</i> Grunow	
	2. Chroococcaceae	2. <i>Aphanocapsa</i>	2. <i>A. grevillei</i> (Hass.) Rabenh.	
		3. <i>Aphanothece</i>	3. <i>A. saxicola</i> Nag	
		4. <i>Chroococcus</i>	4. <i>C. limneticus</i> Lemm.	
		5. <i>Merismopedia</i>	5. <i>M. elegans</i> A. Braun	
		6. <i>Microcystis</i>	6. <i>M. aeruginosa</i> (Kutz.)	
		7. <i>Synechocystis</i>	7. <i>S. aquatilis</i> Sauv.	
	3. Oscillatoriaceae	8. <i>Lyngbya</i>	8. <i>L. martensiana</i> Menegh.	
			9. <i>L. hieormymusii</i> Lemm.	
			10. <i>L. aerugineo-coerulea</i> (Kutz.) Gom	
11. <i>O. granulata</i> Gardner				
12. <i>O. formosa</i> Bory				
13. <i>O. anguina</i> (Bory)				
14. <i>O. formosa</i> Bory				
15. <i>O. agardhii</i> (Gomont)				
16. <i>O. tenuis</i> Agardh				
17. <i>O. limnetica</i> Lemmermann.				
18. <i>O. sp.</i>				
19. <i>O. prolifica</i> N.				
20. <i>O. amoema</i> (Kützing) Gomont.				
	10. <i>Phormidium</i>	21. <i>P. retzii</i> N.		
		22. <i>P. ambiguum</i> Gom.		
4. Nostocaceae	11. <i>Anabaena</i>	23. <i>A. affinis</i> Lemm.		
	12. <i>Aphanizomenon</i>	24. <i>A. flos-aque</i> (L.) Ralfs.		
Chlorophyta	5. Ulothrichaceae	13. <i>Ulothrix</i>	25. <i>U. zonata</i> (Weber & Mohr) Kuetz.	
			26. <i>U. tenerrima</i>	
		14. <i>Hormidium</i>	27. <i>H.sp</i>	
	6. Chaetopeltidiae	15. <i>Chaetosphaeridium</i>	28. <i>C. globosum</i> (Nordst.)nKlebahn	
	7. Chaetophoraceae	16. <i>Stigeoclonium</i>	29. <i>St. flagelliferum</i> Kuetz	
	8. Cladophoraceae	17. <i>Cladophora</i>		30. <i>glumerata</i> (L.)Kutz
				31. <i>C. fracta</i> (Müll. ex Vahl.) Kützing
			18. <i>Rhizoclonium</i>	32. <i>R. hieroglyphicum</i> (Ag.) Kuetz
			19. <i>Oedogonium</i>	33. <i>O. sp</i>
	9. Oedogoniaceae	20. <i>Botryococcus</i>	34. <i>B. brounii</i> Kutz.	
10. Botryococcaceae	21. <i>Characium</i>	35. <i>C. subulatum</i> A. Br.		
11. Characieae	22. <i>Pediastrum</i>	36. <i>P. boryanum</i> Menegh.		
12. Hydrodictyaceae		37. <i>P. duplex var. rolundatum</i>		
		38. <i>P. biradiatum</i> Meyen.		
		39. <i>P. sculptatum</i> G. M. Smith		
		40. <i>H. reticulatum</i> (L.) Lagerheim		
	23. <i>Hydrodictyon</i>	41. <i>M. Loeigrenii</i> (Nordst.) Lag.		
13. Microsporaceae	24. <i>Microspora</i>			
14. Coelastraceae	25. <i>Coelastrum</i>	42. <i>C. acerosum</i> (Schr.)		
		43. <i>C. monilliferum</i> Ehr		
		44. <i>C. lunula</i> (Mull)		
		45. <i>C. cynthia</i> De Not.		
		46. <i>C. parvulum</i> Nag		
		47. <i>Cr. rectangularis</i> (A. Braun) Gay.		
15. Scenedesmaceae	26. <i>Crucigenia</i>	48. <i>S. bijuga</i> (Turp.)		
	27. <i>Senedesmus</i>	49. <i>S. dimorphus</i> (Tarp.)		
		50. <i>S. quadricauda</i> Breb.		
		51. <i>S. obliquus</i> (Turp.)		
		52. <i>S. incrassatus</i> var. <i>mononae</i> G. M. Smith.		
		53. <i>G. paucispina</i> W.G.S.		
16. cholorococcaceae	28. <i>Golenkinia</i>	54. <i>M. punctata</i>		
17. Zygnemataceae	29. <i>Mougeotia</i>			

Phylum	Family	Genus	Species
		30. <i>Spirogyra</i>	55. <i>S. rhizobrachalis</i> Jao. 56. <i>S. borgeana</i> Transeau
	18. Desmidiaceae	31. <i>Cosmarium</i>	57. <i>C. panamenes</i> Presc. 58. <i>C. botrytis</i> Menegh
Bacillariophyta	19. Coscinodiscaceae	32. <i>Stephanodiscus</i> 33. <i>Cyclotella</i>	59. <i>S. hantzchii</i> Grun 60. <i>C. comta</i> (Ehr.)
	20. Melosiraceae	34. <i>Melosira</i>	61. <i>M. granulata</i> (Ehr.) Ralfs 62. <i>M. varians</i> C. A. Ag
	21. Diatomaceae	35. <i>Diatoma</i>	63. <i>D. vulgare</i>
	22. Fragilariaceae	36. <i>Fragilaria</i> 37. <i>Synedra</i>	64. <i>F. capucina</i> Desm. 65. <i>S. ulna</i> (Nitsch.) Ehr.
	23. Tabellariaceae	38. <i>Tabellaria</i>	66. <i>T. flocculosa</i> (Roth) Kutz. 67. <i>T. fenestrata</i> (Lyngb.) Kuetz
	24. Cymbellaceae	39. <i>Cymbella</i>	68. <i>C. affinis</i> Kützig 69. <i>C. cistula</i> (Hemp.)
	25. Naviculaceae	40. <i>Navicula</i>	70. <i>N. gracilis</i> (Ehr.) 71. <i>N. cryptocephala</i> (Kutz.) 72. <i>N. confervoacea</i> (Ehr.)
	26. Cocconeidaceae	41. <i>Cocconeis</i> 42. <i>Encyonema</i>	73. <i>C. placentula</i> Ehrenb. 74. <i>E. caespitosum</i> Kütz
	27. Pleurosigmaaceae	43. <i>Gyrasigma</i> 44. <i>Pleurosigma</i>	75. <i>G. attenuatum</i> (Kutz.) 76. <i>G. scalproides</i> 77. <i>P. delicatum</i> .
	28. Catenulaceae	45. <i>Amphora</i>	78. <i>A. ovalis</i> (Kützing) Kützing
	29. Bacillariaceae	46. <i>Nitzschia</i>	79. <i>N. paradoxa</i> Gmelin.
	30. Diploneidaceae	47. <i>Rhoicosphenia</i>	80. <i>R. curvata</i>
Euglenophyta	31. Euglenaceae	48. <i>Phacus</i>	81. <i>P. curvicauda</i> Swir
Xanthophyta	32. Vaucheriaceae	49. <i>Vaucheria</i>	82. <i>V. geminata</i> (Vauch.)
Rhodophyta	33. Porphyridiaceae	50. <i>Porphyridium</i>	83. <i>P. cruentum</i> Naeg.
	34. Helminthocladia	51. <i>Chantransia</i>	84. <i>C. pygymaea</i> Kutz.

Table. 3 List of taxa recorded at shakhezr dome
(Abbreviations; S: spring; Su: summer; A: Autumn ; W: Winter)

phylum	Family	Genus	Species	S	SU	F	W
Cyanophyta	Chamaesiphonaceae	<i>Chamaesiphon</i>	<i>C. incrustans</i> Grunow			*	*
	Chroococaceae	<i>Aphanothece</i>	<i>A. saxicola</i> Nag		*	*	
	Chroococaceae	<i>Chroococcus</i>	<i>C. limneticus</i> Lemm		*		
	Chroococaceae	<i>Merismopedia</i>	<i>M. elegans</i> A. Braun		*		
	Chroococaceae	<i>Microcystis</i>	<i>M. aeruginosa</i> Kuetz.	*		*	*
	Oscillatoriaceae	<i>Lyngbya</i>	<i>L. hieronymusii</i> Lemm. <i>L. martensiana</i> Menegh.	*		*	*
	Oscillatoriaceae	<i>Oscillatoria</i>	<i>O. anguina</i> (Bory) <i>O. formosa</i> Bory <i>O. agardhii</i> (Gomont) <i>O. tenuis</i> Agardh <i>O. limnetica</i> Lemmermann. <i>O. sp.</i> <i>O. prolifica</i> N. <i>O. amoema</i> (Kützing) Gomont.	*	*	*	*
	Oscillatoriaceae	<i>Phormidium</i>	<i>Phormidium retzii</i> N. <i>Phormidium biguam</i> Gom.	*	*	*	*
	Nostocaceae	<i>Aphanizomenon</i>	<i>A. flos-aque</i> (L.) Ralfs	*			*
Chlorophyta	Ulothrichaceae	<i>Ulothrix</i>	<i>U. zonata</i> (W e b e r & Mohr) Kuetz.			*	*
	Chaetophoraceae	<i>Stigeoclonium</i>	<i>St. flagelliferum</i> Kuetz		*		
	Chroococaceae	<i>Golenkinia</i>	<i>G. paucispina</i> W. G.S. West	*	*	*	
	Cladophoraceae	<i>Cladophora</i>	<i>C. glomerata</i> (L.) Kutz <i>C. fracta</i> (Müll. ex Vahl.) Kützing		*	*	*
	Cladophoraceae	<i>Rhizoclonium</i>	<i>R. hieroglyphicum</i> (Ag.) Kuetz		*	*	

phylum	Family	Genus	Species	S	SU	F	W
	Oedogoniaceae	<i>Oedogonium</i>	<i>O. sp.</i>	*	*		*
	Botryococcaceae	<i>Botryococcus</i>	<i>B. brounii</i> Kutz.		*		
	Characiaceae	<i>Characium</i>	<i>C. subulatum</i> A. Br.		*	*	
	Hydrodictyaceae	<i>Pediastrum</i>	<i>P. boryanum</i> Menegh. <i>P. duplex</i> var. <i>rolundatum</i> <i>P. biradiatum</i> Meyen. <i>P. sculptatum</i> G. M. Smith		*	*	
	Hydrodictyaceae	<i>Hydrodictyon</i>	<i>H. reticulatum</i> (L.) Lagerheim		*	*	
	Microsporaceae	<i>Microspora</i>	<i>M. Loeigrenii</i> (Nordst.) Lag.		*	*	
	Coelastraceae	<i>Coelastrum</i>	<i>C. acerosum</i> (Schr.) <i>C. moniliferum</i> Ehr <i>C. lunula</i> (Mull)		*	*	*
	Scenedesmaceae	<i>Crucigenia</i>	<i>Cr. rectangularis</i> (A. Braun) Gay.			*	
	Scenedesmaceae	<i>Scenedesmus</i>	<i>S. bijuga</i> (Turp.) <i>S. dimorphus</i> (Tarp.) <i>S. quadricauda</i> Breb. <i>S. obliquus</i> (Turp.) <i>S. incrassatulus</i> var. <i>mononae</i> G. M. Smith.		*	*	*
	Zygnemataceae	<i>Mougeotia</i>	<i>M. punctata</i> Wittrock			*	*
	Zygnemataceae	<i>Spirogyra</i>	<i>S. rhizobranchialis</i> Jao. <i>S. borgeana</i> Transeau		*	*	*
	Desmidiaceae	<i>Cosmarium</i>	<i>C. panamenes</i> Presc. <i>C. botrytis</i> Menegh		*	*	
	Coscinodiscaceae	<i>Stephanodiscus</i> <i>Cyclotella</i>	<i>S. hantzchii</i> Grun <i>C. comta</i> (Ehr.)		*	*	
Bacillariophyta	Melosiraceae	<i>Melosira</i>	<i>M. granulata</i> (Ehr.)Ralfs <i>M. varians</i> C. A.Ag	*	*	*	*
	Diatomaceae	<i>Diatoma</i>	<i>D. vulgare</i>			*	*
	Fragilariaceae	<i>Fragilaria</i>	<i>F. capucina</i> Desm.		*	*	
	Fragilariaceae	<i>Synedra</i>	<i>S. ulna</i> (Nitsch.) Ehr.		*	*	*
	Tabellariaceae	<i>Tabellaria</i>	<i>T. flocculosa</i> (Roth)Kutz. <i>T. fenestrata</i> (Lyngb.) Kuetz		*	*	*
	Cymbellaceae	<i>Cymbella</i>	<i>C. affinis</i> <i>C. cistula</i> (Hemp.)			*	*
	Naviculaceae	<i>Navicula</i>	<i>N. gracilis</i> (Ehr.) <i>N. cryptocephala</i> (Kutz.) <i>N. confervacea</i> (Ehr.)	*	*	*	*
	Cocconeidaceae	<i>Cocconeis</i>	<i>C. placentula</i> Ehrenb.			*	
	Pleurosigmataceae	<i>Gyrasigma</i>	<i>G. attenuatum</i> (Kutz.) <i>G. scalproides</i> (Rabenhorst) Cleve	*	*	*	
	Catenulaceae	<i>Amphora</i>	<i>A. coffeaeformis</i> (C.Agardh) Kützing		*	*	*
	Bacillariaceae	<i>Nitzschia</i>	<i>N. paradoxa</i> Gmelin.			*	
	Diploneidaceae	<i>Rhoicosphenia</i>	<i>R. curvata</i> (Kützing) Grunow	*			
Xanthophyta	Vaucheriaceae	<i>Vaucheria</i>	<i>V. geminata</i> (Vauch.)	*	*	*	*
Euglenophyta	Euglenaceae	<i>Phacus</i>	<i>P. caudatus</i> Hubner.		*		

Table 4 List of taxa recorded at Gashtroodkhan
(Abbreviations; S: spring; Su: summer; A: Autumn ; W: Winter)

phylum	Family	Genus	Species	S	SU	F	W
Cyanophyta	Chamaesiphonaceae	<i>Chamaesiphon</i>	<i>C. incrustans</i> Grunow		*	*	
	Chroococcaceae	<i>Aphanocapsa</i>	<i>A. Grevillei</i> (Hass.) Rabenh.		*		
	Chroococcaceae	<i>Aphanothece</i>	<i>A. saxicola</i> Nag		*		*
	Chroococcaceae	<i>Chroococcus</i>	<i>C. limneticus</i> Lemm.		*		
	Chroococcaceae	<i>Microcystis</i>	<i>M. aeruginosa</i> (Kutz.)		*	*	

phylum	Family	Genus	Species	S	SU	F	W
	Chroococaceae	<i>Synechocystis</i>	<i>S. aquatilis</i> Sauv.		*		
	Oscillatoriaceae	<i>Lyngbya</i>	<i>L. martensiana</i> Menegh. <i>L. hieormymusii</i> Lemm. <i>L. aerugineo-coerulea</i> (Kutz.)	*	*	*	
	Oscillatoriaceae	<i>Oscillatoria</i>	<i>O. granulata</i> Gardner <i>O. formosa</i> Bory	*	*	*	
	Oscillatoriaceae	<i>Phormidium</i>	<i>P. retzii</i> N.		*		
	<u>Nostocaceae</u>	<i>Aphanizomenon</i>	<i>A. flos-aque</i> (L.) Ralfs.	*	*	*	
Chlorophyta	Chaetophoraceae	<i>Stigeoclonium</i>	<i>St. farctum</i> Berthold	*		*	
	Cladophoraceae	<i>Cladophora</i>	<i>C. glumerata</i> <i>C. fracta</i> (O.F.Müll. ex Vahl.) Kützing	*		*	
	Characieae	<i>Characium</i>	<i>C. Subulatum</i> A. Br.				*
	Chlorococaceae	<i>Golenkinia</i>	<i>G. paucispina</i> W.G.S.	*	*	*	
	Zygnemataceae	<i>Spirogyra</i>	<i>S. rhizobrachalis</i> Jao. <i>S. borgeana</i> Transeau	*	*	*	
	Scenedesmaceae	<i>Closterium</i>	<i>C. acerosum</i> (Schr.) <i>C. moniliferum</i> Ehr <i>C. lunula</i> (Mull)	*	*	*	
Bacillariophyta	Melosiraceae	<i>Melosira</i>	<i>M. varians</i> C. A.Ag <i>M. granulata</i> (Ehr.)Ralfs	*	*	*	
	Fragilariaceae	<i>Synedra</i>	<i>S. ulna</i> (Nitsch.) Ehr.	*	*		
	Cymbellaceae	<i>Amphora</i>	<i>A. ovalis</i> (Kützing) Kützing		*		
	Cocconeidaceae	<i>Encyonema</i>	<i>E. caespitosum</i> (Kützing) Brun				*
	Naviculaceae	<i>Navicula</i>	<i>N. cryptocephala</i> (Kutz.)	*	*	*	*
	Tabellariaceae	<i>Tabellaria</i>	<i>T. fenestrata</i> (Lyngb.) Kuetz. <i>T. flocculosa</i> (Roth) Kützing	*	*	*	*
	Diatomaceae	<i>Diatoma</i>	<i>D. vulgare</i> Bory de Saint-Vincent	*	*		
Xanthophyta	Vaucheriaceae	<i>Vaucheria</i>	<i>V. sessilis</i> (Vauch.)	*			
Rhodophyta	Porphyridiaceae	<i>Porphyridium</i>	<i>P. cruentum</i> Naeg.	*			

Table. 5 List of taxa recorded at Talab laleh
(Abbreviations; S: spring; Su: summer; A: Autumn ; W: Winter)

phylum	Family	Genus	Species	S	SU	F	W
Cyanophyta	Chroococaceae	<i>Microcystis</i>	<i>M. aeruginosa</i> Kuetz. emend Elenkin	*		*	*
	<u>Nostocaceae</u>	<i>Anabaena</i>	<i>A. affinis</i> Lemm.	*			
	Oscillatoriaceae	<i>Lyngbya</i>	<i>L. martensiana</i> Menegh.	*		*	
Chlorophyceae	Ulothrichaceae	<i>Ulothrix</i>	<i>U. zonata</i> (Weber & Mohr) Kuetz.	*		*	*
	Ulothrichaceae		<i>U. tenerrima</i>		*		
		<i>Hormidium</i>	<i>H. sp.</i>				
	Chaetopeltidiaceae	<i>Chaetosphaeridium</i>	<i>C. globosum</i> (Nordst.) Klebahn		*	*	*
	Chaetophoraceae	<i>Stigeoclonium</i>	<i>St. flagelliferum</i> Kuetz		*	*	
	Cladophoraceae	<i>Cladophora</i>	<i>C. glumerata</i>		*	*	*
	Cladophoraceae	<i>Rhizoclonium</i>	<i>R. hieroglyphicum</i> (Ag.) Kuetz.		*	*	*
	Microsporaceae	<i>Microspora</i>	<i>M. sp</i>			*	

phylum	Family	Genus	Species	S	SU	F	W
	Oedogoniaceae	<i>Oedogonium</i>	<i>O. sp</i>		*	*	
	Botryococcaceae	<i>Botryococcus</i>	<i>B. branii</i> Kützing,	*	*		
	Coelastraceae	<i>Coelastrum</i>	<i>C. acerosum</i> (Schr.) <i>C. cynthia</i> De Not. <i>C. parvulum</i> Nag.	*	*		
	Zygnemataceae	<i>Mougeotia</i>	<i>M. genuflexa</i> (Dill w.) C. A. Agardh,			*	*
	Zygnemataceae	<i>Spirogyra</i>	<i>S. rhizobrachalis</i> Jao.	*	*	*	
	Desmidiaceae	<i>Cosmarium</i>	<i>C. panamenes</i> Presc.	*	*		
Bacillariophyta	Melosiraceae	<i>Melosira</i>	<i>M. granulata</i> (Hhr.)	*	*	*	*
	Fragilariaceae	<i>Synedra</i>	<i>S. ulna</i> (Nitsch.) Ehr.	*	*		
	Naviculaceae	<i>Navicula</i>	<i>N. cryptocephala</i> (Kutz.)	*		*	
	Pleurosigmataceae	<i>Gyrosigma</i>	<i>G. attenuatum</i> (Kutz)	*			
	Pleurosigmataceae	<i>Pleurosigma</i>	<i>P. delicatum</i> Smith		*		
Rhodophyta	Helminthocladia	<i>Chantrasia</i>	<i>C. pygymaea</i> Kutz.			*	
Euglenophyta	Euglenaceae	<i>phucus</i>	<i>P. curvicauda</i> Swir		*		

REFERENCES

- Abbasi, K., Moraadkha, S. and Sarpanah, A (2006) Identification and distribution of fish fauna in Siahdarvishan River (Anzali Wetland basin). *Pajouhesh and Sazandegi Journal*. 74: 27-39.
- Adams, V.D (2000) *Water and waste water examination manual*. Department of the Environment. pp. 260.
- Addy, K. and Green, L (1966) *Algae in Aquatic Ecosystems*. University of Rhode Island College of Resource Development Department of Natural Resources Science. Cooperative Extension. Fact Sheet .96: 1-4.
- Afsharzadeh, S (2003) *Study of algal flora in Zayandeh Rood River*. Ph. D. Thesis University of Isfahan. pp. 195 (In Persian).
- Apha (1986) *Standard method for the examination of water and wastewater* (17th ed.). American Public Health Association, Washington. pp.1265.
- Altieri, A.H. , Silliman, B.R. and Bertness, M.B (2007) Hierarchical organization via a facilitation cascade in intertidal cord grass bed communities . *American naturalist*. 169: 198-206.
- Andrejic, J.Z. , Krizmanic, J. and Cvijan, M (2012) Diatom species composition of the Nisava River an its tributaries Jerma and Temska Rivers (Southern Serbia). *Archives of Biological Science Belgrade*. 64 : 1127-1140.
- Anderson, J.B (1966) *A Guide to the Common Diatoms at Water Pollution Surveillance System Stations*. U.S. Department of Interior Federal Water Pollution Control Administration. pp.1-98.
- Çolak, Ö. and Kaya, Z (1988) Alglerin atik suların biyolojik arıtılmasında kullanılma olanakları, *Doğa Bilimleri Dergisi*. 12: 18-29.
- Compere, P (1981) Algues des desers d, Iran. *Bulletin du Jardin botanique national de Belgique*, Belgium. 51: 3-40.
- Devika, R. , Rajendran, A. and Selvapathy, P (2006) Variation studies on the physico-chemical and biological characteristics at different depths in model waste stabilisation tank. *Pollution Research* . 24: 771-774.
- Dayton, P.K (1972) Toward an understanding of community resilience and the potential effects of enrichment to the benthos at McMurdo Sound, Antarctic (ed. B.C. Parker). Proceedings of the colloquium on conservation problems in Antarctica. Allen Press, Lawrence, Kansas, USA. pp. 81-96
- El-Awamri, A., El-Salam, A., Shaaban M. and Saleh A (2007) Floristic Study on Benthic Diatoms of the Groundwater Seepages at Kobri El-kobba (Cairo, Egypt). *Journal of Applied Sciences Research*. 3: 1809-1818.
- Ersanli, E. and Gönülol, A. I (2003) Study on the Phytoplankton and Seasonal Variation of Lake Simenit (Terme - Samsun, Turkey) , *Turkish Journal of Fisheries and Aquatic Sciences*.3: 29-39.
- Ezekiel, E.N., Ogamba, E.N. and Abowei, J.F.N (2011) The Distribution and Seasonality of Phytoplankton in Sombreiro River, Niger Delta, Nigeria *Asian Journal . of Agriculture Science*. 3 : 192-199.
- Fumanti, B., Alfinito, S. and Cavacini, P (1995) Dipartimento di Biologia Vegetale,

- Universitd di Roma 'La Sapienza', Floristic studies on freshwater algae of Lake Gondwana, Northern Victoria Land (Antarctica) *Hydrobiologia*. 316 : 81-90.
- Gomont, M (1972) *Monographie des Oscillariees (Nostocacees Homocysteas, Deuxieme partie-Lyngbyees*. Edi: Weinheim, J. Cramer. 92-256.
- Hirano, M (1973) Freshwater algae from Mesopotamia. *Control Biology Laboratory, Kyoto. Univ.* 24:105-119.
- Izaguirre, I., O'Farrell, I and Tell, G (2001) Variation in phytoplankton composition and limnological features in a water-water ecotone of Lower Parana Basin (Argentina) . *Freshwater Biology*. 46: 63-74.
- Kim, Y. S., Choi, H. G. and Nam K. M (2008) Seasonal variations of marine algal in the community vicinity of Uljin nuclear power plant, Korea *Journal of Environmental Biology*. 29: 493-499.
- Kiyanmehr, H (1994) *Identification of fresh water algae of Mashhad provinces*, Ferdosi University.
- Kolayli, S., Baysal, A. and Shahin, B (1998) Study on the Epipellic and Epilithic Algae of Sana River (Trabzon/Turkey) , *Turkish Journal of Botany*, **22**, 163-170
- Komulainen, S (2009) Diatoms of periphyton assemblages in small rivers in Northwestern Russia, *Studi trentini di scienze naturali*. 84: 153-160.
- Löffler, H (1961) Beitrage zur Kenntnis der Iranischen Binnengewasser. 2ed Int. Rev. Ges. *Hydrobiol.* 46: 309-406.
- Maosen, H (1983) *Illustration of fresh water plankton*. Agriculture press. pp.1-44.
- Moghadam, F (1976) Diatoms as indicator of pollution Zayandeh Rive, Iran. -Proceeding of the Academy of Natural Science of Philadelphia. 127: 281- 297
- Nejadsattari, T., Noroozi, M. and Fallahi M (2005) The composition and seasonal distribution of epiphytic algae in Anzali lagoon, Iran. *Cryptogamie. Algal*. 26: 387-398.
- Noroozi, M., Naqinezhad, A. and Mehrvarz, S. Sh (2009) Algal flora in first Iranian land-marine the Boujagh National Park. *International Journal on Algae*. 11:276-288.
- Oguni, A., Tanaka, Y., Uemura, M., Yamamoto, M. and Takahashi, E (1987) Floristic Studies on Algae from inland waters of Ongul islands and Vicinity, Antarctica: East Ongul island Proc. NIPR Symp. *Polar Biology*. 1: 233-254.
- Patrick, R and Reimer, C. W (1966). *The Diatoms of the United States V-I & 2*. The Academy of Natural Science of Philadelphia. USA.
- Pfiester, L. A., Lynch, R. and Wright, T.L (1980) Species composition and diversity of Phytoplankton in the Grand River Dam area ,Oklahoma. Proceedings of the Oklahoma Academy of Science. 60:63-68.
- Pundhir, P. and Rana, K.S (2002) Population dynamics of phytoplankton in the wetland area of Keoladeo National Park, Bharatpur (Rajasthan). *Ecology Environment and Conservation*. 8: 253-255.
- Prescott, G.W. (1945) *How to know fresh water algae*. W.M. G Brown Company Publishers. pp. 1-217.
- Rajagopal, T., Thangamani, A. and Archunan G (2010) Comparison of physico-chemical parameters and phytoplankton species diversity of two perennial ponds in Sattur area, Tamil Nadu *Journal of Environmental Biology*. 31: 787-794.
- Richardson, T.L., Gibson, C.E. and Heaney, S.I (2000) Temperature, growth and seasonal succession of Phytoplankton in Lake Baikal, Siberia. *Freshwater Biology* . 44: 431-440.
- Sahin, B and Akar, B (2004) Species Composition and Diversity of Epipellic Algae in CatalLake(Şebinkarahisar-Giresun,Turkey) ,*Turkish Journal of Biology*. 28 : 103-109.
- Sen, B. and Sonmez, F (2006) A Study on the Algae in Fish Ponds and Their Seasonal Variations. *International Journal of Science and Technology*. 1: 25-33.
- Škaloud, P (2009) Species composition and diversity of aero-terrestrial algae and cyanobacteria of the Boreč Hill ventaroles. *Fottea*. 9 : 65-80.
- Smith, G.M (1920) *Phytoplankton of the Inland Lakes of Wisconsin Part I Myxophyceae, Phaeophyceae, Heterokontae, and Chlorophyceae exclusive of the Desmidiaceae*. Published by The state. pp.1-243.
- Smith, G.M (1924) *Phytoplankton of the Inland Lakes of Wisconsin Part II Desmidiaceae* Madison. Published by the University. pp.1-227.
- Starmach, K.(1972) *Flora Slodkowodna Polski*, Freshwater Biological Association, Windermere.UK. 10: 227-263.
- Thebault, J.M. and Rabouille, S (2003) Comparison between two mathematical formulation of the phytoplankton specific growth rate as a function of light and temperature, in

- two simulation models (Aster and YoYo). *Ecological Modelling*. 163: 145-151.
- Tyor , A. K. and Chawla, D (2012) Survey and study of Phytoplankton ecology in Sukhna Lake, Ghandigarh, (India). *International Journal of Applied Biology and Pharmaceutical Technology*. 3: 229-238.
- West, G.S. (1904) *Cambridge biological series, The British freshwater algae*. Cambridge University Press. pp. 372
- Wu, J.T. and Chou, J.W (1999) Dinoflagellate associations in feitsu ireservoir. Taiwan. *Botanical Bull. Academia Sinica*. 39: 137-145.
- Zarei-Darki, B (2009) Marine Species in the Algal Flora of the Anzali Swamp (Iran). *Russian Journal of Marine Biology*.35: 200-205.
- Zimba, P.V. and Hopson, M.S (1997) Quantification of epiphyte removal efficiency from submersed aquatic plants. *Aquatic Botany*. 58: 173-179.

مطالعه فلوریستیک جلبک های رودخانه سیاه درویشان ، استان گیلان، شمال ایران م. ب. فقیر* و ش. شفیع

(تاریخ دریافت: ۹۰/۶/۲۸ - تاریخ پذیرش: ۹۰/۱۱/۱۷)

در تحقیق حاضر جلبک های آب شیرین رودخانه سیاه درویشان طی یک سال (۱۳۸۵ - ۱۳۸۴) در ماه های متوالی نمونه برداری شد. سه ایستگاه نمونه برداری (روستای گشت رودخان، سد شاخزر، تلاب لاله) مشخص شدند. برخی از پارامترهای فیزیکی و شیمیایی (دمای آب، سختی کل (برحسب درجه آلمانی d°)، میزان نیترات، نیتریت، و فسفات و pH) اندازه گیری شد. در مجموع ۳۴ تیره، ۵۱ جنس و ۸۴ گونه از جلبک های آب شیرین مورد شناسایی قرار گرفت که شامل ۳۶ گونه از Cholorophta، ۲۴ گونه از Cyanophyta، ۲۰ گونه از Rhodophyta، با ۲ گونه از Euglenophyta و Xanthophyta هر کدام با ۱ گونه می باشد. از میان تیره ها، Oscillatoriaceae با ۱۴ گونه بیشترین غنای گونه ای را دارد. به منظور سهولت شناسایی فلور جلبکی منطقه، کلید های تاکسونومیکی در سطح جنس و گونه ارائه شده است.

*مؤلف مسئول