

[Research]

## Effect of dietary estradiol-17 $\beta$ on growth performance, body composition and blood indices in Stellate sturgeon, *Acipenser stellatus*

H. Khara<sup>1\*</sup>, B. Meknatkhah<sup>2</sup>, B. Falahatkar<sup>3</sup>, M. Ahmadnezhad<sup>4</sup>, S. Poursaeid<sup>5</sup>

1- Dept. of Fisheries, Lahijan Branch, Islamic Azad University, Lahijan, Iran

2- Dr. Yousefpour Fish Hatchery Center, Siahkal, Iran

3-Dept. of Fisheries, Faculty of Natural Resources, University of Guilan, Sowme Sara, Iran

4- Inland Waters Aquaculture Institute, Bandar Anzali, Iran

5- Young Researchers and Elite Club, Rasht Branch, Islamic Azad University, Rasht, Iran

\* Corresponding author's E-mail: h.khara1974@yahoo.com

(Received: Feb. 25. 2014, Accepted: Jul. 05. 2014)

### ABSTRACT

This study was investigated the effects of dietary estradiol-17 $\beta$  (E2) on growth, body composition and blood indices in *Acipenser stellatus*. Fish (40.9  $\pm$  1.1 g average initial weight; n = 60 per group) were fed with three different diets containing 0 (control), 25 and 50 mg kg<sup>-1</sup> dietary estradiol contents to apparent satiation for seven months. The results suggested that growth rate were decreased as the E<sub>2</sub> level was increased. No significant difference was observed in condition factor among dietary treatments. The highest survival rate was observed in fish fed control diet, but was not significantly different among the treatments (P>0.05). Body composition did not show significant changes among dietary treatments. Number of white blood cells and red blood cells, hemoglobin and hematocrit values were significantly decreased as the E<sub>2</sub> levels were raised (P<0.05). Plasma biochemical parameters including glucose, total protein, cholesterol, triglyceride, calcium and phosphorus levels were dose dependent with the lowest levels in control with the highest levels in fish treated with 50 mg E<sub>2</sub>. The results of the present study revealed growth suppression of dietary E<sub>2</sub> and changes of blood indices with providing some basic information on the effect of estrogen hormone on physiology of sturgeon.

**Keywords:** Estradiol, Blood indices, body composition, Growth, Stellate sturgeon, *Acipenser stellatus*

### INTRODUCTION

In general, the principle that dictates which type of hormone to be used is the need to supplement or replace the particular hormone type that is deficient in the animals to be treated. The estradiol-17 $\beta$  (E<sub>2</sub>) is one of the most important compounds which commonly used for anabolic purposes in food animals. Gender and maturity of an animal influence its growth rate and body composition. E<sub>2</sub>, on the other hand, may act by stimulation of the somatotrophic axis to increase growth hormone (Degani, 1986). In theory, growth enhancement can be able to occur through improvement in appetite, digestion, absorption of metabolic re-arrangements leading to overall anabolism, and the contribution of each of these factors varies somewhat among different species and type of steroid used (Woo *et al.*, 1993).

E<sub>2</sub> promoted growth in yellow perch *Perca flavescens* by stimulating appetite (Malison *et al.*, 1988). On the other hand, E<sub>2</sub> has been proven to depress growth in many fish species (Johnstone *et al.*, 1978; Blázquez *et al.*, 1998). Akhundov and Fedorov (1994) examined the effect of estradiol dipropionate on ovarian development in Sterlet, *Acipenser ruthenus*. Juvenile paddlefish, *Polyodon spathula*, implanted with 17-methyltestosterone showed changes in sex ratio (Shelton and Mims 1998). Also, dietary E<sub>2</sub> feminized all shortnose sturgeon, *Acipenser brevirostrum*, after a 9-month trial (Flynn & Benfey 2007). Sturgeons are valuable species, which are currently highly endangered. The Caspian populations are under massive pressure from overfishing (including poaching) and loss of spawning sites and the stocks are declining very fast. The rearing of these

species has seen considerable progress during past two decades. Stellate sturgeon (*Acipenser stellatus*) is one of the Caspian sturgeon species with relatively suitable growth rates and appears to be very suitable for aquaculture (Falahatkar *et al.*, 2014). Despite the economic importance of Stellate sturgeon, there is little information on using of E2 supplementation in fish diet and its effect on physiological functions. The present study, for the first time, assesses the effectiveness of E2 supplementation to growth performance, body composition and blood indices in Stellate sturgeon.

## MATERIALS AND METHODS

### Experimental fish

One hundred and eighty, five-month old, juvenile Stellate sturgeon reared at Dr. Yousefpour Fish Hatchery Center in Siahkal, Guilan, Iran, were used for this study. Twenty fish within average weight of  $40.9 \pm 1.1$  g were randomly distributed into each of nine circular concrete tanks containing a water volume of 900 L and supplied with flow-through freshwater at  $17 \pm 0.5$  L/min. During the experimental period, water was aerated continuously with compressed air delivered through an air stone in each tank. The fish were kept and reared under this condition for approximately seven months. Water parameters were recorded daily; average water temperature and dissolved oxygen concentrations were  $18 \pm 2$  °C and  $8.7 \pm 1.5$  mg/L.

### Diet preparation

E2 (Sigma- Aldrich St. Louis, MO, USA) was dissolved in alcohol, sprayed on the feed, and then stored at 4 °C in a refrigerator. The experiments were divided into three group treatments with different concentrations of dietary E2: 0 (control treatment), 25, and 50 mg/kg (Lord *et al.*, 2009). The fish was kept under natural photoperiod and fed 2-6 times a day depending on water temperature according to their satiation with a formulated diet (Biomar, France). Every day, each tank was partially cleaned of the fish feces and the water was partially changed (about 50%). Fish were weighed every three weeks intervals during the experimental period to monitor the growth performance.

### Growth performance

Body weight (g) and total body length (cm) of individual fish were measured to the nearest of 1 g and 0.1 cm, respectively, every three weeks. The individual fish from each tank was anesthetized by 300 mg l<sup>-1</sup> of clove powder and then weighted. Fish were deprived of food for 24 h before each weighing. At the end of the trial, specific growth rate (SGR), body weight increase (BWI), weight gain (WG), feed conversion ratio (FCR) and condition factor (CF) were calculated using the following formulas to compare growth performance among experimental groups (Biswas *et al.*, 2008):

$SGR (\% \text{ day}^{-1}) = 100 \times (\ln \text{ final body weight} - \ln \text{ initial body weight}) / \text{time (days)}$

$BWI (\%) = 100 \times [\text{final body weight (g)} - \text{initial body weight (g)}] / \text{initial body weight (g)}$

$WG \text{ (g)} = \text{final body weight (g)} - \text{initial body weight (g)}$

$FCR = \text{feed intake (g)} / \text{wet weight gain (g)}$

$CF = 100 \times [\text{final body weight (g)} / \text{total body length (cm)}^3]$

### Chemical analysis

At the end of the experiment, five fish were randomly sampled from each replicate and then pooled in plastic bags and stored at -20 °C for whole body composition analysis. The proximate body composition (moisture, crude protein, crude lipid, and ash) was determined using the standard methods of the Association of Official Analytical Chemists (AOAC, 1995). Protein was determined by estimating the Kjeldahl method ( $N \times 6.25$ ), moisture by heating at 105 °C to constant weight, ash by incinerating in a crucible at 600 °C and crude lipid was determined by using Soxhlet apparatus.

### Blood sampling

At the end of the trial, five fish from each tank were quickly captured and sampled after being anesthetized. Then, approximately 2 mL of blood samples were collected from the caudal vasculature using a 5 mL heparinized syringe equipped with a 25 gauge needle. Blood samples were divided into two portions. Half of the blood was used for separating plasma and the remaining blood was used for hematological analysis. Plasma was

separated by centrifugation ( $1500 \times g$  for 10 min) and stored at  $-70^\circ\text{C}$  until subsequent analysis. Hematocrit (Hct) value was determined by the standard microhematocrit method and expressed in percentage. Hemoglobin (Hb) concentration was determined using the cyanmethemoglobin method. The numbers of white blood cells (WBC), red blood cells (RBC), mean corpuscular hemoglobin (MCH), mean corpuscular volume (MCV) and mean corpuscular hemoglobin concentration (MCHC) were calculated according to Ranzani-Paiva *et al.* (2004). Plasma glucose was analyzed based on the colorimetric glucose oxidase-peroxidase reaction (Bayunova *et al.*, 2002). Plasma cholesterol and triglyceride levels were determined by the CHOD-PAP and GPO-PAP methods, respectively (Chatzifotis *et al.*, 2004), using commercial available kits (Pars Azmun, Karaj, Iran). Total protein levels were measured by the Biuret method (Ziest Chem Diagnostics, Tehran, Iran). Total plasma calcium was determined by the colorimetric method (Sigma-Aldrich procedure no. 587). Plasma phosphorus was measured using an endpoint colorimetric assay (Sigma-Aldrich procedure no. 360).

### Statistical analysis

All data were subjected to a one-way analysis of variance (ANOVA) after confirmation of normality of data and homogeneity of variance based on Kolmogorov-Smirnov and Levene's tests, respectively. Significance of the differences between means was tested using Duncan's multiple range test ( $P < 0.05$ ). All assays were performed in triplicates and data are shown as mean  $\pm$  standard deviation (SD) for each dietary group.

### RESULTS

The final body weight of the groups fed 25 and 50 mg E<sub>2</sub> kg<sup>-1</sup> diet were significantly lower than those of control group ( $P > 0.05$ ). The FCR and SGR of the control fish were significantly higher ( $P < 0.05$ ) than those of all E<sub>2</sub> treated groups (Table 1). There was no significant difference in survival rate and CF among the dietary treatments. Different concentrations of E<sub>2</sub> in the diets significantly affected blood indices. Highest Hb and Hct were observed in control group compared to groups treated with E<sub>2</sub> (Table 2). Levels of calcium, cholesterol, triglyceride, protein and phosphorus were significantly increased as the E<sub>2</sub> levels were raised (Table 3). In all experimental groups, body composition including crude protein, crude lipid, moisture and ash content did not show significant differences (Table 4).

**Table 1.** Growth performance of Stellate sturgeon fed with different levels of E<sub>2</sub> in diet.

E2 levels (mg kg <sup>-1</sup> )	Initial body weight (g)	Final body weight (g)	FCR	SGR (%/day)	CF	Survival rate (%)
Control	40.9 $\pm$ 1.1	171.7 $\pm$ 6.5 <sup>a</sup>	1.15 $\pm$ 0.04 <sup>b</sup>	1.38 $\pm$ 0.02 <sup>a</sup>	0.24 $\pm$ 0.001	100
25	40.9 $\pm$ 1.1	119.4 $\pm$ 3 <sup>b</sup>	1.46 $\pm$ 0.07 <sup>b</sup>	1.2 $\pm$ 0.02 <sup>b</sup>	0.26 $\pm$ 0.002	95 $\pm$ 5
50	40.9 $\pm$ 1.1	80.6 $\pm$ 1.4 <sup>c</sup>	2.01 $\pm$ 0.18 <sup>a</sup>	1.03 $\pm$ 0.04 <sup>c</sup>	0.25 $\pm$ 0.007	81.7 $\pm$ 7.3

Different superscripts in columns show the significant differences between data ( $P < 0.05$ ).

**Table 2.** Blood indices in Stellate sturgeon fed with different levels of E<sub>2</sub> in diet.

Parameters	Control	25 mg kg <sup>-1</sup>	50 mg kg <sup>-1</sup>
RBC ( $\times 10^3$ iL <sup>-1</sup> )	525.3 $\pm$ 43.7 <sup>a</sup>	435.8 $\pm$ 23.2 <sup>a</sup>	304.4 $\pm$ 22.6 <sup>b</sup>
WBC ( $\times 10^3$ iL <sup>-1</sup> )	13.6 $\pm$ 0.86 <sup>a</sup>	12.8 $\pm$ 0.68 <sup>a</sup>	7.9 $\pm$ 0.78 <sup>b</sup>
Hemoglobin (mg dL <sup>-1</sup> )	6.55 $\pm$ 0.46 <sup>a</sup>	5.54 $\pm$ 0.28 <sup>a</sup>	3.8 $\pm$ 0.3 <sup>b</sup>
Hematocrit (%)	26.6 $\pm$ 1.7 <sup>a</sup>	21.1 $\pm$ 1 <sup>a</sup>	15.4 $\pm$ 1.2 <sup>b</sup>
MCV (fL)	517 $\pm$ 11	511 $\pm$ 8	538 $\pm$ 7
MCH (pg)	12.6 $\pm$ 0.22	12.7 $\pm$ 0.11	12.4 $\pm$ 0.21
MCHC (g dL <sup>-1</sup> )	24.5 $\pm$ 0.34	25 $\pm$ 0.34	24.8 $\pm$ 0.40
Lymphocyte (%)	77.7 $\pm$ 2.1	78.7 $\pm$ 2.2	79.1 $\pm$ 1.9
Monocyte (%)	1.9 $\pm$ 0.3	2.1 $\pm$ 0.3	1.9 $\pm$ 0.3
Neutrophil (%)	17.8 $\pm$ 1.3	17.5 $\pm$ 1.7	18.7 $\pm$ 1.7
Eosonophil (%)	2.6 $\pm$ 0.9	1.7 $\pm$ 0.5	1 $\pm$ 0.2

Different superscripts in columns show the significant differences between data ( $P < 0.05$ ).

**Table 3.** Blood biochemistry parameters in Stellate sturgeon fed with different levels of E<sub>2</sub> in diet.

E2 levels (mg kg <sup>-1</sup> )	Calcium (mg dL <sup>-1</sup> )	Glucose (mg dL <sup>-1</sup> )	Cholesterol (mg dL <sup>-1</sup> )	Triglyceride (mg dL <sup>-1</sup> )	Protein (g dL <sup>-1</sup> )	Phosphorus (mg dL <sup>-1</sup> )
Control	6.8 ± 0.7 <sup>c</sup>	75.6 ± 3.7 <sup>b</sup>	59.2 ± 3.1 <sup>b</sup>	526.7 ± 28.2 <sup>c</sup>	2.21 ± 0.04 <sup>c</sup>	15.6 ± 0.78 <sup>c</sup>
20	113.9 ± 5.5 <sup>b</sup>	97.3 ± 7.3 <sup>a</sup>	255.9 ± 12.4 <sup>a</sup>	1427 ± 111 <sup>b</sup>	12.64 ± 1.5 <sup>b</sup>	61.7 ± 4.6 <sup>b</sup>
50	136.9 ± 9.1 <sup>a</sup>	88.7 ± 8.5 <sup>b</sup>	298.1 ± 23.4 <sup>a</sup>	2070 ± 217 <sup>a</sup>	21.11 ± 2.6 <sup>a</sup>	135.2 ± 16.9 <sup>a</sup>

Different superscripts in columns show the significant differences between data (P<0.05).

**Table 4.** Whole body composition (%) of Stellate sturgeon with different levels of E<sub>2</sub> in diet.

E2 levels (mg kg <sup>-1</sup> )	Moisture	Crude protein	Crude lipid	Ash
Control	75.6 ± 0.4	14.62 ± 0.19	9.2 ± 0.4	2.91 ± 0.1
25	74.6 ± 0.2	15.05 ± 0.12	8.7 ± 0.5	2.57 ± 0.1
50	74.7 ± 0.4	14.45 ± 0.17	8.4 ± 0.4	2.54 ± 0.1

## DISCUSSION

The results of the present study showed that E<sub>2</sub> significantly increased the growth of Stellate sturgeon. Fish in E<sub>2</sub> treated groups grew significantly slower than control fish during the period of treatment, indicating significant growth depression of E<sub>2</sub>, especially in the high dose of 50 mg E<sub>2</sub>. This result is consistent with most of previous reports. Funk *et al.*, (1973) in pink salmon *Onchorhynchus gorbuscha*, Johnstone *et al.* (1978) in rainbow trout *Onchorhynchus mykiss*, Blázquez *et al.* (1998) in European sea bass *Dicentrarchus labrax*, Woo *et al.* (1993) in red sea bream *Chrysophrys major*, Hendry *et al.* (2003) in Atlantic halibut, *Hippoglossus hippoglossus* and Flynn & Benfey, (2007) in shortness Sturgeon, *Acipenser brevirostrum* have reported E<sub>2</sub>-related growth depression. Goetz *et al.*, (1979) also found a dose dependent decrease in the size of Coho salmon treated with high E<sub>2</sub> concentrations. However, estrogen enhanced growth in *Pleuronectes platessa* (Cowey *et al.* 1973), Coho salmon, *Oncorhynchus kisutch* (Yu *et al.*, 1979), yellow perch and Japanese eel *Anguilla japonica* (Satoh & Nimura, 1991). In European eels, *Anguilla anguilla* an increase of the concentration of E<sub>2</sub> from 1 mg kg<sup>-1</sup> to 15 mg kg<sup>-1</sup> increased the growth of fish (Degani, 1986). These discrepancies in the growth effects caused by E<sub>2</sub> may indicate difficulties in the method of hormone treatment in hormone dosage, duration of

hormone treatment and age of the fish receiving the hormone.

The results of present study showed that diet supplemented with E<sub>2</sub> resulted in significant changes in blood indices of Stellate sturgeon. These changes in WBC, RBC, Hb and Hct values were clearly observed in E<sub>2</sub> treated groups. With increasing doses of hormone in the diet, WBC, RBC, Hb and Hct values were decreased. However, other blood indices were not significantly influenced by E<sub>2</sub> treatment. The results of this study are in line with other studies which reported Hb and Hct values were significantly lower in lake trout *Salvelinus fontinalis* (Schafhauser-Smith and Benfey, 2003), chum salmon *Oncorhynchus keta* (Haney *et al.*, 1992), *Carassius auratus* (Wang & Belosevic, 1994) and *Cyprinus carpio* (Watanuki *et al.*, 2002) when E<sub>2</sub> administrated in the diet.

In the present study, we found that values of blood biochemical parameters except glucose in E<sub>2</sub> treated groups were considerably higher than control treatment. Levels of these parameters significantly increased as the E<sub>2</sub> levels were raised. The results of this study are inconsistent with Mei-Ping *et al.* (2009) in Siberian sturgeon, *Acipenser baerii* fed with doses of 1 and 10 mg kg<sup>-1</sup> E<sub>2</sub> for 90 days. Chiba *et al.* (1993) stated that E<sub>2</sub> may play a role in food consumption and conversion in fish. Other roles of E<sub>2</sub> in growth may involve protein and fat synthesis. Japanese eel (Satoh & Nimura, 1991) and slow growing eel *Anguilla anguilla* (Degani,

1986) showed alternations in amount of protein and fat in the muscle following hormone treatment. In contrast, we did not observe any alteration in body composition of *A. stellatus*.

In summary, considering the effects of E2 on survival, growth, blood indices and body composition, we conclude that the E2 in even low dose can suppress the growth and changes in blood and biochemical parameters. Further study is also required to understand the growth improvement mechanism in the Stellate Sturgeon.

#### ACKNOWLEDGMENTS

The authors express their sincere appreciation to the people who gave their time, advice, and support to this study. The authors also thank the staff of Dr. Yousefpour Fish Hatchery Center for their technical help during this work.

#### REFERENCES

- Akhundov, M.M. & Fedorov, K.E. (1994) Effects of exogenous estradiol on the formation of ovaries in juvenile starlet *Acipenser ruthenus*. *Journal of Ichthyology*, 35: 109-120.
- Association of Official Analytical Chemists (AOAC). (1995) *Official methods of analysis of the AOAC*. 16th ed. Arlington (VA).
- Bayunova, L., Barannikova, I. & Semenkova, T. (2002) Sturgeon stress reactions in aquaculture. *Journal of Applied Ichthyology*, 18: 397-404.
- Biswas, A.K., Seoka, M., Ueno, K., Yong, A.S.K., Biswas, B.K., Kim, Y., Takii, K. & Kumai, H., (2008) Growth performance and physiological responses in striped knife jaw, *Oplegnathus fasciatus*, held under different photoperiods. *Aquaculture*, 279: 42-46.
- Blázquez, M., Zanuy, S., Carrillo, M. & Piferrer, F. (1998) Structural and functional effects of early exposure to estradiol-17 $\beta$  and 17 $\alpha$ -ethynylestradiol on the gonads of the gonochoristic teleost *Dicentrarchus labrax*. *Fish Physiology and Biochemistry*, 18: 37-47.
- Chatzifotis, S., Muje, P. & Pavlidis, V. (2004) Evolution of tissue composition and serum metabolites during gonadal development in the common dentex (*Dentex dentex*). *Aquaculture*, 236: 557-573.
- Chiba, H., Iwatsuki, K., Hayam, K. & Yamauchi, K. (1993) Effects of dietary estradiol-17 $\beta$  on feminization, growth and body composition the Japanese eel (*Anguilla japonica*). *Comparative Biochemistry and Physiology, Part A*. 106: 367-371.
- Cowey, C.B., Pope, J., A., Andron, J.W. & Blair, A. (1973) Studies on the nutrition of marine flatfish. The effect of oral administration of diethylstilbesterol and cyroheptadiene on the growth of *Pleuronectes platessa*. *Marine Biology*, 19: 14-21.
- Degani, G. (1986) Effect of dietary 17- $\beta$ -estradiol and 17 $\alpha$ -methyltestosterone on growth and body composition of slow growing elvers. *Anguilla anguilla* (L.). *Comparative Biochemistry and Physiology, Part A*. 85: 243-247.
- Falahatkar, B., Poursaeid, S., Meknatkhah, B., Khara, H. & Efatpanah, I. (2014) Long-term effects of intraperitoneal injection of estradiol-17 $\beta$  on the growth and physiology of juvenile stellate sturgeon *Acipenser stellatus*. *Fish Physiology and Biochemistry*, 40: 365-373.
- Flynn, S.R. & Benfey, T.J. (2007) Sex differentiation and aspects of gametogenesis in shortnose sturgeon, *Acipenser brevirostrum*, Lesueur. *Journal of Fish Biology*, 70(4): 1027-1044.
- Funk, J.D., Donaldson, E.M. & Dye H.M. (1973) Induction of precocious sexual development in female pink salmon (*Oncorhynchus gorbuscha*). *Canadian Journal of Zoology-Revue Canadienne De Zoologie*, 5: 493-500.
- Goetz, F.W., Donaldson, E.M. & Hunter, G.A. (1979) Effects of estradiol-17 $\beta$  and 17 $\alpha$ -methyltestosterone on gonadal differentiation in the coho salmon, *Oncorhynchus kisutch*. *Aquaculture*, 17: 267-278.
- Haney, D.C., Hursh, D.A., Mix, M.C. & Winton, J.R. (1992) Physiological and haematological changes in chum salmon artificially infected with erythrocytic necrosis virus. *Journal of Aquatic Animal Health*, 4: 48-57.

- Hendry, C.I., Martin-Robichaud, D.J. & Benfey, T.J. (2003) Hormonal sex reversal of Atlantic halibut (*Hippoglossus hippoglossus* L). *Aquaculture*, 219: 769-781.
- Johnstone, R., Simpson, T.H. & Youngson, A.F. (1978) Sex reversal in salmonid culture. *Aquaculture*, 13: 115-134.
- Johnstone, R., Simpson, T.H. & Youngson, A.F. (1979) Sex reversal in salmonid culture. Part II. The progeny of sex-reversed rainbow trout. *Aquaculture*, 18: 13-19.
- Kubokawa, K., Watanabe, T., Yoshioka, M. & Iwata, M. (1999) Effects of acute stress on plasma cortisol, sex steroid hormone and glucose levels in male and female sockeye salmon during the breeding season. *Aquaculture*, 172: 335-349.
- Lord, L.D., Bond, J. & Thompson, R.R. (2009) Rapid steroid influences on visually-guided sexual behavior in goldfish. *Hormones and Behavior*, 56: 519-526.
- Malison, I.A., Kayes, T.B., Wentworth, B.C. & Amundson, C.H. (1988) Growth and feeding responses of male versus female yellow perch (*Perca flavescens*) treated with estradiol-17 $\beta$ . *Canadian Journal of Fisheries and Aquatic Science*, 45: 1942-1948.
- Mei-Ping, T., Ping, Z., Tao, Z., Long-Zhen, Z. & Shi-Wei, Y. (2009) Effects of dietary estradiol-17 $\beta$  on reared juvenile Siberian sturgeon *Acipenser baerii*. 6<sup>th</sup> international symposium on sturgeon, Wuhan (China) Sept. 23-26, pp: 157-172.
- Ranzani-Paiva, M.J.T., Ishikawa, C.M., das Eiras, A.C. & da Silveira, V.R. (2004) Effects of an experimental challenge with *Mycobacterium marinum* on the blood parameters of Nile Tilapia, *Oreochromis niloticus* (Linnaeus, 1757). *Brazilian Archives of Biology and Technology*, 47:945-953.
- Satoh, H., Nimura, Y. (1991) Growth promotion in Japanese eel by the oral administration of an estrogen (diethylstilbestrol). *Nippon Suisan Gakkaishi*, 57: 21-27.
- Schafhauser-Smith, D. & Benfey, T.J. (2003) The effects of long-term estradiol-17 $\beta$  treatment on the growth and physiology of female triploid brook trout (*Salvelinus fontinalis*). *General and Comparative Endocrinology*, 131: 9-20.
- Shelton, W.L. & Mims, S.D. (1998) Induced sex reversal in gynogenetic paddlefish. *Aquaculture* 98, Book of Abstracts.
- Wang, R. & Belosevic, M. (1994) Estradiol increases susceptibility of goldfish to *Trypanosoma danilewskyi*. *Developmental & Comparative Immunology*, 18: 377-387.
- Watanuki, H., Yamaguchi, T. & Sakai, M. (2002) Suppression in function of phagocytic cells in common carp *Cyprinus carpio* L. injected with estradiol, progesterone or 11-ketotestosterone. *Comparative Biochemistry and Physiology, Part C*, 132: 407-413.
- Woo, N.Y.S., Chung, A.S.B. & Ng, T.B. (1993) Influence of oral administration of estradiol-17 $\beta$  and testosterone on growth, digestion, food conversion and metabolism in the under yearling red sea bream, *Chrysophrys major*. *Fish Physiology and Biochemistry*, 10: 377-387.
- Yu, T.C., Shinnhuber, R. & Hendricks, J.D. (1979) Effect of steroid hormones on the growth of juvenile coho salmon (*Oncorhynchus kisutch*). *Aquaculture*, 16: 351-359.

## اثر ۱۷-بتا استرادیول جیره بر عملکرد رشد، ترکیب بدن و شاخص های خونی ازون برون، *Acipenser stellatus*

ح. خارا<sup>۱\*</sup>، ب. مکننت خواه<sup>۲</sup>، ب. فلاحتکار<sup>۳</sup>، م. احمدنژاد<sup>۴</sup>، س. پورسعید<sup>۵</sup>

- ۱- گروه شیلات، واحد لاهیجان، دانشگاه آزاد اسلامی، لاهیجان، ایران
- ۲- مرکز تکثیر و بازسازی ذخایر ماهیان دریایی شادروان دکتر یوسف پور، سیاهکل، ایران
- ۳- گروه شیلات، دانشکده منابع طبیعی، دانشگاه گیلان، صومعه سرا، ایران
- ۴- پژوهشکده آبی پروری آبهای داخلی، بخش آبی پروری، بندر انزلی، ایران
- ۵- باشگاه پژوهشگران جوان و نخبگان، واحد رشت، دانشگاه آزاد اسلامی، رشت، ایران

(تاریخ دریافت: ۹۲/۱۲/۶ - تاریخ پذیرش: ۹۳/۵/۲۰)

### چکیده

این مطالعه روی اثرات ۱۷-بتا استرادیول جیره بر عملکرد رشد، ترکیب بدن و شاخص های خونی ماهی ازون برون انجام شد. ماهیان (میانگین وزن اولیه ۱/۱ ± ۴۰/۹ گرم، تعداد = ۶۰ عدد در هر گروه) با مقادیر مختلف ۱۷-بتا استرادیول در جیره شامل صفر (کنترل)، ۲۵ و ۵۰ میلی گرم در کیلوگرم غذا به مدت ۷ ماه تغذیه شدند. نتایج نشان دادند که رشد با افزایش سطح استرادیول کاهش می یابد. بالاترین نرخ بقا در ماهیان گروه کنترل مشاهده شد، ولی اختلاف در میان تیمارها معنی دار نبود ( $P > 0.05$ ). ترکیب بدن ماهیان در تیمارهای مختلف جیره تفاوت معنی دار نشان نداد. تعداد گلبول های سفید و قرمز، هموگلوبین و هماتوکریت با افزایش سطح ۱۷-بتا استرادیول تفاوت معنی دار کاهشی داشت ( $P < 0.05$ ). سطوح پارامترهای بیوشیمیایی پلازما شامل گلوکز، پروتئین تام، کلسترول، تری گلیسرید، کلسیم و فسفر به سطوح مختلف ۱۷-بتا استرادیول بستگی داشت، بطوریکه در گروه کنترل کمترین مقدار و در گروه با دوز ۵۰ میلی گرم در کیلوگرم غذا بیشترین مقدار را به خود اختصاص داده بودند. نتایج مطالعه حاضر سرکوب رشد با جیره حاوی سطوح بالای ۱۷-بتا استرادیول و تغییرات شاخص های خونی را به همراه برخی اطلاعات پایه بر روی اثر هورمون استروژن در فیزیولوژی ماهیان خاویاری را نشان داد.

\* مؤلف مسئول