

[Research]

Primary culture of ovarian follicular cells of Sterlet, *Acipenser ruthenus* to develop an *in vitro* system

M. R. Nowruzfashkhami^{1, 2*}, M. Sudagar¹, M. Bahmani³, N. Salamat⁴, M. Mazandrani¹, M. A. Yazdani Sadati²

1- Breeding and Rearing Department, Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran

2- International Sturgeon Research Institute, Agricultural Research Education and Extension Organization (AEERO), Rasht, Iran

3- Agricultural Research Education and Extension Organization (AEERO), Tehran, Iran

4- Marine Biology Department, Marine Sciences Faculty, Khorramshahr University of Marine Sciences and Technology, Khorramshahr, Iran

* Corresponding author's E-mail: Nowruzfashkhami@yahoo.com

(Received: May. 19.2015 Accepted: Oct. 14.2015)

ABSTRACT

The aim of the present study was to develop an *in vitro* system for functional investigation of ovarian follicular cells in Sterlet, *Acipenser ruthenus*. Oocytes for the primary culture were obtained from the ovaries of a 6 years old Sterlet 729 g in weight and 47 cm in total length. The oocytes were in advanced vitellogenesis stage (PI >10). A part of the ovary (containing about 300 follicles) was removed, ovarian follicles isolated by manually removing those from the interstitial tissue and washed with sterile phosphate buffered saline (PBS) containing antibiotics and Amphotericin B. Follicular cells were separated by treating oocytes with 0.25% trypsin-EDTA in Ca²⁺ and Mg²⁺ free PBS and cultured in medium L-15 supplemented with 20% FBS, streptomycin sulphate (Gibco, 100 µg.ml⁻¹), penicillin G potassium (Gibco, 100 IU.ml⁻¹) and Amphotericin B (Gibco, 2.5 µg.ml⁻¹) at 22 °C. The concentrations of Testosterone (T), Estradiol-17β (E2), Progesterone (P4) and 17α-hydroxyprogesteron (17αOHP) in the medium were measured at days 3, 5 & 7 by the Enzyme-Linked Immunosorbent Assay. According to the results, the ovarian follicular cells of Sterlet proliferated in L-15 medium were steroidogenically active as expressed by the secretion of T, E2, P4 & 17αOHP. Testosterone was the dominant hormone secreted by cultivated follicular cells, which was correlated closely with the end of vitellogenesis in the isolated oocytes. Decrease in production of these hormones was greater at days 3 & 4 in comparison with those at days 5 & 6. By successfully culturing ovarian follicular cells of Sterlet in L-15 culture medium, an *in vitro* system was developed which enables functional studies to be carried out similar to the *in vivo* situation in the ovarian follicles.

Key words: Sterlet, *Acipenser ruthenus*, Primary culture, Follicular cells, *In vitro*

INTRODUCTION

Fish oocytes, like other vertebrates, are surrounded by follicular cells (Unal *et al.* 2005). These cells play a vital role in fish reproduction as well as in the synthesis of steroid hormones, growth and final maturation of oocytes (Nagahama 1994). Several studies have evaluated the synthesis and secretion of steroid hormones by follicular cells in fish (Nagahama *et al.* 1982; Lessman 1991; Venkatesh *et al.* 1992;

Kime *et al.* 1992). It has also been documented that the fish follicular cells remained alive and continued to secrete hormones in culture medium (Stoklosowa & Epler 1985; Petrino *et al.* 1989 a, b; Salamat *et al.* 2010). The primary culture of ovarian follicular cells throughout the year can serve as a valuable tool to study the seasonal variations in hormones secreted by these cells (Galas *et al.* 1999). Reproduction in sturgeons is regulated by the hypothalamus-

pituitary-gonadal (HPG) axis (Dettlaff *et al.* 1993; Doroshov *et al.* 1997). Gonadotropin releasing hormone (GnRH) is secreted from the hypothalamus under the influence of environmental factors (such as photoperiod and temperature). This hormone regulates the synthesis and release of gonadotropins (GTH I and GTH II) by the pituitary (Lescheid *et al.* 1995; Moberg *et al.* 1995; Querat *et al.* 2000). These hormones are transported to the follicular cells through bloodstream and induce them to secrete steroid hormones (Tyler *et al.* 1991). These hormones are involved in vitellogenesis, growth and final maturation of oocytes (Moberg *et al.* 1995).

With regard to the drastic decline in natural stocks of sturgeons during the recent years, the artificial breeding of these fish provides effective means for the restoration and sustainable use of sturgeon stocks. Artificial breeding of sturgeons first started with Sterlet, *Acipenser ruthenus* (Webb & Doroshov 2011). The Sterlet, belonging to the family Acipenseridae, is a native species in Volga River. It is also found in riverine regions of the Black, Azov and Balkan Seas. In their natural habitats, females attain sexual maturity between 4 and 7 years which can be reduced to 3 or 4 years under artificial rearing conditions. The eggs of Sterlet are 1.91 to 2 mm in diameter (Chebanov & Galich 2013). This species adapts easily to rearing conditions and does not require more special attention. It also feeds on a wide range of food items (Holcik 1989). Success in artificial breeding programs of sturgeon depends on the selection of female breeders that respond to hormone injection and produce the suitable quality post-vitellogenic oocytes, understanding of the factors that significantly affect the growth and maturation of oocytes, gamete quality, synthesis and secretion of the related hormones. Follicular cells are the key of the success in the sturgeon aquaculture. The present study was aimed at developing an *in vitro* system for functional studies of ovarian follicular cells in order to achieve suitable guidelines for the success in artificial breeding programs of sturgeons in the

future. In this regards, the purpose of the present experiment was to culture the oocytes follicular cells in *in vitro* condition to study the viability and secretion of these cells.

MATERIALS AND METHODS

A sixty years old sterlet sturgeon (729 g in weight and 47.6 cm in length), was provided from International Sturgeon Research Institute, (Rasht, north of Iran) in March 2015. Fish was anesthetized by clove powder (200 ppm), abdominal area was cleaned with 70% alcohol and dissected to remove a part of the ovary (containing about 300 follicles). The ovary was attained at advanced stage of vitellogenesis (PI>10) according to Chapman *et al.* (2007). The protocol developed by Stoklosowa and Epler (1985) was followed for isolating of ovarian follicular cells with some modifications. The follicles were washed in a petri dish containing 5 ml sterile phosphate-buffered saline (PBS), streptomycin sulphate (200 µg.ml⁻¹), penicillin G potassium (Gibco 200 IU.ml⁻¹) and Amphotericin B (5 µg.ml⁻¹). After a minute the follicles were transferred to another petri dish containing 5 ml L-15 Medium (Sigma), streptomycin sulphate (Gibco, 200 µg.ml⁻¹), penicillin G potassium (Gibco, 200 IU.ml⁻¹) and Amphotericin B (Gibco, 5 µg.ml⁻¹). After 10 minutes incubation at 22 °C, the culture medium was drained and 5 ml of fresh culture medium of the same composition was added to the petri dish.

The culture medium was drained after 10 minutes and 5 ml of fresh culture medium of the same composition was added to the petri dish. The interstitial connective tissue of the oocytes was removed manually under a stereomicroscope (Nikon). The culture medium was drained and the connective tissue free follicles were washed in PBS solution containing streptomycin sulphate (200 µg.ml⁻¹), penicillin G potassium (200 IU.ml⁻¹) and Amphotericin B (5 µg.ml⁻¹) for three times. PBS was replaced by 5 ml 0.25 % Trypsin-EDTA (Gibco) in Ca²⁺ and Mg²⁺ free PBS (Gibco). Trypsinization was carried out at the optimum temperature of 37 °C (Galas & Epler 2002).

The process of trypsinization and the state of follicular envelop were monitored by an inverted microscope (Nikon, Model Eclipse-Ti). Trypsinization was carried out twice for 25 minutes and once for 10 minutes ($2 \times 25'$ and $1 \times 10'$). After each incubation time, supernatant fractions (containing the follicular cells) were pooled and centrifuged, then the supernatant was discarded and 1 ml Fetal Bovine Serum (FBS) was added to the sediment to inhibit trypsin action. After the third trypsinization and centrifugation, supernatant was removed and the deposited follicular cells were re-suspended in the medium L-15 and then diluted to a concentration of 1.25×10^6 cells.ml⁻¹ culture medium. After assessing viability of cells by trypan blue (0.4%) exclusion test (Doyle & Griffins 1998), the follicular cells were cultured (1.25×10^6 cells.ml⁻¹) in medium L-15 supplemented with 20% FBS, streptomycin sulphate (Gibco, 100 µg.ml⁻¹), penicillin G potassium (Gibco, 100 IU.ml⁻¹) and Amphotericin B (Gibco, 2.5 µg.ml⁻¹) and then transferred into an incubator at 22 °C. Culture medium was collected on days 3, 5 and 7, centrifuged and supernatant was stored at -20

°C (Epler *et al.* 1997) for further analysis of steroid hormones produced including testosterone (T), progesterone (P4), 17α-hydroxyprogesterone (17αOHP) and estradiol-17β (E2). Steroids analyses were carried out by Enzyme-Linked Immune Sorbent Assay (ELISA).

After determining the concentrations of the steroids, the data were analyzed by One - Way analysis of variance (ANOVA) test (Holm 1979) and the significance of the differences between the means was determined using LSD test (Fisher 1935) performed by SPSS (Version 17) at 0.95% confidence level. Differences were considered significant at $p < 0.05$ for all analyses. Graphs were plotted using Excel 2007 software package.

RESULTS

By removing the interstitial connective tissue, separated oocytes surrounded only by the follicular layers were obtained (Fig. 1), and follicular cells were observed as a thin layer when oocytes were examined under an inverted microscope (Fig. 2). Oocytes and the follicular layers were expanded after first trypsinization (Fig. 3).




Fig. 1. Separated oocytes from interstitial tissue of the ovary in L-15 Medium.

Further trypsinization resulted in detaching of follicular layers from the rest of the oocytes and formation of the cell suspension containing theca and granulosa cells. The isolated follicular cells were observed under an inverted microscope as separated cells (Fig. 4A) and

Sometimes formed clumps (Fig. 4B). The number of obtained follicular cells was 1.25×10^6 cells.ml⁻¹, and the viability of cells, determined by the trypan blue exclusion test, was 92%. After 48 hours, follicular cells were attached to the bottom of the culture vessel and

proliferated. The cells were mostly fibroblast-like in shape (Fig. 5).




Fig. 2. Sterlet oocyte ($\times 600$).




Fig. 3. Trypsinized oocyte of Sterlet ($\times 600$).

The propagation of cells increased in the following days (Fig. 6) and after one week the cultured cells had almost completely covered the bottom of the culture vessel (Fig. 7). Based on the results, the concentrations of T secreted by cultivated follicular cells after 2, 4 and 6 days were 15.93 ± 1.57 , 10.30 ± 0.20 and 9.63 ± 0.24 ng.ml⁻¹, respectively (Fig. 8). According to Fig. 8, T secretion in the culture medium decreased over time. T levels on days 5 and 7 were significantly lower than that on day 3. However T level on days 5 and 7 did not differ significantly. E2 concentration in culture

medium after 2, 4 and 6 days were recorded as 443.33 ± 12.01 , 170.0 ± 11.54 , and 130.0 ± 15.8 pg.ml⁻¹, respectively (Fig. 9). E2 levels gradually decreased in the culture medium and were significantly lower on days 5 and 7 than on day 3, although no significant differences were detected in E2 levels between days 5 and 7.

Levels of P4 in the culture medium increased gradually from 260.0 ± 17.32 pg.ml⁻¹ on day 3 to 303.33 ± 34.80 pg.ml⁻¹ on day 5 and further increased to 330.3 ± 36.05 pg.ml⁻¹ on day 7, although no significant differences were

recorded in P4 levels between at the beginning and at the end of the culture (Fig. 10).




Fig. 4. Follicular cells of Sterlet oocytes 1 h after culturing (A: $\times 120$; B: $\times 1200$).



Fig. 5. Fibroblast-like cells after two days of plating ($\times 300$).

Similarly, $17\alpha\text{OHP}$ levels in the culture medium also increased after days 2, 4 and 6. $17\alpha\text{OHP}$ level was $106.67 \pm 8.81 \text{ pg.ml}^{-1}$ on day 3 which significantly increased to 180.0 ± 5.78 and $193.33 \pm 12.01 \text{ pg.ml}^{-1}$ on days 5 and 7, respectively (Fig. 11). No significant differences were detected between $17\alpha\text{OHP}$ levels on days 5 and 7. Based on the results obtained from this

study, cultivated follicular cells of Sterlet in L-15 medium, containing 20% FBS, streptomycin sulphate ($100 \text{ }\mu\text{g.ml}^{-1}$), penicillin G potassium (100 IU.ml^{-1}) and amphotericin B ($2.5 \text{ }\mu\text{g.ml}^{-1}$) at $22 \text{ }^\circ\text{C}$, responded and adapted to the culture, attached to the bottom of the culture vessel, proliferated and secreted steroids, T, E2, P4 and $17\alpha\text{OHP}$.




Fig. 6. Fibroblast - like cells after four days ($\times 300$).




Fig. 7. Fibroblast - like cells after a week ($\times 120$).




Fig. 8. Concentrations of T in the follicular cells culture medium (as mean \pm SEM; $P = 0.700$).




Fig. 9. Concentrations of E2 in the follicular cells culture medium (as mean \pm SEM; P =0.001).




Fig. 10. Concentrations of P4 in the follicular cells culture medium (as mean \pm SEM; P =0.010).




Fig. 11. Concentrations of 17 α -OHP in the follicular cells culture medium (as mean \pm SEM; P=0.287).

DISCUSSION

Primary culture of ovarian follicular cells was previously reported in common carp, *Cyprinus carpio* (Epler et al. 1997; Galas et al. 1999; Salamat

et al. 2010) and European sea bass, *Dicentrarchus labrax* (Crespo et al. 2012). Epler et al. (1997) cultured interstitial cells of carp ovary and follicular cells isolated from the carp oocytes in

medium M199 and compared the levels of secreted hormones, P_4 , E_2 and androgens (A_s) in the two seasonal periods pre-spawning, active time (April), and resting time (December). In another study Galas *et al.* (1999) cultured the follicular cells of common carp in culture medium 199 after isolating them and compared the seasonal variation of steroid secretions by the dispersed carp ovarian follicular cells as influenced by carp pituitary homogenate (CPH), human chorionic gonadotropin (HCG) and 17α , 20β -dihydroxy-4-pregnen-3-one ($17,20\beta$ -P) and without the influence of these factors. Similarly Salamat *et al.* (2010) successfully cultured follicular cells of common carp in M199 and reported that isolated follicular cells continued to grow and proliferate in the culture medium and also E_2 and 17α OHP were the main steroids secreted by the cultivated cells. Crespo *et al.* (2012) isolated ovarian follicular cells of European sea bass and set up a procedure to culture them to study their steroidogenic capacity at difference temperatures. All of these studies made employing of the medium 199 for culturing the follicular cells of the fish under study. However, in the present study, L-15 medium (Sigma) was employed to support follicular cells growth in medium without carbon dioxide equilibration. According to several researchers L-15 is the ideal culture medium for fish cell growth (Fernandez *et al.* 1993a; Sathe *et al.* 1995). In the present study, the isolated Sterlet ovarian follicular cells were successfully cultured and the cultivated cells secreted higher levels of T in the culture medium than the other hormones. In fish, T hormone is produced by theca cells. In most fish, follicular cells are made up of two layers, an outer thecal layer and an inner granulosa layer. These two layers of cells are separated by the basement membrane. The cells of these two layers show aromatase activity and the aromatizing potential of the granulosa cells is higher than that of theca (Kagawa *et al.* 1982b). Nagahama (1994) found that the theca and granulosa cells are the major sites of the production of steroid hormones. However it is difficult to recognize

and differentiate these two types of cells in the culture medium (Galas *et al.* 1999; Galas & Epler 2002). Crespo *et al.* (2012) reported that granulosa cells were darker than theca cells. Stoklosowa & Epler (1985) after tripsinizing the ovarian follicles of common carp identified both types of cells, theca and granulosa, in the cell suspension. Darker cells with stronger aromatase activity were identified as granulosa cells, and lighter cells with weaker enzyme activity were reported as theca cells. In the present study, although large and small cells were observed among the follicular cells, it was not possible to clearly distinguish theca cells from granulosa cells in the culture medium. The production of T by follicular cells in the culture medium was more than that of other hormones. At the beginning of vitellogenesis, T is initially synthesized by the theca cells, and is later transformed to E_2 in the granulosa cells under the influence of the enzyme P_{450} aromatase (Kagawa *et al.* 1983). This hormone is then transported to the liver in the bloodstream where it is transformed into vitellogenin. From the liver, vitellogenin circulates in the bloodstream and reaches the ovaries where it is incorporated in the developing oocytes (Patino & Sullivan 2002). Early vitellogenesis is accompanied by an increase in T secretion by follicular cells. Plasma T levels in fish increase during this stage and decrease at the end of vitellogenesis. However plasma T levels remained higher than other steroid hormones until the end of vitellogenesis, before oocyte maturation (Webb *et al.* 2001; Barannikova *et al.* 2002a), and even until the beginning of follicular atresia and oocyte degeneration (Webb *et al.* 2001; Barannikova *et al.* 2002b; Linares- Casenave *et al.* 2002). In addition to being the precursor to E_2 , Testosterone has a feedback effect on the H-P-G axis (Bukovskaya *et al.* 1999; Safi *et al.* 1999; Bayunova *et al.* 2002; Ceapa *et al.* 2002), and also plays an important role in inducing maturation (Kime, 1993) and germinal vesicle breakdown (GVBD) in sturgeon (Semenkova *et al.* 2006). Therefore the plasma levels of this hormone after vitellogenesis have been reported to be

higher than those of other sex steroids including E2 in sturgeon as well as in white sturgeon (Webb *et al.* 2001; Webb *et al.* 2002).

It appears that testosterone remains stored in follicular cells before employing (Epler *et al.* 1997).

Sampling of Sterlet oocytes in the present study was carried out in late March 2015 when oocytes were at the end of vitellogenesis (post vitellogenesis) or before final maturation (PI>10) and this may explain the higher levels of T as compared to E2 in the culture medium. In the present study, the levels of T and E2 at the first 48 h were higher than at the following days.

However the levels of these hormones decreased from day 3, with T secretions decreasing faster than E2. In addition, decrease in hormone secretions on days 3 and 4 were greater than that on days 5 and 6. In all other teleosts as in vertebrates (Stoklosowa & Epler 1985; Benninghoff & Thomas 2006) the secretions of T and E2 by theca and granulosa cells gradually decreased from the beginning of primary culture due to the decreased aromatase potential of follicular cells (Stoklosowa *et al.* 1982).

Similar results were reported in common carp by Epler *et al.* (1997) who found that the aromatase activity of follicular cells decreased from day 4 through 6 of culture.

They believed that follicular cells dedifferentiated and lost their characteristic secretory pattern.

Increased levels of P4 and 17 α OHP in the culture medium from day 3 onwards seems to be associated with the role of these hormones in inducing oocyte maturation.

Research on several sturgeon species has shown that P4 exhibits a strong potential to induce oocyte maturation *in vitro* (Lutes 1985; Distaff *et al.* 1993; Williot *et al.* 1997; Mojazi Amiri 1999a). Semenikova *et al.* (2006) reported P4 to be the most effective sex steroid hormone in inducing final oocyte maturation in sterlet sturgeon with levels reaching 3.3 times the initial level during final maturation.

Galas *et al.* (1999) reported ovarian follicular cells of common carp in culture secreted steroid hormones for 48 hours *in vitro* in a manner reflecting *in vivo* pattern at the time of cell harvest, similar to the data obtained from the culture cells in mammals (Szoltys *et al.* 1982). Similarly, Galas *et al.* (1999) reported that the *in vitro* changes of steroid fluctuations by ovarian follicular cells in common carp during different seasons were similar to the annual fluctuations described in the blood plasma and ovarian tissue homogenates by Galas & Bieniarz (1989). The ovarian follicular cells isolated in the present study belonged to post vitellogenesis and hormonal data suggest that the steroid secretions by follicular cells in culture *in vitro* are similar to the pattern reported *in vivo*.

The secretion of T and E₂ was high for 48 hours and decreased from day 3, while secretion of T remained higher than the other sexsteroid hormones. However, levels of P4 and 17 α OHP gradually increased after day 2.

Although several studies have been conducted on oocyte endocrinology and the effect of various factors including steroid hormones on the *in vitro* oocyte maturation in various sturgeon species including white sturgeon, *Acipenser transmontanus* (Webb *et al.* 2000); A hybrid sturgeon, Bester (Mojazi Amiri *et al.* 2001; Baranikova 2006; Smenekova *et al.* 2006; Goncharov *et al.* 2009; Azarine *et al.* 2013); Persian sturgeon *Acipenser persicus* (Taneh *et al.* 2012; Azarine *et al.* 2013), but there was no report on the primary culture of isolated ovarian follicular cells in sturgeons.

CONCLUSION

In the present study, the follicular cells of sterlet sturgeon cultured in the supplemented L-15 medium adapted well as an adherent monolayer culture, remained viable and continued to secrete steroid hormones into the culture medium.

These results indicate we have successfully developed an *in vitro* homologous system for follicular cells of oocyte in sterlet sturgeon that resembles the *in vivo* situation.

ACKNOWLEDGEMENTS

The authors would like to thank employees of International Sturgeon Research Institute, Ali Hushyar and Hushang Yeganeh, in the Breeding and Rearing Department for helping us to handle the fish, chief and employees from departments of Genetic and Biotechnology, Ecology, Aquatic Animal Health, Physiology and Biochemistry for allowing us to use their laboratory and equipments during the experiment, and also Mr. Jalil Jalilpour from the Aquatic Animal Health Department for microbial controlling of samples.

REFERENCES

- Azarin, H, Imanpoor, MR & Pourdehghani, M 2013, Effect of 17α , 20β -dihydroxy progesterone on *in vitro* oocyte maturation in Persian sturgeon, *Acipenser persicus* and Sterlet, *Acipenser ruthenus*. *Journal of Aquaculture Research and Development*, 4: 1-4.
- Barannikova, IA, Bayunova, LV, Saenko, II & Semenkova, TB 2002a, Steroid hormones content in the blood serum of Russian sturgeon, *Acipenser gueldenstaedtii* at reproductive function disturbances in comparison with normal sexual cycle. *Journal of Ichthyology*, 42: 221-224 (In Russian).
- Barannikova, IA, Dyubin, VP, Bayunova, LV & Semenkova, TB 2002b, Steroids in the control of reproductive function of Fish. *Neuroscience and Behavioral Physiology*, 32: 141-148.
- Barannikova, IA 2006, Sex steroids and oocyte maturation in the sterlet, *Acipenser ruthenus* L. *Journal of Applied Ichthyology*, 22: 340-355.
- Bayunova, L, Barannikova, I & Semenkova, T 2002, Sturgeon stress reaction in aquaculture. *Journal of Applied Ichthyology*, 18: 397-404.
- Benninghoff, AD & Thomas, P 2006, Gonadotropin regulation of testosterone synthesis by theca and granulosa cells of Atlantic croaker, *Micropogonias undulatus*: II. Involvement of a mitogen-activated protein kinase pathway. *General and Comparative Endocrinology*, 147: 288-296.
- Bukovskaya, OS, Bayunova, LV, Blokhin, SV & Boev, AA 1999, the effects of acute stress on hormonal serum levels in Russian and Stellate sturgeon during inducing maturation. *Journal of Applied Ichthyology*, 15: 308-309.
- Ceapa, C, Willot, P, Le Menn, F & Davail-Cnisset, B 2002, Plasma sex steroids and vitellogenin levels in stellate sturgeon, *Acipenser stellatus* (Pallas) during spawning migration in the Danube River. *Journal of Applied Ichthyology*, 18: 391-396.
- Chapman, F & Van Eenennaam, J 2007, *Sturgeon aquaculture specialized techniques: Determining the stage of sexual maturity in female sturgeon for artificial spawning: The egg polarization index or PI*. University of Florida, IFAS Extension, document FA153.
- Chebanov, MS & Galich, EV 2013, *Sturgeon Hatchery Manual*. FAO Fisheries and aquaculture technical paper, 338 p.
- Crespo, B, Zanuy, S & Go'mez, A 2012, Development of an *in vitro* system for functional studies of ovarian follicular cells in European sea bass, *Dicentrarchus labrax*. *Cytotechnology*, 65: 273-286.
- Detlaff, TA, Ginsburg, AS & Schmalhausen, OI 1993, *Sturgeon fishes: developmental biology and aquaculture*. Springer-Verlag, New York. 300 p.
- Doroshov, SI, Moberg, GP & Van Eenennaam, JP 1997, Observations on the reproductive cycle of cultured white sturgeon, *Acipenser transmontanus*. *Environmental Biology of Fishes*, 48: 265-278.
- Doyle, A & Griffiths, JB 1998, *Cell and tissue culture: Laboratory procedures in biotechnology*. John Wiley & Sons Ltd, 352 p.
- Epler, P, Galas, J & Stoklosowa, S, 1997, Steroidogenic activity of carp ovarian follicular and interstitial cells at the pre-spawning and resting time: A tissue culture approach. *Comparative Biochemistry and Physiology*, 116C: 167-170.
- Fernandez, RD, Yoshimizu, M, Ezura, Y & Kimura, T 1993a, Comparative growth response of fish cell lines in different media, temperature and sodium chloride concentrations. *Fish Pathology*, 28: 27-34.

- Fisher, RA 1935, *The Design of Experiments*. Edinburgh, Oliver and Boyd, 261 p.
- Galas, J & Bieniarz, K 1989, Seasonal changes of sex steroids in mature female and male carp, *Cyprinus carpio* L. *Polish Archives of Hydrobiology*, 36: 407-416.
- Galas, J, Epler, P & Stoklosowa, S 1999, Seasonal response of carp, *Cyprinus carpio* ovarian cells to stimulation by various hormones as measured by steroid secretion. *Tissue culture approach and Endocrine Regulation*, 33: 125-132.
- Galas, J & Epler, P 2002, Does prolactin affect steroid secretion by isolated rainbow trout ovarian cells. *Comparative Biochemistry and Physiology*, 132: 287-297.
- Goncharov, BF, Skoblina, MN, Trubnikova, OB & Chebanov, MS 2009, Influence of temperature on the Sterlet, (*Acipenser ruthenus* L.) ovarian follicles state. biology, conservation and sustainable development of sturgeons. *Fish and Fisheries Series*, 29: 205-214.
- Holcik, J 1989, *The freshwater fishes of Europe, Vol I/II. General Introduction to fishes Acipenseriformes*. Aula-Verlag Wiesbaden pp. 342-362.
- Holm, S 1979, A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, 6: 65-70.
- Kagawa, H, Young, G, Adachi, S & Nagahama, Y 1982b, Estradiol-17 β production in amago salmon *Oncorhynchus rhodurus* ovarian follicle: role of thecal and granulosa cells. *General and Comparative Endocrinology*, 47: 440-448.
- Kagawa, H, Young, G & Nagahama, Y 1983, Relationship between seasonal plasma estradiol- 17 β and testosterone levels and in vitro production by ovarian follicles of Amago salmon, *Oncorhynchus rhodurus*. *Boilogy of Reproduction*, 29: 301-309.
- Kime, D, Scott, AP & Canario, A 1992, *In vitro* biosynthesis of steroid including 11-deoxycortisol and 5 α -pregnane-3 β , 7 α , 17, 20 β -tetrol by ovaries of the goldfish *Carassius auratus* during the stage of oocyte final maturation. *General and Comparative Endocrinology*, 87:375-384.
- Kime, D 1993, 'Classical' and 'non-classical' reproductive steroids in fish. *Reviews in Fish Biology and Fisheries*, 3: 160-180.
- Lessman, ChA, 1991, Metabolism of progestagens during *in vitro* meiotic maturation of follicle enclosed oocytes of the goldfish *Carassius auratus*. *Journal of Experimental Zoology*, 259: 59 -68.
- Lescheid, DW, Powell, JF, Fischer, WH, Park, M, Craig, A, Bukovskaya, O, Barannikova, IA & Sherwood, NM 1995, Mammalian gonadotropin-releasing hormone (GnRH) identified by primary structure in Russian sturgeon, *Acipenser gueldenstaedtii*. *Regulatory Peptides*, 55: 299-309.
- Linares-Casenave, J, Van Eenennaam, JP & Doroshov, SI 2002, Ultrastructural and histological observations on temperature-induced follicular ovarian atresia in the white sturgeon. *Journal of Applied Ichthyology*, 18: 382-390.
- Lutes, PB, 1985, Oocyte maturation in white sturgeon, *Acipenser transmontanus*: some mechanisms and applications. In: *North American sturgeons*. Edited by FP Binkowski & SI Doroshov, W, Junk Publishers, Dordrecht, Netherlands, pp. 87-92.
- Moberg, GP, Watson, JG, Doroshov, SI, Papkoff, H & Pavlick, RJJr 1995, Physiological evidence for two sturgeon gonadotropins in *Acipenser transmontanus*. *Aquaculture*, 135: 27-39.
- Mojazi Amiri, B, Maebayashi, M, Adachi, S, Moberg, GP, Doroshov, SI & Yamauchi, K 1999a, *In vitro* Steroidogenesis by testicular fragments and ovarian follicles in a hybrid sturgeon (Bester). *Journal of Fish Physiology and Biochemistry*, 21: 1-14.
- Mojazi Amiri, B, Maebayashi, M, Omoto, N, Adachi, S & Yamauchi, K 2001, *In vitro* oocyte Maturation in a hybrid sturgeon (Bester): changes in the germinal vesicle breakdown and 17, 20 β -dihydroxy-4-pregnen-3-one Production. *Journal of Agricultural Science and Technology*, 3: 199-207.

- Nagahama, Y, Kagawa, H & Young, G 1982, Cellular sources of sex steroids in teleost gonads. *Canadian Journal of Fisheries and Aquatic Sciences*, 39:48-55.
- Nagahama, Y, 1994, Endocrine regulation of gametogenesis in fish. *International Journal of Developmental Biology*, 38: 217-229.
- Patino, R & Sullivan, CV 2002, ovarian follicle growth, maturation, and ovulation in teleost fish. *Fish Physiology and Biochemistry*, 26: 57-70.
- Petrino, TR, Lin, Y-WP & Wallace, RA 1989a, Steroidogenesis in *Fundulus heteroclitus* I. Production of 17 α -hydroxy, 20 β -dihydroprogesterone, testosterone and 17 β -estradiol by pre-maturational follicles *in vitro*. *General and Comparative Endocrinology*, 73: 147-156.
- Petrino, TR, Greeley, MS JR, Selman, K, Lin, Y-WP & Wallace, RA 1989b, Steroidogenesis in *Fundulus heteroclitus* II. Production of 17 α -hydroxy-20 β -dihydroprogesterone, testosterone and 17 β -estradiol by various components of the ovarian follicle. *General and Comparative Endocrinology*, 76: 230-240.
- Querat, B, Sellouk, A & Salmon, C 2000, Phylogenetic analysis of the vertebrate glycoprotein hormone family including new sequences of sturgeon (*Acipenser baerii*) β subunits of the two gonadotropins and the thyroid-stimulating hormone. *Journal of Biology of Reproduction*, 63: 222-228.
- Safi, S, Mojabi, A, Takami, G, Nowrouzian, I, Mahmoodi, M & Bokaei, S 1999, Evaluation of sturgeon follicle-stimulating hormone, luteinizing hormone, estradiol, progesterone and testosterone in *Acipenser persicus* serum to identify fertile broodstock. *Journal of Applied Ichthyology*, 15: 196-198.
- Salamat, N, Erfani Majd, N, Hashemitabar, M, Mesbah M & Ahangarpour, A 2010, Isolation of common carp ovarian follicular cells and evaluation of their endocrine activity in primary cell culture. *Iranian Journal of Fisheries Sciences*, 9: 305-314.
- Sathe, PS, Maurya, DT, Basu, A, Gogate, SS & Banerjee, K, 1995, Establishment and characterization of a new Fish cell line, MG-3, from the gills of *Mrigal Cirrhinus mrigala*. *Indian Journal of Experimental Biology*, 33: 589-594.
- Semenkova, TB, Barannikova, I, Kime, D, McAllister, B, Bayunova, L, Dyubin, V & Kolmakov, NN 2002, Sex steroid profiles in female and male stellate sturgeon (*Acipenser stellatus* Pallas) during final maturation induced by hormonal treatment. *Journal of Applied Ichthyology*, 18: 375-381.
- Semenkova, TB, Bayunova, LV, Webb, MAH, Kolmakov, NN, Romanov, AG & Barannikova, IA 2006, Effect of progestins on germinal vesicle break down in sturgeon follicles *in vitro*. *Journal of Applied Ichthyology*, 22: 353-357.
- Stoklosowa, S, Gregoraszczyk, E & Channing, CP 1982, Estrogen and progesterone secretion by isolated cultured porcine thecal and granulosa cells. *Biology of Reproduction*. 26: 943-952.
- Stoklosowa, S & Epler, P 1985, The endocrine activity of isolated follicular cells of the carp vary in prime culture. *General and Comparative Endocrinology*, 58: 386-393.
- Szoltys, M, Stoklosowa, S & Casten, F 1982, Hormonal secretion of cultured rat ovarian follicles isolated at various hours of proestrus. *In vitro*, 18: 463-468.
- Taneh, B, Abtahi, B & Nazari, RM 2012, *in vitro* assessment of final oocyte maturation index (GVBD) in Persian sturgeon and broodstock selection. *Experimental Animal Biology*, 1: 35-46. (In Persian)
- Trant, JM & Thomas, P 1988, Structure-activity relationships of steroids in inducing germinal vesicle breakdown of Atlantic croaker *Micropogonias undulatus* oocytes *in vitro*. *General and Comparative Endocrinology*, 71: 307-317.
- Tung LC, Chen, SN & Kou, GH 1991, Three cell lines derived from spleen and kidney of black porgy *Acanthopagrus schlegelii*. *Fish Pathology*, 26: 109-117.

- Tyler, CR, Sumpster, JP, Kawauchi, H & Swanson, P 1991, Involvement of gonadotropin in the uptake of vitellogenin into vitellogenic oocytes of the rainbow trout, *Oncorhynchus mykiss*. *General and Comparative Endocrinology*, 84: 291-299.
- Ueda, H, Hiroi, O, Yamauchi, K, Hara, A, Kagawa, H, Adachi, S & Nagahama, Y 1991, Changes in serum steroid levels in *in vitro* steroid hormone production in the ovaries of female chum salmon *Oncorhynchus keta* during spawning migration. *Nippon Suisan Gakkaishi*, 57: 881-884.
- Unal, G, Karakisi, H & Elp, M 2005, Ovarian follicle ultrastructure and changes in levels of ovarian steroids during oogenesis in *Chalcalburnus tarichi* (Pallas). *Turkish Journal of Veterinary and Animal Sciences*, 29: 645-653.
- Venkatesh, B, Tan, CH & Lam, TJ 1992, Effects of neurohypophyseal and adenohipophyseal hormones, steroids, eicosanoids, and extrafollicular tissue on ovulation *in vitro* of guppy, *Poecilia reticulata* embryos. *General and Comparative Endocrinology*, 87: 20-27.
- Webb, MAH, Van Eenennaam, JP & Doroshov, SI, 2000, Effects of steroid hormones on *in vitro* oocyte maturation in white sturgeon, *Acipenser transmontanus*. *Fish Physiology and Biochemistry*, 23: 317-325.
- Webb, MAH, Van Eenennaam, JP, Feist, G, Linares-Casenave, J, Fitzpatrick, M, Schreck, CB & Doroshov, SI 2001, Effects of thermal regime on ovarian maturation and plasma sex steroids in farmed white sturgeon. *Aquaculture*, 201: 137-151.
- Webb, MAH, Feist, GW, Trant, JM, Van Eenennaam, JP, Fitzpatrick, MS, Schreck, CB & Doroshov, SI 2002, Ovarian steroidogenesis in white sturgeon, *Acipenser transmontanus* during oocyte maturation and induced ovulation. *General and Comparative Endocrinology*, 129: 27-38.
- Webb, MAH & Doroshov, SI 2011, Importance of environmental endocrinology in fisheries management and aquaculture of sturgeons. *General and Comparative Endocrinology*, 170: 313- 321.
- Williot, P 1997, Effects of incubation media on maturation of isolated ovarian follicles of Siberian sturgeon, *Acipenser baeri* (Brandt) induced by sturgeon gonadotropic preparation or 17 α , 20 β - dihydroxy progesterone. *Comparative Biochemistry and Physiology*, 118: 285-293.

کشت اولیه سلول‌های فولیکولی تاسماهی استرلیاد *Acipenser ruthenus*

به منظور ایجاد یک سیستم آزمایشگاهی

م. ر. نوروزفشخامی^{۱،۲*}، م. سوداگر^۱، م. بهمنی^۳، ن. سلامت^۴، م. مازندرانی^۱، م. ع. یزدانی ساداتی^۲

۱- گروه تکثیر و پرورش آبزیان، دانشکده شیلات و محیط زیست، دانشگاه علوم کشاورزی و منابع طبیعی گرگان، گرگان، ایران

۲- موسسه تحقیقات بین المللی تاسماهیان دریای خزر، سازمان تحقیقات، آموزش و ترویج کشاورزی، رشت، ایران

۳- سازمان تحقیقات، آموزش و ترویج کشاورزی، تهران، ایران

۴- گروه زیست شناسی دریا، دانشکده علوم دریایی، دانشگاه علوم و فنون دریایی خرمشهر، خرمشهر، ایران

(تاریخ دریافت: ۹۴/۲/۲۹ تاریخ پذیرش: ۹۴/۷/۲۲)

چکیده

هدف از انجام این تحقیق ایجاد یک سیستم آزمایشگاهی (*in vitro*) برای بررسی عملکرد سلول‌های فولیکولی تخمک تاسماهی استرلیاد *Acipenser ruthenus* بود. تخمک‌های مورد نیاز برای کشت اولیه از تخمدان یک عدد ماهی استرلیاد ۶ ساله پرورشی به وزن ۷۲۹ گرم و به طول ۴۷ سانتی متر به دست آمد. تخمک‌ها در مرحله پیشرفته ویتلوژنز بودند ($10 > PI$). یک قسمت از تخمدان (دارای حدود ۳۰۰ عدد تخمک‌ها) از تخمدان خارج شد، تخمک‌ها از بافت‌های بنابینی جدا شدند و با محلول بافر نمکی فسفات (PBS) استریل حاوی آنتی‌بیوتیک‌ها و آمفوتریسین B شستشو شدند. سلول‌های فولیکولی از طریق تیمار تخمک‌ها با محلول trypsin-EDTA (۰/۲۵٪) حل شده در PBS عاری از یون‌های کلسیم و منیزیم جدا شدند و در محیط کشت L-15 حاوی ۲۰٪ سرم جنین گاو (FBS)، استرپتومایسین سولفات ($100 \mu g.ml^{-1}$; Gibco)، پنی‌سیلین پنتاسیم ($100 IU.ml^{-1}$ G; Gibco) و آمفوتریسین B ($2/5 \mu g.ml^{-1}$; Gibco) تحت دمای ۲۲ درجه سانتی - گراد کشت داده شدند. غلظت هورمون‌های تستوسترون (T)، ۱۷-بتا استرادیول (E2)، پروژسترون (P4) و ۱۷-آلفا هیدروکسی پروژسترون ($17\alpha OHP$) موجود در محیط کشت در روزهای سوم، پنجم و هفتم با استفاده از روش ELISA اندازه‌گیری شدند. بر اساس نتایج به دست آمده، سلول‌های فولیکولی تکثیرشده تاسماهی استرلیاد در محیط کشت L-15 از نظر تولید هورمون‌های استروئیدی فعال بودند بطوری که این موضوع با ترشح هورمون‌های T، E2، P4 و $17\alpha OHP$ نمایان شد. هورمون T بیشترین مقدار هورمون ترشح شده توسط سلول‌های فولیکولی کشت داده شده بود که نشان دهنده اواخر مرحله ویتلوژنز در تخمک‌های جدا شده بود. کاهش تولید این هورمون‌ها طی روزهای سوم و چهارم بیش از روزهای پنجم و ششم بود. با کشت موفقیت‌آمیز سلول‌های فولیکولی تخمک تاسماهی استرلیاد در محیط کشت L-15 یک سیستم آزمایشگاهی (*in vitro*) ایجاد شد که انجام مطالعات عملکردی را مقدور می‌سازد و شبیه شرایط این سلول‌ها در تخمک‌ها است.

* مولف مسئول