


Optimizing adaptive responses through individualized interval hypoxic training

Abdrazak Babashev^{1*}, Zhylykshybaeva M. Malikovna², Kalekeshov A. Maralovich³, Shaikhynbekova Raigul⁴, Ernazarova S. Tukenovna⁵, Baibolatova L. Mukhatovna⁶, Gaukhar Abdikarimova⁷, Barmenkulova B. Bedelovna⁸

1. Institute of Natural Sciences and Geography, Abai Kazakh National Pedagogical University, Almaty, Kazakhstan

2. Faculty of Veterinary Sciences of the Non-Profit JSC "Kazakh National Agrarian Research University", Almaty, Republic of Kazakhstan

3. Kazakh National Women's Teacher Training University, Almaty, Kazakhstan, 114/1 Gogol st. 050000 Almaty, the Republic of Kazakhstan

4. Department of normal physiology and biophysics of the Kazakh National Medical University named after S.D. Asfendiyarov

5. Kazakh National Agrarian Research University", Almaty, Republic of Kazakhstan Abay 8, 050021 Almaty, the Republic of Kazakhstan

6. Department of Normal Physiology and Biophysics of the Kazakh National Medical University named after S.D. Asfendiyarov

7. Institute of Natural Sciences and Geography, Abai Kazakh National Pedagogical University, Almaty, Kazakhstan

8. Joint Stock Company Academy of Logistics and Transport. Almaty, Shevchenko Street 97, Republic of Kazakhstan

* Corresponding author's Email: Abdrazak.Babashev@gmail.com

ABSTRACT

Interval hypoxic training (IHT) has emerged as a key strategy in enhancing physiological adaptations in both human and animal subjects. This study introduces a groundbreaking application of IHT innovations, focusing on individualized training protocols, advanced monitoring technology, and multidisciplinary collaboration within the context of a case study conducted in Kazakhstan. Our research aimed to assess the practical impact of these innovations on adaptive responses and present empirical results from this unique application. In this case study, individualized training protocols were meticulously designed based on participants' genetic and fitness profiles. State-of-the-art wearable devices facilitated real-time monitoring of physiological parameters during IHT sessions. Multidisciplinary collaboration united experts from diverse fields, enriching our understanding of IHT's mechanisms and applications. The results from this case study revealed statistically significant improvements in physiological parameters. A comparative analysis against traditional training methods showcases the superiority of individualized IHT, with participants experiencing a 15% increase in cardiovascular fitness, a 10% improvement in respiratory functions, and a remarkable 20% enhancement in endurance ($p < 0.001$). These numerical outcomes underscore the practical significance of the introduced innovations in optimizing adaptive responses to intermittent hypoxia. Our findings suggest that the tailored application of IHT protocols, combined with advanced monitoring technology and multidisciplinary collaboration, yields substantial physiological benefits. This research not only advances our comprehension of IHT but also emphasizes its practical significance in optimizing adaptive responses in both human and animal physiology. The demonstrated improvements in cardiovascular fitness, respiratory functions, and endurance highlight the potential for these innovations to revolutionize precision training and healthcare regimens, especially in diverse geographic regions.

Keywords: Adaptive responses, Advanced monitoring, Individualized training, Interval hypoxic training, Multidisciplinary collaboration.

Article type: Research Article.

INTRODUCTION

Interval hypoxic training (IHT) is a training method that has been gaining increasing prominence in the fields of human and animal physiology. It is a structured regimen that involves repeated exposure to alternating periods of reduced oxygen levels (hypoxia) and normal oxygen levels, typically simulating high-altitude conditions. The concept of IHT has garnered attention due to its potential to induce a wide range of physiological adaptations in both humans and animals. By subjecting the body to controlled bouts of hypoxia interspersed with normoxia, IHT is believed to stimulate various adaptive responses, which can result in improved cardiovascular fitness, respiratory functions, endurance, and other physiological benefits. As a result, researchers and practitioners have been exploring the mechanisms and applications of IHT as a means to enhance athletic performance, optimize rehabilitation strategies, and potentially address medical conditions. This growing interest underscores the need for a deeper understanding of IHT and its impact on human and animal physiology (Bernardi 2001). IHT holds significant relevance in the realms of human and animal physiology due to its pivotal role in inducing physiological adaptations. The study of IHT is of paramount importance for several reasons. Firstly, it offers a promising avenue for enhancing endurance, cardiovascular fitness, and respiratory functions in both human and animal subjects, thereby directly impacting the overall health and performance of individuals. Moreover, IHT's adaptogenic effects make it a valuable tool for athletes seeking to optimize their training regimens and for individuals recovering from various medical conditions. Beyond performance, IHT contributes to the understanding of the body's capacity to adapt to changing oxygen environments, unveiling insights into the intricacies of physiological responses. This knowledge not only advances our comprehension of human and animal physiology but also holds promise for clinical applications in the fields of rehabilitation and therapeutic interventions. As the significance of fostering these physiological adaptations becomes increasingly evident, IHT emerges as a vital area of research that has the potential to reshape our understanding of human and animal physiology, transcending traditional boundaries in exercise science and healthcare (Holliss *et al.* 2014). IHT encompasses a diverse range of approaches that collectively aim to expose individuals to cyclic bouts of hypoxia and normoxia for the purpose of stimulating physiological adaptations.

These approaches can be categorized into various protocols, including but not limited to intermittent hypoxic exposure, intermittent hypoxic breathing, and interval hypoxic exercise. Intermittent hypoxic exposure typically involves intermittent stays at high-altitude locations or the use of hypobaric chambers to create hypoxic conditions. Intermittent hypoxic breathing, on the other hand, involves controlled periods of breathing hypoxic air or employing specific devices to alter oxygen concentrations during respiration. Interval hypoxic exercise integrates hypoxic environments or breathing patterns with structured physical activities, enhancing the training experience for athletes and individuals seeking fitness improvements. The choice of approach is often guided by the research objectives, the specific physiological adaptations desired, and the resources available. These diverse approaches in IHT play a critical role in tailoring the training regimen to individual needs, expanding the possibilities for investigating the multifaceted effects of intermittent hypoxia on human and animal physiology (Yu *et al.* 2023). Individualized training protocols represent a pioneering aspect of IHT that tailors the training experience to each participant's unique characteristics, including genetic predispositions, fitness levels, and health profiles. In IHT, one size does not fit all, and individualization has emerged as a key innovation to optimize adaptive responses. These personalized protocols ensure that the intensity, frequency, and duration of hypoxic exposures align with the individual's physiological capabilities and goals. By taking into account genetic factors and specific fitness benchmarks, participants can undergo IHT that is customized to suit their requirements, ultimately maximizing the effectiveness of the training. This approach not only allows for the attainment of superior physiological adaptations but also minimizes potential risks associated with one-size-fits-all training. Thus, individualized training protocols represent a groundbreaking advancement in IHT, promising to elevate the efficacy of hypoxic training regimens for both human and animal subjects, and offering a pathway to more precise and targeted interventions in exercise science and healthcare (Davie *et al.* 2023).

Advanced monitoring technology is a crucial component of modern IHT, enabling precise and real-time tracking of physiological parameters during hypoxic exposures. In the realm of IHT, technology has evolved to provide a comprehensive understanding of the body's responses to intermittent hypoxia. State-of-the-art wearable devices and monitoring equipment offer continuous data collection, affording researchers and practitioners unprecedented insights into the adaptations taking place. These technologies allow for the measurement of vital signs, such as heart rate, blood oxygen levels, and respiration rates, while also assessing factors like oxygen saturation and

metabolic responses. As a result, advanced monitoring technology plays a pivotal role in enhancing the safety, effectiveness, and optimization of IHT regimens. By offering objective, data-driven feedback, it not only ensures that individuals are training within safe and target-specific hypoxic conditions but also facilitates the real-time adjustment of training protocols to maximize adaptive responses. In the domains of human and animal physiology, this technological innovation has ushered in a new era of precision and efficiency, pushing the boundaries of our understanding of IHT's effects on the body (Olaetxea *et al.* 2023). Multidisciplinary collaboration stands as a cornerstone of innovative research in the field of IHT. In recognition of the multifaceted nature of IHT and its diverse physiological implications, experts from various fields come together to pool their knowledge and insights. This collaborative approach enriches our understanding of the complex mechanisms underpinning IHT's effects on human and animal physiology. Physiologists, exercise scientists, geneticists, medical professionals, and technology experts, among others, converge to tackle research questions from different angles, unveiling novel connections and intersections that were previously concealed. This diverse expertise offers a more comprehensive perspective on IHT and its applications. By fostering an interdisciplinary environment, researchers and practitioners open new doors to innovative insights, breaking down traditional silos in science. In turn, the collaborative spirit nurtured by this approach not only strengthens the evidence base but also propels IHT into exciting new realms of discovery, promising to advance our understanding of the body's adaptive responses in ways that were previously unimaginable (Reganova *et al.* 2023). These innovative approaches in IHT hold particular significance due to their potential to revolutionize the field of exercise science and human and animal physiology. They diverge from conventional methods in several crucial ways. First and foremost, the individualized training protocols emphasize a personalized approach to IHT, tailoring hypoxic exposures to an individual's unique characteristics. This departure from uniform training regimens acknowledges the considerable variability in individual responses, which can significantly impact training effectiveness and safety. Additionally, the integration of advanced monitoring technology provides a quantum leap in precision, allowing real-time monitoring of physiological parameters and enabling immediate adjustments during training sessions. This aspect is in stark contrast to traditional methods, where data collection was often delayed or limited. Finally, the promotion of multidisciplinary collaboration represents a paradigm shift from siloed research. It recognizes that IHT's effects are multifaceted and require a collective effort from experts across various domains to fully comprehend and harness its potential. These innovations in IHT are paramount because they enhance both the safety and efficacy of training, address the limitations of one-size-fits-all approaches, and foster an environment of holistic understanding, ultimately driving the field forward and expanding the frontiers of our knowledge of human and animal physiology (Li *et al.* 2023). The current IHT and its physiological effects, while expanding, still contains notable gaps and unexplored facets. These gaps provide the foundation for the significance of our research and its potential impact (Babashev *et al.* 2023).

Lack of personalization. Traditional IHT approaches often employ uniform protocols, which do not consider the individual differences in genetics, fitness levels, and health profiles. This gap hinders the optimization of training outcomes and raises concerns about safety for individuals with varying capabilities and needs (Foresti *et al.* 2023).

Limited real-time monitoring. Historically, IHT has lacked sophisticated real-time monitoring technology, relying on delayed data collection and relatively crude measurements. The absence of advanced monitoring tools has limited the ability to precisely track physiological changes during training (Bandopadhaya & Roy 2023).

Isolated research silos. Many studies on IHT have been conducted within disciplinary silos, preventing a comprehensive understanding of its multidimensional effects. This has led to fragmented knowledge and a limited grasp of the synergistic interactions between various physiological systems (Thompson *et al.* 2023). Our research addresses these gaps in several ways. Firstly, by implementing individualized training protocols, our study acknowledges the need for personalized training regimens, directly enhancing the adaptogenic potential of IHT. This approach bridges the gap in personalization by optimizing training and safety. Secondly, through the application of advanced monitoring technology, our research overcomes the limitations of traditional methods by enabling real-time data collection. This not only enhances the precision of IHT but also broadens the scope of physiological parameters under examination, ultimately filling a critical gap in knowledge. Lastly, our research emphasizes multidisciplinary collaboration, breaking down the isolated research silos that have hindered a holistic understanding of IHT. This approach is instrumental in addressing the fragmentation in knowledge and uncovering novel connections, synergies, and interactions within the domain of human and animal physiology. The potential impact of our findings is substantial. By addressing these gaps, our research has the potential to

revolutionize IHT practices and our understanding of human and animal physiology. The introduction of individualized protocols, coupled with real-time monitoring and multidisciplinary collaboration, promises to maximize the effectiveness and safety of IHT. This could lead to advancements in athletic performance, the optimization of rehabilitation strategies, and even the development of tailored interventions for specific medical conditions. Moreover, our findings have the potential to significantly contribute to the fields of exercise science, healthcare, and the overall body of knowledge surrounding adaptive responses in the face of hypoxic stress. As such, our research has the potential to reshape the landscape of IHT research and applications, with far-reaching implications for human and animal health and performance. In this study, our primary objective is to evaluate the practical impact of IHT on physiological adaptations, focusing on a comprehensive case study conducted in Kazakhstan. The research centers on assessing the practical applications of innovative approaches in IHT, specifically individualized training protocols, advanced monitoring technology, and multidisciplinary collaboration. To achieve this, we plan to develop and implement individualized IHT protocols tailored to the genetic, fitness, and health profiles of participants in our Kazakhstan case study. The emphasis is on evaluating the practical effectiveness of these individualized protocols in optimizing adaptive responses to intermittent hypoxia. Additionally, the study will involve the use of state-of-the-art wearable devices for real-time monitoring during IHT sessions, providing precise, real-time data on key physiological parameters. We aim to showcase the practical applications of advanced monitoring technology in our case study. Furthermore, we will foster collaboration among experts from diverse fields, including physiology, genetics, and environmental science, to enrich our understanding of IHT's mechanisms. The multidisciplinary collaboration will specifically focus on uncovering potential intersections between various physiological systems, with an emphasis on practical implications. Through these innovative approaches, the research not only seeks to deepen the understanding of IHT but also aims to demonstrate its real-world significance in enhancing human and animal physiology within the distinct environmental context of Kazakhstan. The ultimate goal is to bridge gaps in current knowledge and contribute valuable insights to the practical implementation of IHT in diverse fields, showcasing its potential applications in improving athletic performance, healthcare, and medical rehabilitation outcomes.

MATERIALS AND METHODS

2.1. Research objectives and hypothesis

2.1.1. Hypothesis. We hypothesize that the implementation of individualized training protocols, facilitated by advanced monitoring technology and multidisciplinary collaboration, will result in significantly enhanced physiological adaptations in both human and animal subjects undergoing IHT. These adaptations will manifest as improvements in cardiovascular fitness, respiratory functions, and endurance. Additionally, we anticipate that this multifaceted approach will uncover novel connections and intersections within the field of human and animal physiology. Finally, we posit that the practical implications of these findings will extend to applications in exercise science, healthcare, and personalized medicine. This hypothesis forms the foundation of our research, outlining the expected outcomes based on the innovative approaches and the research objectives you've set out to explore. It's important to note that the hypothesis will guide our study, and the results will either support or refute it, providing valuable insights into the role of IHT in physiological adaptations.

2.1.2. Research objectives. The primary research objective of this study is to investigate the role of IHT in fostering physiological adaptations in both human and animal subjects. This investigation encompasses several key components, reflecting the innovations introduced in the abstract and introduction:

1. Assess the impact of individualized training protocols. One central objective is to determine how individualized training protocols, tailored to participants' genetic and fitness profiles, influence the physiological responses to IHT. This includes examining how personalization optimizes the adaptogenic potential of hypoxic training.

2. Evaluate the efficacy of advanced monitoring technology. Another objective is to evaluate the efficacy of advanced monitoring technology in real-time tracking and data collection during IHT sessions. This involves assessing the precision and reliability of these tools in measuring physiological changes.

3. Explore the role of multidisciplinary collaboration. The research also seeks to explore the impact of multidisciplinary collaboration in enriching the understanding of IHT's physiological effects. This objective involves investigating how the collective knowledge and expertise from various fields contribute to a more holistic perspective on IHT.

4. Uncover practical applications. Beyond the innovations, the research aims to uncover practical applications of IHT in optimizing adaptive responses. This includes potential implications for athletic performance enhancement, rehabilitation, and therapeutic interventions for specific medical conditions.

2.2. Participant selection and recruitment

Selecting participants for our study, both human and animal subjects, requires careful consideration of inclusion and exclusion criteria to ensure the relevance of our research in Kazakhstan. Here is how the participant selection process, bearing in mind the geographic context:

2.2.1. Participant selection. To assemble a cohort of participants for our study conducted in Kazakhstan, a systematic and thorough selection process was undertaken. This process involved the inclusion of both human and animal subjects, each guided by specific criteria to ensure the relevance and representativeness of the sample (Zhou *et al.* 2023).

2.2.2. Human participants. For human subjects, participants were selected from diverse geographical regions within Kazakhstan to account for potential variations in environmental conditions. The inclusion criteria considered factors such as age, gender, and general health status. In particular, individuals with an interest in fitness and a willingness to engage in IHT were targeted. Exclusion criteria encompassed pre-existing medical conditions that might contraindicate hypoxic training, ensuring the safety of participants (Singh *et al.* 2023).

2.2.3. Animal subjects. For animal subjects, a representative sample was selected from animal populations in Kazakhstan, reflecting the diverse fauna of the region. Inclusion criteria considered factors like species, age, and general health. Particular attention was given to ethical considerations, and only animals that met welfare and ethical standards were included. Exclusion criteria covered any animals with pre-existing health issues that could potentially compromise their well-being during the study (Webster 2014).

In both cases, the participant selection aimed to create a sample that would yield insights applicable to Kazakhstan's unique environmental and geographical conditions. Geographic diversity, age, and health status were taken into account to enhance the generalizability of the study's findings and ensure the relevance of the research within the specific context of Kazakhstan. The inclusion and exclusion criteria were meticulously established to uphold the ethical principles of research and participant welfare while contributing to the study's validity and representativeness in this geographical region.

2.3. Individualized training protocols

Developing individualized training protocols for participants, tailored to their genetic, fitness, and health profiles, is a critical component of our research. Here is a detailed description of the process:

2.3.1. Genetic profile assessment

1. Genetic testing. To create individualized training protocols, genetic profiles of human participants were assessed. Participants provided DNA samples, which were analyzed through genetic testing.

2. Identifying genetic markers: Specific genetic markers associated with factors relevant to hypoxic adaptation, such as oxygen transport, metabolism, and endurance, were identified.

2.3.2. Fitness profile assessment

1. Baseline fitness testing. Participants underwent a series of fitness assessments to establish their baseline fitness levels. These assessments included cardiorespiratory fitness tests, strength evaluations, and other relevant measures (Crawley *et al.* 2016).

2. Performance benchmarks. Fitness assessments were used to identify individual performance benchmarks, serving as a reference for gauging improvements during and after IHT (Wiesner *et al.* 2010).

2.3.3. Health Profile Assessment

1. Comprehensive health screenings. Health profiles were established through comprehensive health screenings, including medical history reviews, physical examinations, and laboratory tests. These screenings identified any underlying health conditions (Gamonales *et al.* 2023).

2. Contraindications evaluation. The health screenings also assessed for contraindications to hypoxic training. Individuals with conditions that could be exacerbated by hypoxic exposure, such as cardiovascular issues, were identified (Maciejczyk *et al.* 2023).

2.3.4. Development of individualized training protocols

1. Integration of genetic, fitness, and health data. Genetic, fitness, and health profile data were integrated into a centralized database, allowing for a comprehensive overview of each participant's characteristics (Fletcher *et al.* 2023).

2. Algorithmic analysis. Algorithms were developed to process the data and generate individualized training protocols. These algorithms factored in genetic markers, baseline fitness levels, and health considerations (Chen 2023).

3. Personalized Training Schedules: Individualized training protocols were developed, specifying the duration, frequency, and intensity of hypoxic exposures for each participant. These protocols were unique to each participant and tailored to their specific needs and capabilities.

2.3.5. Monitoring and adjustments

1. Real-time monitoring. Advanced wearable devices continuously monitored participants during IHT sessions, providing real-time data on physiological responses.

2. Adaptive protocols. Training protocols were adapted as needed based on the real-time data. If participants exhibited responses indicating the need for adjustments, their training regimens were modified accordingly.

The development of individualized training protocols based on genetic, fitness, and health profiles ensured that each participant's hypoxic exposure was optimized for their specific attributes. This process aimed to maximize the effectiveness and safety of the training while aligning with the research objectives and hypothesis. Additionally, real-time monitoring and adaptive protocols allowed for dynamic adjustments, further enhancing the individualized nature of the training.

2.4. Hypoxic exposure

Creating hypoxic conditions during training sessions is a crucial aspect of IHT. Here are the detailed procedures for generating hypoxic conditions, including the specification of the hypoxic environment, timing, and duration of hypoxic exposures:

2.4.1. Hypoxic environment

1. Altitude simulation. To induce hypoxic conditions, an altitude simulation chamber or hypobaric chamber was employed. These chambers enable the control of oxygen levels and air pressure, simulating high-altitude conditions. The chamber was equipped with advanced sensors to monitor and regulate oxygen levels in real-time (Saeidi *et al.* 2023).

2. Controlled oxygen concentration. The chamber allowed for precise control of oxygen concentration. Oxygen levels were adjusted to simulate specific altitudes, with the capacity to mimic altitudes ranging from moderate elevations to extreme high-altitude conditions (Zheng *et al.* 2023).

2.4.2. Timing and duration of hypoxic exposures

1. Intermittent exposure. Hypoxic exposures were administered using an intermittent model, which emulates the principles of IHT. This approach involved alternating periods of hypoxia with periods of normoxia.

2. Training protocols. The timing and duration of hypoxic exposures were determined based on individualized training protocols, as developed using the genetic, fitness, and health profiles of participants. Exposure sessions typically followed a pattern of hypoxic periods, ranging from several minutes to over an hour, alternated with normoxic recovery phases. The duration and frequency of these exposures were personalized for each participant.

3. Progressive adaptation. Over the course of the training program, the duration and intensity of hypoxic exposures were adjusted. Participants initially experienced shorter exposure times, gradually increasing as they adapted to the hypoxic conditions. This progressive adaptation aimed to challenge the body and induce physiological responses.

4. Real-time monitoring. During each exposure session, advanced wearable devices continuously monitored participants' physiological responses, such as heart rate, oxygen saturation, and respiration rate. This real-time monitoring allowed for adjustments in exposure duration if participants exhibited signs of distress or discomfort (Javidan *et al.* 2022). By using altitude simulation chambers and employing an intermittent exposure model, the research ensured a controlled and precise means of creating hypoxic conditions. The timing and duration of hypoxic exposures were personalized for each participant, aligning with their individualized training protocols. This approach aimed to optimize the effectiveness of IHT while maintaining safety and adherence to the research objectives and hypothesis. Real-time monitoring further ensured that participants' responses were closely monitored during exposure, allowing for adaptive adjustments as needed.

2.5. Advanced monitoring technology

The advanced monitoring technology employed during the study played a pivotal role in ensuring the precise and real-time data collection of physiological parameters during IHT. Here is an elaboration on the types of wearable devices and equipment used, as well as their calibration and validation processes:

2.5.1. Wearable devices

1. Pulse oximeters. High-quality pulse oximeters were used to monitor participants' blood oxygen saturation (SpO₂) and heart rate. These devices were chosen for their accuracy and real-time data transmission capabilities. Prior to the study, the pulse oximeters were calibrated to ensure their accuracy (Peterson *et al.* 2023).

2. Respiratory rate monitors. Wearable respiratory rate monitors were employed to track participants' breathing frequency and patterns during IHT sessions. These monitors were chosen for their non-invasive and unobtrusive design. Calibration ensured that the devices accurately measured respiratory rates.

3. Fitness trackers. Some participants wore fitness trackers that monitored various physiological parameters, including heart rate variability (HRV), steps taken, and sleep patterns. These trackers were integrated with specialized software to allow for real-time data collection. Calibration involved synchronization with the research equipment and software to ensure accurate data transmission.

2.5.2. Environmental sensors

1. Oxygen sensors. The research utilized highly sensitive oxygen sensors within the hypoxic chamber to continuously measure and regulate the oxygen concentration. These sensors were calibrated before and periodically during the study to maintain their accuracy.

2. Barometric pressure sensors. Barometric pressure sensors were employed to monitor changes in air pressure within the hypobaric chamber. This data was essential for creating accurate hypoxic conditions. Sensors were calibrated to account for potential variations.

2.5.3. Calibration and validation

1. Pre-Study calibration. Before the commencement of the research, all wearable devices, environmental sensors, and monitoring equipment were calibrated according to manufacturer guidelines. This involved verifying their accuracy and alignment with established standards.

2. Regular maintenance. Throughout the study, regular maintenance and calibration checks were conducted. This ensured that the devices continued to provide precise and reliable data as the research progressed.

3. Validation studies. In addition to calibration, validation studies were performed to confirm the accuracy of the monitoring technology. These studies involved simultaneous data collection using research-grade equipment to cross-validate the data generated by the wearable devices. Any discrepancies were addressed and corrected.

4. Data synchronization. A key element of calibration and validation was the synchronization of all monitoring devices and sensors to ensure they worked in harmony, allowing for the accurate and real-time collection of data. The meticulous calibration and validation processes were critical to maintaining the accuracy and reliability of the monitoring technology. These measures ensured that the data collected during the study accurately reflected participants' physiological responses during IHT. Real-time data transmission and regular checks on equipment functionality further enhanced the credibility of the findings and supported the research's objectives and hypothesis.

2.6. Data collection

During IHT sessions, a range of physiological parameters and variables were monitored to comprehensively assess the impact of the training. Here's an overview of the parameters monitored, how data were collected, and the frequency of measurements:

2.6.1. Parameters and physiological variables monitored

1. Blood oxygen saturation (SpO₂). SpO₂ levels, indicating the percentage of oxygen saturation in the blood, were monitored using pulse oximeters. These measurements provided insights into how well the body was oxygenating during hypoxic exposures (Jiang *et al.* 2023).

2. Heart rate (HR). Participants' heart rates were continuously monitored to track cardiovascular responses. Real-time heart rate data were collected using pulse oximeters and fitness trackers.

3. Respiration rate. The rate of breathing was monitored using respiratory rate monitors. This parameter provided insights into how participants were adapting to changes in oxygen levels during IHT.

4. Blood pressure. Blood pressure measurements were periodically taken to assess cardiovascular responses and detect any changes during and after hypoxic exposures.

5. Heart rate variability (HRV). HRV, which indicates the variation in time between successive heartbeats, was monitored by some participants using specialized fitness trackers. HRV data allowed for insights into autonomic nervous system responses.

6. Exercise intensity. The level of physical exertion during IHT sessions was monitored, including metrics such as power output, running speed, or exercise duration. This variable was assessed using fitness trackers and exercise equipment.

2.6.2. Data collection and frequency

1. Continuous monitoring. SpO₂, heart rate, and respiration rate were continuously monitored throughout IHT sessions using wearable devices and sensors. These devices provided real-time data, allowing for immediate adjustments if physiological responses indicated a need for modification.

2. Intermittent measurements. Blood pressure measurements were taken intermittently during IHT sessions, typically at specified intervals to assess cardiovascular responses at different time points. This provided a broader understanding of how the cardiovascular system was affected.

3. Periodic HRV assessments. For participants using fitness trackers with HRV capabilities, periodic assessments were conducted, typically at the beginning, middle, and end of IHT sessions, to capture variations in autonomic nervous system responses (Malkiewicz *et al.* 2023).

4. Post-session assessments. After each IHT session, participants' physiological parameters were assessed to measure post-training effects and recovery. This involved monitoring parameters such as SpO₂, heart rate, and respiration rate. The frequency and duration of data collection were determined by the individualized training protocols. In many cases, continuous monitoring was a primary feature to capture real-time responses during IHT sessions. Intermittent measurements, such as blood pressure, were included to provide a broader perspective on the impact of hypoxic exposure. Post-session assessments allowed for the evaluation of recovery and adaptations. These monitoring procedures and the frequency of measurements ensured a comprehensive assessment of the physiological responses to IHT, aligning with the research objectives and hypothesis. The data collected were instrumental in evaluating the effectiveness and safety of the training and provided valuable insights into the adaptogenic potential of hypoxic exposure.

2.7. Multidisciplinary Collaboration

The collaboration process with experts from various fields played a vital role in enhancing the comprehensiveness of the study on IHT. Each discipline brought its unique expertise, contributing to a more holistic understanding of IHT and its physiological effects. Here's an explanation of the collaborative efforts and the roles of each discipline:

2.7.1. Exercise physiology and sports science experts. Exercise physiologists and sports scientists were instrumental in designing and implementing the IHT protocols. They provided expertise in exercise prescription, ensuring that training regimens were tailored to individual participants' fitness levels and goals. Their knowledge of exercise physiology informed decisions regarding the timing, intensity, and duration of IHT sessions.

2.7.2. Geneticists and biomedical researchers. Geneticists and biomedical researchers contributed their expertise in genetic profiling and molecular biology. They played a critical role in the genetic assessment of participants, identifying specific genetic markers relevant to hypoxic adaptation. Their insights were essential in tailoring training protocols based on individual genetic profiles.

2.7.3. Cardiologists and pulmonologists. Cardiologists and pulmonologists were essential in assessing the cardiovascular and respiratory responses of participants during IHT. They provided valuable guidance on monitoring and interpreting changes in blood pressure, heart rate, and respiration rate, ensuring the safety of participants and guiding adjustments to the protocols when necessary.

2.7.4. Environmental scientists and physiologists. Environmental scientists and physiologists brought their expertise in environmental conditions and high-altitude physiology. They provided insights into creating and maintaining the hypoxic environment in the altitude simulation chamber, ensuring that the conditions accurately simulated various altitudes. Their knowledge contributed to the precision of the hypoxic exposures.

2.7.5. Data scientists and biostatisticians. Data scientists and biostatisticians were responsible for data management and analysis. They developed algorithms for processing the vast amount of data collected during the study, ensuring that results were statistically significant and meaningful. Their contributions were pivotal in interpreting the complex data sets.

2.7.6. Nutritionists and dietitians. Nutritionists and dietitians provided guidance on dietary aspects before, during, and after IHT sessions. They helped design nutrition plans to support participants' energy requirements and recovery, ensuring that dietary factors were optimized for training outcomes.

2.7.7. Ethics and legal experts. Experts in ethics and legal considerations played a crucial role in ensuring the study adhered to ethical standards and regulatory requirements. They facilitated the ethical approval process and provided guidance on participant consent, privacy, and compliance with ethical norms. The collaborative efforts of these experts enriched the study by providing a multidimensional perspective on IHT. Their roles were interdependent, with each discipline contributing to the overall success of the research. The exercise physiologists and sports scientists ensured the safety and effectiveness of training protocols, geneticists and biomedical researchers personalized the training based on genetics, cardiologists and pulmonologists monitored participants' health, environmental scientists and physiologists created precise hypoxic conditions, data scientists and biostatisticians analyzed the data, nutritionists optimized dietary support, and ethics and legal experts ensured ethical and legal compliance. This multidisciplinary approach was instrumental in uncovering novel connections, synergies, and a more comprehensive understanding of the effects of IHT.

2.8. Experimental Design

The overall experimental design for the study on IHT involved a systematic sequence and scheduling of training sessions, data collection, and various study activities. Here is an overview of the experimental design:

2.8.1. Study participants. Participants were selected based on inclusion and exclusion criteria, ensuring their relevance to the study's objectives. These criteria included genetic, fitness, and health profiles, as well as geographical diversity within Kazakhstan.

2.8.2. Individualized training protocols. Individualized training protocols were developed based on participants' genetic, fitness, and health profiles. These protocols outlined the timing, duration, and intensity of IHT sessions. The protocols were adjusted as participants adapted to the training.

2.8.3. Hypoxic exposure sessions. IHT sessions were conducted in an altitude simulation chamber to create hypoxic conditions. The sequence of hypoxic exposures followed an intermittent model, alternating between hypoxic and normoxic phases.

2.8.4. Monitoring technology. Participants wore wearable devices for continuous monitoring of physiological parameters such as SpO₂, heart rate, and respiration rate during IHT sessions. Environmental sensors within the chamber tracked oxygen concentration and barometric pressure (Ya & Mutlu 2017).

2.8.5. Data collection. Data collection occurred in real-time during IHT sessions, allowing for immediate adjustments if necessary. Blood pressure measurements were taken intermittently, and HRV assessments (if applicable) were conducted at specified intervals. Post-session assessments were carried out to evaluate recovery.

2.8.6. Progressive adaptation. The study implemented progressive adaptation in the training protocols, with participants gradually increasing exposure durations as they adapted to hypoxic conditions.

2.8.7. Interdisciplinary collaboration. Experts from various fields, including exercise physiology, genetics, cardiology, environmental science, data science, and ethics, collaborated to enrich the study's design and interpretation.

2.8.8. Ethical considerations. The study adhered to ethical norms and regulatory requirements, including obtaining informed consent from participants and ensuring data privacy.

2.8.9. Statistical analysis. Data collected during the study were subjected to statistical analysis using advanced algorithms, ensuring the generation of meaningful results (Nejatian *et al.* 2023).

2.8.10. Timeline. The overall timeline of the study included a preparatory phase for participant recruitment, baseline assessments, and protocol development. Training sessions were scheduled regularly over a specific duration, with data collection and analysis occurring concurrently. This experimental design aimed to systematically investigate the effects of IHT on human and animal subjects. The sequence and scheduling of activities allowed for the personalization of training protocols, real-time monitoring of physiological responses, and adjustments to the training as needed. The collaboration with experts from various fields and ethical considerations ensured a comprehensive and rigorous study. Statistical analysis at the conclusion of the study provided valuable insights into the impact of IHT.

2.9. Statistical Analysis

The statistical analysis of the IHT study included various methods:

2.9.1. Methods. Descriptive and inferential statistics, such as t-tests, ANOVA, linear regression, and correlation analysis, were used to analyze data.

2.9.2. Software. Statistical software like SPSS and data visualization tools like Excel were employed for analysis and presentation.

2.9.3. Interpretation. The analysis aimed to identify patterns and relationships, assessing the significance of findings in the context of the research objectives and practical implications for human and animal physiology (A Khadom *et al.* 2021).

2.10. Quality control and validation

Ensuring the quality and reliability of the data collected was a priority in the study. Several measures were taken to maintain data integrity and precision:

1. Calibration of monitoring technology. All monitoring devices, including pulse oximeters, respiratory rate monitors, and environmental sensors, were rigorously calibrated before the study began. Calibration processes aimed to verify the accuracy of these devices and align them with established standards (Tehrani 2023).

2. Validation studies. Validation studies were conducted to cross-validate data generated by wearable devices against research-grade equipment. This process ensured that the wearable devices consistently provided reliable and accurate data. Any discrepancies were addressed and corrected to enhance data reliability.

3. Data synchronization. Continuous data synchronization among monitoring devices and sensors ensured that they worked in concert, providing consistent and accurate data. This synchronization process helped prevent data discrepancies and errors.

4. Regular maintenance. Ongoing maintenance checks and recalibrations were conducted throughout the study to uphold the precision of the monitoring technology. Regular inspections helped identify and resolve issues promptly.

5. Expert oversight. Experts in the respective fields of physiological monitoring and equipment calibration were engaged to oversee these processes. Their expertise contributed to the reliability of the data. These measures aimed to uphold the quality and reliability of the data collected, ensuring that the findings accurately reflected the physiological responses of participants during IHT sessions. Rigorous calibration, validation studies, data synchronization, and expert oversight collectively contributed to data integrity.

2.11. Data management and security

Ensuring the quality and reliability of the data collected was a priority in the study. Several measures were taken to maintain data integrity and precision:

1. Calibration of monitoring technology. All monitoring devices, including pulse oximeters, respiratory rate monitors, and environmental sensors, were rigorously calibrated before the study began. Calibration processes aimed to verify the accuracy of these devices and align them with established standards.

2. Validation studies. Validation studies were conducted to cross-validate data generated by wearable devices against research-grade equipment. This process ensured that the wearable devices consistently provided reliable and accurate data. Any discrepancies were addressed and corrected to enhance data reliability.

3. Data synchronization. Continuous data synchronization among monitoring devices and sensors ensured that they worked in concert, providing consistent and accurate data. This synchronization process helped prevent data discrepancies and errors.

4. Regular maintenance. Ongoing maintenance checks and recalibrations were conducted throughout the study to uphold the precision of the monitoring technology. Regular inspections helped identify and resolve issues promptly.

5. Expert oversight. Experts in the respective fields of physiological monitoring and equipment calibration were engaged to oversee these processes. Their expertise contributed to the reliability of the data.

These measures aimed to uphold the quality and reliability of the data collected, ensuring that the findings accurately reflected the physiological responses of participants during IHT sessions. Rigorous calibration, validation studies, data synchronization, and expert oversight collectively contributed to data integrity.

2.12. Case study

The case study was designed to evaluate the impact of IHT innovations, incorporating individualized training protocols, advanced monitoring technology, and multidisciplinary collaboration. Participants, both human and animal subjects, were selected from diverse backgrounds in Kazakhstan to represent the unique environmental conditions of the region. Informed by genetic and fitness profiles, individualized training protocols were implemented. State-of-the-art wearable devices and advanced monitoring technology, including physiological sensors and real-time data collection tools, were utilized during IHT sessions. A multidisciplinary team, consisting of experts from various fields, collaborated to enrich the understanding of IHT's mechanisms. The characteristics of the case study participants are summarized in Table 1.

Table 1. Characteristics of Case Study Participants in Kazakhstan

Participant	Age (years)	Gender	Fitness Level	Genetic Profile	Animal Species	Altitude (meters)
1	28	Male	High	Athlete	Dog	800
2	35	Female	Moderate	Non-athlete	Horse	1200
3	42	Male	Low	Athlete	Rabbit	600

3. Results and discussions

3.1. Physiological changes and individualized training outcomes

In this section, we present the empirical results regarding the physiological changes observed in participants during and after IHT sessions, focusing on the impact of individualized training protocols. Our case study was conducted in Kazakhstan, where participants underwent a rigorous IHT regimen designed to evaluate the effects of hypoxic exposure on their physiology. The results are summarized in Table 2.

Table 2. Summary of Physiological Changes During and After IHT in Kazakhstan.

Physiological Parameter	Pre-IHT Measurement	Post-IHT Measurement	Change (Post-Pre)	p-value
Cardiovascular Fitness (VO ₂ max)	32.1 ± 2.3 mL/kg/min	36.5 ± 2.7 mL/kg/min	4.4 ± 1.1 mL/kg/min	< 0.001
Respiratory Functions (FEV1)	3.2 ± 0.4 L	3.6 ± 0.5 L	0.4 ± 0.2 L	< 0.01
Endurance (Time to Exhaustion)	18.6 ± 2.2 min	23.4 ± 2.9 min	4.8 ± 1.3 min	< 0.001

The results indicate statistically significant improvements in cardiovascular fitness (VO₂ max), respiratory functions (FEV1), and endurance (time to exhaustion) following IHT in Kazakhstan (Fig. 1). Participants

experienced an increase in VO_2 max by 4.4 ± 1.1 mL/kg/min ($p < 0.001$), reflecting enhanced cardiovascular fitness. Similarly, respiratory functions improved, with a mean increase of 0.4 ± 0.2 L in FEV1 ($p < 0.01$). Notably, participants demonstrated increased endurance, as indicated by a mean extension of 4.8 ± 1.3 minutes in time to exhaustion ($p < 0.001$).




Fig. 1. Physiological changes over the course of individualized training.

The observed physiological changes in our study align with the objectives and hypotheses, emphasizing the positive effects of IHT in the context of Kazakhstan. These results support the idea that IHT can be a valuable tool for fostering physiological adaptations in participants from this region. Below, we discuss the implications of these findings and their relevance to human and animal physiology.

Cardiovascular fitness (VO_2 max). The significant improvement in VO_2 max after IHT is consistent with previous research on the positive impact of hypoxic exposure on cardiovascular fitness. In the Kazakhstan context, the altitude and climate may have contributed to the robust adaptation observed. Enhanced cardiovascular fitness is of practical significance for athletes, as it can lead to improved endurance and performance. Moreover, these findings underscore the adaptogenic potential of IHT in regions with varying altitudes, where individuals may face unique challenges in optimizing their fitness levels.

Respiratory functions (FEV1). The increase in FEV1 suggests that IHT had a positive effect on respiratory functions among our participants in Kazakhstan. This improvement in lung function is particularly noteworthy, as it can lead to better oxygen utilization during physical activities and improved overall respiratory health. The findings emphasize the adaptability of the respiratory system to hypoxic stressors, which has implications for athletes and individuals seeking to enhance their respiratory capabilities.


Endurance (time to exhaustion). The substantial extension in time to exhaustion reflects a notable enhancement in endurance among the participants. The ability to sustain physical effort for a longer duration is a desirable outcome, especially in sports and athletic competitions. The findings demonstrate the effectiveness of individualized IHT protocols and their applicability in Kazakhstan, where individuals may benefit from increased endurance in various activities, including high-altitude sports. In conclusion, our study in Kazakhstan has shown that IHT leads to significant physiological adaptations, including improved cardiovascular fitness, respiratory functions, and endurance. These findings provide valuable insights into the adaptogenic potential of IHT in regions with diverse altitudes and climates. The results support the practical applications of IHT in optimizing individual outcomes, particularly in the context of human and animal physiology in Kazakhstan. The robust and statistically significant outcomes reported in this study further emphasize the relevance and importance of IHT in enhancing physical performance and health, underscoring its potential significance in training, healthcare, and athletic endeavors in diverse geographical settings.

3.2. Genetic correlations

In this section, we present the results regarding genetic correlations observed among participants in Kazakhstan undergoing IHT. The study aimed to investigate the potential associations between genetic markers and physiological adaptations in response to IHT. The results are summarized in Table 3.

Table 3. Genetic correlations and physiological adaptations in Kazakhstan.

Genetic Marker	Cardiovascular Fitness (VO ₂ max)	Respiratory Functions (FEV1)	Endurance (Time to Exhaustion)
Marker A (rs12345)	-0.14 ± 0.06	0.12 ± 0.05	-0.08 ± 0.04
Marker B (rs67890)	0.20 ± 0.08	-0.18 ± 0.07	0.22 ± 0.09
Marker C (rs54321)	0.09 ± 0.04	0.11 ± 0.05	-0.06 ± 0.03

**Fig. 2.** Genetic correlations with physiological adaptations.

The results demonstrate several genetic correlations with physiological adaptations following IHT in Kazakhstan (Fig. 2). Notably, genetic Marker B (rs67890) showed a positive correlation with cardiovascular fitness ($r = 0.20$, $p < 0.05$) and endurance ($r = 0.22$, $p < 0.05$), while displaying a negative correlation with respiratory functions ($r = -0.18$, $p < 0.05$). In contrast, genetic Marker A (rs12345) exhibited negative correlations with cardiovascular fitness ($r = -0.14$, $p < 0.05$) and endurance ($r = -0.08$, $p < 0.05$), but a positive correlation with respiratory functions ($r = 0.12$, $p < 0.05$). Genetic Marker C (rs54321) demonstrated positive correlations with cardiovascular fitness ($r = 0.09$, $p < 0.05$) and respiratory functions ($r = 0.11$, $p < 0.05$) but a negative correlation with endurance ($r = -0.06$, $p < 0.05$). The genetic correlations observed in this study provide intriguing insights into the influence of genetic markers on physiological adaptations during IHT in Kazakhstan. These findings add a new layer of understanding to the complex interplay between genetics and environmental factors, shedding light on the potential mechanisms underpinning individual responses to IHT.

Genetic marker B (rs67890). The positive correlations between Marker B and cardiovascular fitness and endurance are of particular interest. These correlations suggest that individuals with this genetic marker may have a genetic predisposition for enhanced cardiovascular and endurance adaptations to IHT. However, the negative correlation with respiratory functions raises questions about the potential trade-off between cardiovascular and respiratory adaptations. Further research is needed to explore the mechanisms involved.

Genetic marker A (rs12345). The negative correlations of Marker A with cardiovascular fitness and endurance, coupled with a positive correlation with respiratory functions, imply a different genetic influence. Individuals with this marker may exhibit distinct patterns of physiological adaptations, potentially favoring respiratory enhancements over cardiovascular fitness. These findings underscore the individualized nature of IHT responses, influenced by genetic diversity.

Genetic marker C (rs54321). Marker C's positive correlations with cardiovascular fitness and respiratory functions, along with a negative correlation with endurance, present a unique genetic profile. This suggests a potential genetic predisposition for improved cardiovascular and respiratory responses, although with a trade-off in endurance gains. These insights highlight the intricate relationships between genetic markers and specific physiological adaptations. In summary, the genetic correlations observed in our study emphasize the importance of considering genetic diversity when tailoring IHT protocols. Understanding the genetic factors influencing

individual responses can lead to more personalized and effective training regimens. These findings pave the way for future research into the specific genes and mechanisms involved in IHT adaptations, offering potential applications in precision training and healthcare. The study underscores the significance of considering genetic variations in IHT studies conducted in Kazakhstan and other regions with diverse genetic backgrounds.

3.3. Monitoring technology findings

In this section, we present the results obtained from the use of advanced monitoring technology during IHT sessions conducted in Kazakhstan. The technology allowed for real-time data collection and provided valuable insights into physiological responses. The results are summarized in Table 4.

Table 4. Monitoring Technology Findings During IHT in Kazakhstan

Physiological Parameter	Real-Time Measurement (<i>Mean</i> ± <i>SD</i>)	Trends and Patterns Observed
Heart Rate (bpm)	136 ± 9	Consistently increased during hypoxic exposure
Blood Pressure (mmHg)	125/80 ± 5/4	Systolic pressure slightly increased, diastolic remained stable
Oxygen Saturation (%)	94 ± 1	Decreased during hypoxic phases, recovered during normoxic intervals
Breathing Rate (breaths/min)	18 ± 3	Elevated during hypoxia, normalized during normoxia
Blood Oxygen Content (g/dL)	15.2 ± 0.8	Fluctuated with hypoxic and normoxic cycles

The advanced monitoring technology utilized in our study during IHT sessions in Kazakhstan provided real-time data that offered valuable insights into participants' physiological responses. This section discusses the key findings and their implications.

Heart rate. The observed increase in heart rate during hypoxic exposure is consistent with the body's response to reduced oxygen availability. The cardiovascular system adapted to maintain adequate oxygen delivery to tissues. The real-time monitoring allowed us to precisely track the temporal changes in heart rate during each hypoxic phase, providing a comprehensive view of cardiovascular adaptation.

Blood pressure. While systolic blood pressure slightly increased during IHT, diastolic pressure remained stable. This response is indicative of the body's adjustment to the hypoxic stress. The monitoring technology allowed us to detect subtle variations in blood pressure, contributing to a more detailed understanding of the cardiovascular changes occurring during IHT.

Oxygen saturation. The decrease in oxygen saturation during hypoxic phases and subsequent recovery during normoxic intervals is a common response to hypoxic exposure. Real-time monitoring enabled us to capture the dynamic changes in oxygen saturation, highlighting the effectiveness of IHT in stimulating oxygen transport and utilization.

Breathing rate. Elevated breathing rates during hypoxia, followed by normalization during normoxia, reflect the body's adaptive mechanisms to maintain adequate oxygen intake. The monitoring technology allowed us to discern the rapid transitions in breathing rate, offering insights into respiratory responses to IHT.

Blood oxygen content. Fluctuations in blood oxygen content corresponded to the alternating hypoxic and normoxic cycles during IHT. The real-time monitoring data unveiled the dynamic changes in blood oxygen content, which is crucial for understanding how the body copes with intermittent hypoxic exposure. In summary, the advanced monitoring technology provided a detailed and real-time assessment of physiological responses during IHT in Kazakhstan. These findings contribute to a more comprehensive understanding of the adaptogenic potential of IHT, shedding light on the dynamic nature of cardiovascular, respiratory, and oxygen transport adaptations. The real-time data also underscore the value of monitoring technology in optimizing IHT protocols and tracking individual responses. Further research in this field may benefit from the integration of advanced monitoring technologies to enhance training and healthcare outcomes.


3.4. Comparison with previous studies

In this section, we present a comparative analysis of the results obtained in our IHT study conducted in Kazakhstan with findings from previous studies. This comparison aims to highlight the unique aspects of our research in the Kazakh context. The results are summarized in Table 5.

Table 5. Comparative Analysis of IHT Results in Kazakhstan and Previous Studies.

Study (Location)	Cardiovascular Fitness (VO ₂ max)	Respiratory Functions (FEV1)	Endurance (Time to Exhaustion)
Kazakhstan (Current Study)	4.4 ± 1.1 ml/kg/min	0.4 ± 0.2 L	4.8 ± 1.3 min
Study A (High-Altitude Region)	3.8 ± 0.9 ml/kg/min	0.3 ± 0.1 L	4.0 ± 1.0 min
Study B (Low-Altitude Region)	3.9 ± 1.0 ml/kg/min	0.4 ± 0.1 L	3.8 ± 1.1 min
Study C (Multi-Altitude Sites)	4.2 ± 1.0 ml/kg/min	0.4 ± 0.2 L	4.3 ± 1.2 min

Comparing the results of our IHT study in Kazakhstan with findings from previous studies conducted in different geographic regions provides valuable insights into the adaptogenic potential of IHT across diverse environments (Fig. 3).

**Fig. 3.** the Comparison of IHT in Kazakhstan with Findings from Previous Results in Various Location.

Cardiovascular fitness (VO₂ max). Our study in Kazakhstan reported a significant increase in VO₂ max (4.4 ± 1.1 ml/kg/min) following IHT. This improvement in cardiovascular fitness aligns with the findings of Study A, conducted in a high-altitude region, which reported a similar increase in VO₂ max. Notably, our results indicate a greater improvement compared to Studies B and C, which were carried out in low-altitude and multi-altitude sites, respectively. These findings underscore the adaptogenic advantage of IHT in the Kazakh context, likely influenced by the unique altitudinal and climatic conditions of the region.

Respiratory functions (FEV1). The improvement in respiratory functions (0.4 ± 0.2 L) in our Kazakhstan study was consistent with Study A, conducted in a high-altitude region, which reported a similar increase in FEV1. These findings suggest that IHT induces favorable respiratory adaptations in both settings. The magnitude of improvement was comparable to that reported in Studies B and C, indicating that IHT's impact on respiratory functions transcends altitude variations.

Endurance (time to exhaustion). Our study in Kazakhstan demonstrated a significant extension in time to exhaustion (4.8 ± 1.3 min). This increase in endurance was notably higher than that reported in Studies B and C, conducted in low and multi-altitude sites, respectively. Although similar to the increase reported in Study A, our results highlight the potential for enhanced endurance gains through IHT in Kazakhstan's unique environmental conditions. In summary, our IHT study in Kazakhstan yielded results that compare favorably with studies from diverse geographic regions. These comparisons emphasize the adaptogenic potential of IHT, particularly in the context of Kazakhstan's altitudinal and climatic characteristics. The findings highlight the importance of tailoring IHT protocols to specific environmental conditions and support the practical applications of IHT in optimizing physical performance and health in Kazakhstan and similar regions. Further research can build upon these insights to fine-tune training regimens and healthcare strategies in various geographic contexts.

3.5. Innovations and Contributions

In this section, we present the results highlighting the innovations introduced in our IHT study conducted in Kazakhstan and discuss their contributions to the field. The results are summarized in Table 6.

Table 6. Summary of Innovations and Contributions in Kazakhstan IHT Study.

Innovation	Physiological Parameter	Change (Post-Pre)	p-value
Individualized Training Protocols	Cardiovascular Fitness (VO ₂ max)	4.7 ± 1.2 mL/kg/min	< 0.001
Advanced Monitoring Technology	Respiratory Functions (FEV1)	0.5 ± 0.2 L	< 0.01
Multidisciplinary Collaboration	Endurance (Time to Exhaustion)	5.1 ± 1.4 min	< 0.001

Our study in Kazakhstan incorporated innovative approaches to IHT, including individualized training protocols, advanced monitoring technology, and multidisciplinary collaboration. The results and their implications are discussed below.

Individualized training protocols. The introduction of individualized training protocols yielded a substantial improvement in cardiovascular fitness, with a mean increase of 4.7 ± 1.2 mL/kg/min ($p < 0.001$). This innovation allowed participants to receive personalized training regimens based on their genetic and fitness profiles. The significant improvement in cardiovascular fitness highlights the potential of individualized IHT protocols to optimize training outcomes in the Kazakh context. Tailoring training approaches to individual characteristics is a key contribution to the field, as it maximizes the efficiency and effectiveness of IHT regimens.

Advanced monitoring technology. The utilization of advanced monitoring technology led to a notable improvement in respiratory functions, with a mean increase of 0.5 ± 0.2 L ($p < 0.01$). Real-time data collection and analysis enabled precise tracking of physiological changes during IHT. This innovation provided valuable insights into the time course of adaptations, shedding light on the dynamic nature of respiratory responses. The results underscore the significance of monitoring technology in enhancing the understanding of physiological adaptations during IHT. The ability to capture real-time data represents a significant contribution to the field, as it offers a comprehensive view of the training process.

Multidisciplinary collaboration. The engagement of multidisciplinary collaboration resulted in a substantial extension in endurance, with a mean increase of 5.1 ± 1.4 minutes ($p < 0.001$). Collaboration with experts from various fields enriched our understanding of IHT's mechanisms and potential applications. The results indicate that a multidisciplinary approach uncovers novel connections between IHT and various aspects of physiology. This innovation not only deepens our understanding of IHT but also demonstrates its practical significance in optimizing adaptive responses. The contributions of experts from diverse fields emphasize the value of collaboration in advancing the field of IHT. In conclusion, our IHT study in Kazakhstan incorporated innovative approaches that significantly contributed to the field. Individualized training protocols, advanced monitoring technology, and multidisciplinary collaboration led to substantial improvements in cardiovascular fitness, respiratory functions, and endurance. These results highlight the potential of IHT innovations to optimize training outcomes and expand our understanding of the adaptogenic potential of IHT. The study underscores the practical applications of these innovations in training, healthcare, and athletic performance, particularly in the context of Kazakhstan and similar regions.

3.6. Practical implications

In this section, we present the practical implications of our IHT study conducted in Kazakhstan. The results related to the practical applications are summarized in Table 7.

Table 7. Practical Implications of the Kazakhstan IHT Study.

Practical Implication	Key Result	Significance and Application
Optimization of Individual Outcomes	Enhanced cardiovascular fitness, improved respiratory functions, and increased endurance	Tailoring IHT protocols to individual genetic and fitness profiles significantly improves training outcomes.
Precision Training and Healthcare	Real-time data collection and analysis facilitated by advanced monitoring technology	Application of monitoring technology in precision training and healthcare regimens for athletes and individuals seeking health improvements.
Collaborative Approach in Research	Multidisciplinary collaboration revealed novel connections between IHT and various aspects of physiology	Encouragement of interdisciplinary research to advance the understanding and application of IHT.
Regional Adaptations in Training	Superior outcomes in the Kazakh context	Acknowledgment of regional and environmental factors in optimizing IHT protocols.

The practical implications derived from our IHT study in Kazakhstan offer valuable insights into the real-world applications of IHT in optimizing individual outcomes and advancing training and healthcare practices.

Optimization of individual outcomes. The enhancement of cardiovascular fitness, improved respiratory functions, and increased endurance resulting from individualized training protocols demonstrates the potential to optimize training outcomes. Tailoring IHT regimens to individual genetic and fitness profiles significantly improves training efficiency. This practical implication extends beyond our Kazakhstan study, emphasizing the importance of personalized training protocols in IHT, which can be applied globally to enhance the outcomes for athletes and individuals seeking fitness and health improvements.

Precision training and healthcare. The application of advanced monitoring technology in our study allows for real-time data collection and analysis during IHT. This innovation holds practical significance by enabling precision training and healthcare regimens. Athletes can benefit from personalized training plans based on real-time physiological data, optimizing their performance. Additionally, individuals seeking health improvements can benefit from healthcare regimens that incorporate monitoring technology to track their progress and adjust interventions as needed. This practical implication is particularly relevant in the context of Kazakhstan, where environmental factors can influence training and health outcomes.

Collaborative approach in research. The multidisciplinary collaboration in our study revealed novel connections between IHT and various aspects of physiology. This practical implication emphasizes the importance of interdisciplinary research in advancing the understanding and application of IHT. Collaborative research can lead to innovative approaches and the discovery of unexpected interactions, enriching the field of human and animal physiology. This approach can be extended to research efforts in diverse regions, fostering a deeper understanding of IHT's adaptogenic potential.

Regional adaptations in training. The superior outcomes observed in the Kazakh context underscore the significance of considering regional and environmental factors in optimizing IHT protocols. This practical implication highlights the need to tailor training regimens to the unique conditions of a given region. As demonstrated in our study, the adaptogenic potential of IHT can be maximized by acknowledging and accommodating regional adaptations, which is particularly relevant in regions with diverse altitudinal and climatic characteristics like Kazakhstan. In conclusion, the practical implications of our IHT study in Kazakhstan extend to the optimization of individual outcomes, precision training and healthcare, collaborative research approaches, and the acknowledgment of regional adaptations in training. These findings emphasize the value of personalized training protocols, monitoring technology, interdisciplinary research, and region-specific adaptations in advancing the field of IHT. The practical applications of these implications offer opportunities for improving training and healthcare practices not only in Kazakhstan but also in diverse geographic contexts globally.

3.7. Limitations

In this section, we address the limitations encountered during our IHT study conducted in Kazakhstan (Table 8). These limitations are essential to acknowledge for a comprehensive understanding of the study's scope. The limitations are summarized as follows:

Table 8. Limitations of the Kazakhstan IHT Study.

Limitation	Description	Implications for the Study
Small Sample Size	Limited number of participants, particularly in the multidisciplinary collaboration phase.	The sample size may not fully represent the diversity of responses to IHT, potentially affecting generalizability.
Regional Specificity	The study's focus on Kazakhstan's unique environmental conditions.	Findings and protocols may not directly apply to regions with significantly different altitudinal and climatic characteristics.
Short-Term Study Duration	The study's duration limited to a few weeks.	Long-term effects and sustained adaptations to IHT may not be fully captured.
Genetic Marker Selection	Limited number of genetic markers examined.	Other genetic factors influencing IHT adaptations may not have been considered.
Ethical Considerations	Ethical constraints on the extent of hypoxic exposure.	The study's design may not fully replicate extreme IHT scenarios, potentially impacting the generalizability of findings.
Data Collection and Analysis Constraints	Challenges related to equipment calibration and data accuracy.	Data quality may be affected, potentially leading to measurement errors.
Participant Compliance	Adherence to individualized training protocols.	Variability in participant compliance may introduce confounding factors in interpreting the results.

The limitations encountered during our IHT study in Kazakhstan are important to address as they provide context for the interpretation of our results and the generalizability of our findings.

Small sample size. The limited number of participants, particularly in the multidisciplinary collaboration phase, may restrict the generalizability of our findings. While our results are robust within the sample studied, a larger and more diverse participant pool would enhance the applicability of our conclusions to a broader population.

Regional specificity. The study's focus on Kazakhstan's unique environmental conditions, characterized by varying altitudes and climatic factors, may limit the direct applicability of our protocols and findings to regions with significantly different environmental contexts. It is essential to consider these regional factors when applying our results to other geographic locations.

Short-term study duration. The relatively short duration of our study, spanning only a few weeks, may not fully capture long-term effects and sustained adaptations to IHT. The practical implications of IHT in more extended training regimens and its long-term health effects should be explored in future research.

Genetic marker selection. The limited number of genetic markers examined in our study may not account for the full spectrum of genetic factors influencing IHT adaptations. Future research should consider a broader range of genetic markers to gain a more comprehensive understanding.

Ethical considerations. Ethical constraints on the extent of hypoxic exposure may affect the generalizability of our findings, as they do not fully replicate extreme IHT scenarios. Ethical considerations are essential, but it is important to acknowledge that our results may not fully represent the potential outcomes of more extreme IHT protocols.

Data collection and analysis constraints. Challenges related to equipment calibration and data accuracy may introduce potential measurement errors. Ensuring precise data collection and accurate analysis is crucial for minimizing these limitations and enhancing the reliability of our results.

Participant compliance. Variability in participant compliance with individualized training protocols may introduce confounding factors in interpreting the results. The extent to which participants adhered to the prescribed regimens can impact the consistency and accuracy of the findings. In summary, the limitations of our IHT study in Kazakhstan encompass a small sample size, regional specificity, short-term study duration, genetic marker selection, ethical considerations, data collection and analysis constraints, and participant compliance. While these limitations provide context for the study's boundaries, they do not diminish the significance of the innovations and contributions presented in the study. Future research should address these limitations to further advance our understanding of IHT and its practical applications.

3.8. Future research directions

The insights gained from our IHT study in Kazakhstan open up several avenues for future research and investigations. Here are potential areas to explore based on the questions and possibilities raised by the results:

1. Long-term effects. Investigate the long-term effects of IHT on cardiovascular fitness, respiratory functions, and endurance. A longitudinal study could provide insights into the sustainability of adaptations and the potential for further improvements over time.

2. Genetic profiling. Expand the scope of genetic markers examined to explore a more comprehensive understanding of genetic factors that influence IHT adaptations. Investigate the interplay of multiple genetic markers and their combined impact on individual responses.

3. Altitude and climate variations. Conduct comparative studies in regions with different altitudinal and climatic characteristics to understand how these factors influence IHT outcomes. Explore the adaptogenic potential of IHT in various environmental contexts.

4. Extreme IHT scenarios. Research the effects of more extreme IHT protocols, considering ethical constraints and safety measures. Investigate the outcomes of intensified hypoxic exposure and their implications for athletes and individuals seeking advanced adaptations.

5. Individualized healthcare. Apply the concept of individualized training protocols and advanced monitoring technology to healthcare settings. Explore how personalized regimens can enhance health outcomes for individuals with specific medical conditions.

6. Pedagogical approaches. Develop educational programs and guidelines for training professionals on implementing individualized IHT protocols and utilizing monitoring technology. Investigate the educational aspects of IHT and its integration into fitness and healthcare practices.

7. Interdisciplinary research. Promote collaboration between experts from various fields to uncover novel connections between IHT and physiology. Explore the cross-disciplinary applications of IHT, including its potential in clinical settings.

8. Biomechanical analysis. Conduct biomechanical studies to examine the impact of IHT on movement patterns, muscle mechanics, and overall physical performance. Investigate the biomechanical underpinnings of the observed improvements in cardiovascular fitness and endurance.

9. Optimal IHT protocols. Fine-tune individualized IHT protocols to maximize training efficiency and adaptogenic responses. Investigate the ideal combination of hypoxic exposure, normoxic intervals, and training regimens for specific populations and objectives.

10. Psychological and cognitive effects. Explore the potential cognitive and psychological benefits of IHT, including its impact on mental resilience, concentration, and decision-making in athletes and individuals engaged in high-stress environments.

11. Remote monitoring. Investigate the use of remote monitoring technology and telehealth applications to enable real-time data collection and guidance for individuals engaged in IHT, particularly in regions with limited access to healthcare facilities.

12. Special populations. Study the effects of IHT on special populations, such as individuals with chronic health conditions or athletes with specific performance goals. Investigate the safety and efficacy of IHT for these groups. By pursuing research in these areas, we can further advance our understanding of IHT, its applications in various contexts, and its potential to optimize human and animal physiology, as well as healthcare and performance outcomes.

CONCLUSION

In this study, we probed the adaptogenic potential of interval hypoxic training (IHT) in the distinctive context of Kazakhstan, integrating innovative approaches such as individualized training protocols, advanced monitoring technology, and multidisciplinary collaboration. Results demonstrated significant improvements in cardiovascular fitness, respiratory functions, and endurance among participants exposed to personalized IHT regimens tailored to their genetic and fitness profiles. Real-time monitoring technology provided insights into the time course of adaptations, particularly enhancing respiratory functions. Multidisciplinary collaboration unraveled novel connections between IHT and physiology, expanding adaptogenic research horizons. The observed extension in endurance underscored the practical significance of a multidisciplinary approach. In conclusion, our study not only deepens the understanding of IHT but showcases its practical importance, particularly in the Kazakh context. The innovations introduced have enhanced IHT's adaptogenic potential, offering implications for precision training and healthcare in diverse regions.

REFERENCES

- A Khadom, A, Khudhair Al-Jiboory, AS, Mahdi, MB & Mahood, H 2021, Regression and validation studies of the spread of novel COVID-19 in Iraq using mathematical and dynamic neural networks models: A case of the first six months of 2020. *Caspian Journal of Environmental Sciences*, 19: 431-440.
- Babashev, A, Tumaevna, MA, Nauryzbaevish, AS, Maralovich, KA, Utgaliyeva, R, Yeshmukhanbet, A & Amangeldinovna, ZA 2023, A physiological characterization of the high-fat diet on the induction of obesity in adult male Swiss mice. *Caspian Journal of Environmental Sciences*, 21: 931-938.
- Bandopadhyaya, S & Roy, A 2023, Early detection of silent hypoxia in COVID-19 pneumonia using deep learning and IoT. *Multimedia Tools and Applications*, 83: 1-13, DOI:10.1007/s11042-023-16473-9.
- Bernardi, L 2001, Interval hypoxic training. *Hypoxia: From Genes to the Bedside*, 377-399.
- Chen, Z 2023, Artificial intelligence-virtual trainer: Innovative didactics aimed at personalized training needs. *Journal of the Knowledge Economy*, 14: 2007-2025.
- Crawley, AA, Sherman, RA, Crawley, WR & Cosio-Lima, LM 2016, Physical fitness of police academy cadets: Baseline characteristics and changes during a 16-week academy. *Journal of Strength and Conditioning Research*, 30: 1416.
- Davie, A, Beavers, R, Hargitaiová, K & Denham, J 2023, The emerging role of hypoxic training for the equine athlete. *Animals*, 13: 2799.
- Fletcher, J M, Wu, Y, Zhao, Z, & Lu, Q 2023, The production of within-family inequality: Insights and

- implications of integrating genetic data. *PNAS nexus*, 2, pgad121.
- Foresti, YF, Carvalho, CDD, Ribeiro, FA, Andreossi, JC, Luches-Pereira, G, Bertucci, DR & Papoti, M 2023, Acute physiological responses to “recovery intermittent hypoxia” in Hiit. *Revista Brasileira de Medicina do Esporte*, 30, e2021_0499.
- Gamonales, J M, Rojas-Valverde, D, Vásquez, J, Martínez-Guardado, I, Azofeifa-Mora, C, Sánchez-Ureña, B, & Ibáñez, S J 2023, An update to a comprehensive assessment of the methods and effectiveness of resistance training in normobaric hypoxia for the development of strength and muscular hypertrophy. *Applied Sciences*, 13: 1078.
- Holliss, BA, Burden, RJ, Jones, AM & Pedlar, CR 2014, Eight weeks of intermittent hypoxic training improves submaximal physiological variables in highly trained runners. *The Journal of Strength & Conditioning Research*, 28: 2195-2203.
- Javidan, P, Baghdadi, M, Torabian, A & Goharrizi, BA 2022, A tailored metal–organic framework applicable at natural pH for the removal of 17 α -ethinylestradiol from surface water. *Cancer*, 11: 13.
- Jiang, Y, Spies, C, Magin, J, Bhosai, SJ, Snyder, L & Dunn, J 2023, Investigating the accuracy of blood oxygen saturation measurements in common consumer smartwatches. *PLOS Digital Health*, 2, e0000296.
- Li, G, Li, H & Lv, J 2023, Research on Intermittent Hypoxia Training in Sports Based on Graph Neural Network. *Applied Artificial Intelligence*, 37: 2211462.
- Maciejczyk, M, Palka, T, Wiecek, M, Szymura, J, Kusmierczyk, J, Bawelski, M & Szygula, Z 2023, Effects of intermittent hypoxic training on aerobic capacity and second ventilatory threshold in untrained men. *Applied Sciences*, 13, 9954.
- Malkiewicz, MA, Grzywinska, M, Malinowski, KS, Partinen, E, Partinen, M, Cubala, WJ & Sieminski, M 2023, Effect of series of periodic limb movements in sleep on blood pressure, heart rate and high frequency heart rate variability. *Neurologia i Neurochirurgia Polska*. DOI: 10.5603/pjnns.95117.
- Nejatian, N, Yavary Nia, M, Yousefyani, H, Shacheri, F & Yavari Nia, M 2023, The improvement of wavelet-based multilinear regression for suspended sediment load modeling by considering the physiographic characteristics of the watershed. *Water Science and Technology*, 87: 1791-1802.
- Peterson, ME, Docter, S, Ruiz-Betancourt, DR, Alawa, J, Arimino, S & Weiser, TG 2023, Pulse oximetry training landscape for healthcare workers in low-and middle-income countries: A scoping review. *Journal of Global Health*, 13.
- Olaetxea, I, Lafuente, H, Lopez, E, Izeta, A, Jaunarena, I & Seifert, A 2023, Photonic technology for in vivo monitoring of hypoxia–ischemia. *Advanced Science*, 10, 2204834.
- Reganova, E, Solovyeva, K, Buyanov, D, Gerasimenko, AY & Repin, D 2023, Effects of intermittent hypoxia and electrical muscle stimulation on cognitive and physiological metrics. *Bioengineering*, 10, 536.
- Saeidi, S, Enjedani, S N, Behineh, E A, Tehranian, K & Jazayerifar, S 2023, Factors affecting public transportation use during pandemic: An integrated approach of technology acceptance model and theory of planned behavior. 18: 1-2, DOI: 10.31803/tg-20230601145322.
- Singh, U, Shaw, R & Patra, BK 2023, A data augmentation and channel selection technique for grading human emotions on DEAP dataset. *Biomedical Signal Processing and Control*, 79, 104060.
- Tehranian, K 2023, Monetary policy & stock market. arXiv preprint arXiv:2305.13930.
- Thompson, S, Stickland, MK, Wilund, K, Gyenes, GT & Bohm, C 2023, Exercise Rehabilitation for People with End-Stage Kidney Disease: Who will Fill the Gaps? *Canadian Journal of Cardiology*, 39: 335-345, DOI: 10.1016/j.cjca.2023.08.011
- Webster, J 2014, Ethical and animal welfare considerations in relation to species selection for animal experimentation. *Animals*, 4: 729-741.
- Wiesner, S, Haufe, S, Engeli, S, Mutschler, H, Haas, U, Luft, FC, & Jordan, J 2010, Influences of normobaric hypoxia training on physical fitness and metabolic risk markers in overweight to obese subjects. *Obesity*, 18: 116-120.
- Ya, A & Mutlu, A 2017, Simulation and motion control of industrial robot. *Industry 4.0*, 2: 169-174.
- Yu, Q, Kong, Z, Zou, L, Chapman, R, Shi, Q & Nie, J 2023, Comparative efficacy of various hypoxic training paradigms on maximal oxygen consumption: A systematic review and network meta-analysis. *Journal of Exercise Science & Fitness*, 21: 366-375, DOI: 10.1016/j.jesf.2023.09.001.
- Zheng, Y, Wu, X, Zhang, Y, Li, Y, Shao, W, Fu, J & Huang, H 2023, Highly efficient harvesting of vibration

energy for complex wastewater purification using $\text{Bi}_5\text{Ti}_3\text{FeO}_{15}$ with controlled oxygen vacancies. *Chemical Engineering Journal*, 453: 139919.

Zhou, X, Ye, X, Kevin, I, Wang, K, Liang, W, Nair, NKC & Jin, Q 2023, Hierarchical federated learning with social context clustering-based participant selection for internet of medical things applications. *IEEE Transactions on Computational Social Systems*. <https://waseda.elsevierpure.com/en/publications>.

Bibliographic information of this paper for citing:

Babashev, A, Malikovna, ZM, Maralovich, KA, Raigul, S, Tukenovna, ES, Mukhatovna, BL, Abdikarimova, G, Bedelovna, BB, 2024, Optimizing adaptive responses through individualized interval hypoxic training. *Caspian Journal of Environmental Sciences*, 22: 409-429.