

## Floristic, life form and chorological studies of the Saqalaksar forest, Rasht, Northern Iran

Mohammadreza Ajamian, Marzieh Beygom Faghir\*, Fatemeh Bazdid Vahdati<sup>id</sup>

Department of Biology, Faculty of Science, University of Guilan, Rasht, Iran

\* Corresponding author's E-mail: faghirmb@guilan.ac.ir

### ABSTRACT

Saqalaksar forest is a relict of Caspian plain forests with an area of 200 ha in northern Iran and an altitude of 60 to 150 m a.s.l. This study was conducted to determine the floristic composition, life form and chorology of all plants of the region. Based on the results, 94 vascular plant species belonging to 80 genera and 46 families were identified in the study area. The most numerous families were Rosaceae (10 species), Fabaceae (8 species), Poaceae (7 species) as well as Asteraceae and Lamiaceae (6 species each). The classification according to the life spectrum showed that geophytes with 28 species (29.79 %) and therophytes with 25 species (26.60 %) had the highest proportion of life forms in the region. Similarly, the results of the chorological studies showed that the largest proportion of the flora consisted of pluri-regional elements (34 species, 36.17 %). As the forests investigated in this study are considered remnants of the Caspian Plain forests, urgent conservation measures are needed to protect these very important, vulnerable and threatened ecosystems.

**Keywords:** Plant, Flora, Guilan Province, Lowland forest, Chorological studies.

**Article type:** Research Article.

### INTRODUCTION

The floristic survey documents all plant species in a given geographical area, provides an essential tool for biodiversity assessment and monitoring (Barans 1969; Chiarucci & Bonini 2005; Ataei *et al.* 2021; Kudryavtsev *et al.* 2021; Mirhashemi *et al.* 2021; Aghajani *et al.* 2023; Muratovna *et al.* 2024) and shows unknown angles of the vegetation of the area (Yousefi 2009). The study of the flora of each area is very important to provide basic information on the ecological capacity of that area, to assess the effectiveness of management in the area and to protect threatened and vulnerable species (Stanford & Robinson 2005; Najafi Tire Shabankar 2008). In addition, plant geography and floristic surveys of each region show the region's position in the International Union for Conservation of Nature (IUCN 2020). Forests are a valuable asset for countries. The existence of forests in the countries creates an enormous wealth that encompasses all aspects. The Hyrcanian region, stretch in an arc along the southern shore of the Caspian Sea, is located in the provinces of Guilan, Mazandaran and Golestan in Northern Iran. They are the most important remnants of Arcto-Tertiary forests (Scharnweber *et al.* 2007) located in the Euro-Siberian geographical vegetation zone (Zohary 1973; Léonard 1988; Akhiani 1998) and in steppe forest ecotones. Unlike arid and semi-arid landscapes in most central and southern regions of Iran (Naqinezhad *et al.* 2008), Hyrcanian forests receive large amount of water vapor from evaporation of the Caspian Sea. As these water vapours meet a large barrier such as the Alborz Mountains and form dense clouds, heavy rainfall in the northern part of the mountain range has given rise to dense forests, in contrast to the southern half (Knapp 2005). Hyrcanian forests comprise 15% of the total forests of Iran and 1.1% of the total area of the country (Sagheb-Talebi 2014), which is scattered from the sea level up to 2800 m and includes several types of plants (Tohidifar *et al.* 2016). So far, numerous studies have been carried out on the floristic composition and vegetation of Hyrcanian forests (e.g. Djazirei 1964-1965; Tregubov & Mobayen 1970; Zohary 1973; Dorostkar 1974; Assadollahi 1980; Rastin 1983; Frey & Probst 1986; Klein 2001; Ghahreman *et al.* 2006; Hamzeh'ee *et al.* 2008; Naqinezhad *et al.* 2008; Atashgahi *et al.* 2009; Akhiani *et al.* 2010; Bazdid vahdati *et al.* 2014; Naqinezhad *et al.* 2015; Bazdid vahdati *et al.* 2017; Poursiadian *et al.* 2022). However, there is no information on the vegetation of the Saqalaksar forest. The aim of this study was to determine the floristic composition, life forms and chorology of each taxon. The

results will also provide (i) an overview of the biodiversity of plant species, (ii) a checklist of all vascular plants found in the study area, (iii) a comparison of the present study with recorded data from other forest areas in the Hyrcanian region, and (iv) the identification of threats to the conservation of the vegetation.

## MATERIALS AND METHODS

### Study Area

The dense forest of Saqalaksar is located beside the ecotourism village of Saqalaksar, on the southeastern border of Rasht City and northeast of Shaft City (Guilan Province). It is noteworthy that the Saqalaksar earth dam has created a natural and tourist-friendly area on the edge of the dense forest. The study area covers an area of 200 ha between latitudes  $37^{\circ}8'33''$ - $37^{\circ}9'25''$  N and longitudes  $49^{\circ}31'48''$ - $49^{\circ}32'46''$  E and includes both aquatic and terrestrial habitats. The minimum and maximum elevation of the study area varies between 60 and 150 m a.s.l., which is one of the few remnants of forest in lowland areas (Fig. 1).



Fig. 1. Location of the Saqalaksar area in Guilan Province, Iran (<https://www.google.com/>).




Fig. 2. Climatological diagram from Rasht station (2010-2020).

According to long-term statistics, climate is the general state of the climate prevailing in a particular place (Bailey 1999), so defining the climate zones of a region requires recognition of climate heterogeneity factors over time and space (Harding *et al.* 2011). Based on the De'Martonne and Emberger climate classification (De Martonne 1926; Emberger 1932), the Saqalaksar ecotourism region has a very humid climate, and according to a recent bioclimatic classification of Iran (Rivas-Martinez *et al.* 1997, 1999), the region is also known as a temperate oceanic climate (sub-Mediterranean type) or TocsM (Djamali *et al.* 2011). The average rainfall and temperature during the last 10 years (2010-2020) were 1315.37 mm and 17.03 °C, respectively. The average minimum and maximum temperatures varied between 2.7 and 27.16 °C, respectively. The maximum rainfall is in late summer and autumn (September to December; Fig. 2), when July is the hottest month of the year.

### Data collection

Data collection was performed during three seasons (spring, summer and autumn) of 2018 and 2019. The voucher specimens were deposited in the Herbarium of Guilan University, Rasht, Iran. The plant specimens were collected

of 25 relevés with an area of 400 m<sup>2</sup> for forest and 25 m<sup>2</sup> for herbaceous habitat, based on Mueller-Dombois & Ellenbreg (1974) and Djazirei (1965) and minimal areas method (Cain 1938). Floristic data were collected using relevés of 25 with an area of 400 m<sup>2</sup> for forest and 25 m<sup>2</sup> for herbaceous habitat, based on Mueller-Dombois & Ellenbreg (1974) and Djazirei (1965). The specimens were identified by Rechinger (1963-2015), Assadi *et al.* (1988-2016), Davis (1965-1985), Ghahreman (1979-2005) and Tutin *et al.* (1964-1980). The classification of flowering plants was coordinated based on APG IV (2016) and the names of the authors of the taxon using IPNI (2020). Life form classification was identified based on the Raunkiaer classification system (Raunkiaer 1934). Detailed information about phanerophytes based on Pears (1985), (megaphanerophyte = more than 30 m height, mesophanerophytes = 8-30 m height, microphanerophytes = 2-8 m, nanophanerophytes = less than 2 m height, liana = no height limit). The distribution of the species was taken from surveys, monographs and floras, especially the Flora Iranica (Rechinger 1963-2015) and Flora of Turkey (Davis 1965-1985). The terminology and delimitation of the main phytochoria followed the concepts used by Zohary (1973) and Takhtajan (1986). The abbreviations used are ES (plants distributed in the Euro-Siberian region), PL (Pluriregional, Plants distributed in three phytogeographical regions), M (plant distributed in the Mediterranean region), IT (plant distributed in the Irano-Turanian region), COS (Cosmopolitan, refers to a plant with a worldwide distribution), SCOS (Sub cosmopolitan, a plant distributed on most continents but not all).

## RESULTS AND DISCUSSION

### Flora

As a result of our fieldwork, the vascular flora of Saqalaksar forest contains a total of 94 taxa from 46 families and 80 genera (Appendix 1). Angiosperms include Eudicots with 72 species, 61 genera and 34 families, followed by monocots with 17 species, 15 genera and 8 families. While Monilophytes (Pteridophytes) represent 5 species, 4 genera and 4 families (Table 1). In terms of the number of genera, the richest families were Rosaceae and Poaceae (7 genera), Asteraceae (6 genera), Fabaceae and Lamiaceae (5 genera; Fig. 3). The genus with the highest number of species was *Trifolium* with 4 species, *Carex* and *Rubus* each with 3 species, *Asplenium*, *Crataegus*, *Geranium*, *Mentha*, *Ranunculus*, *Solanum* and *Viola* with 2 representatives each, and the remaining 70 genera with 1 species. The floristic composition of the region indicated the predominance of angiosperms and the ratio of Eudicots (76.60%) to Monocots (18.08%) = 4: 1, which is typical for the flora of Iran and Central Asia (Kamakhina 1994; Memariani *et al.* 2016). Pteridophytes (5 species, 5.32%), were the least diverse group in the region. The richest families of vascular plants in terms of number of species were Rosaceae with 10 species, Fabaceae with 8 species, Poaceae with 7 species, Asteraceae and Lamiaceae with 6 species (Fig. 3). The presence of Rosaceae as the largest family has been reported in previous studies in lowland and mountain forests (Fallah *et al.* 2009; Siadati *et al.* 2010; Ghahremaninejad *et al.* 2011). The abundance of Rosaceae species can be due to geographical conditions, their impacts on the region and the Hyrcanian climate (Razavi & Abbasi 2009), while presence of Poaceae indicates the role of habitat diversity as well as developed soils in the species richness of grasses and Asteraceae, which consist mainly of xerophytic species (Memariani *et al.* 2016). In this study, the results indicated high plant diversity in a limited area. Moreover, comparison of these results with some previous studies in Hyrcanian lowland forest such as Gisum (Ravanbakhsh & Amini 2012), Sedtrik Rudbar (Haghgooy & Pourbabaei 2012) and Bibi Yanloo Astara (Salahi *et al.* 2018), show similarities in the floristic composition (Table 3).

**Table 1.** Number of families, genera and species of main plant groups of Saqalaksar.

Plant Groups	Families	Genera	Species
Eudicots	34	61	72
Monocots	8	15	17
Monilophytes	4	4	5
Total	46	80	94

### Life form spectrum

The life spectrum of plants provides information about their adaptation to environmental conditions, which leads to a classification based on habitat conditions (Archibold 1995; Memariani *et al.* 2016). It can also describe how plants adapt morphologically to climatic factors (Kent & Coker 1995; Asri 2011). Each plant species has a different biological spectrum, which forms the basic structure of the plant communities (Mobayen 1985). In the

assessment of life forms, geophytes dominated by 28 species (29.79%), followed by therophytes by 25 species (26.60%), phanerophytes by 23 species (24.47%), hemicryptophytes by 13 species (13.82%), hydrophytes by 3 species (3.20%) and helophytes by 2 species (2.12%; Fig. 4). A precise classification of the phanerophytes also shows that microphanerophytes are represented by 10 species (10.62 %), megaphanerophytes by 5 species (5.30 %), nanophanerophytes by 4 species (4.25 %), lianas by 3 species (3.20 %) and mesophanerophytes by 1 species (1.1 %).

**Table 2.** Number of families, genera and species according to APG IV (2016).


		Family	Genus	Species		
Monocots	Commelinids	Poales	3	9	11	
		Alismatales	3	3	3	
		Asparagales	1	2	2	
		Liliales	1	1	1	
Angiosperms	Ceratophyllales		1	1	1	
		Ranunculales	1	1	2	
	Eudicots	Fabids	Fabales	1	5	8
			Rosales	3	9	12
			Fagales	3	4	4
		Malvids	Oxalidales	1	1	1
			Malpighiales	3	4	5
			Geraniales	1	1	2
			Myrtales	1	1	1
			Malvales	1	1	1
Sapindales			2	2	2	
Superrosids			Saxifragales	1	1	1
Campanulids	Superasterids	Caryophyllales	2	3	3	
	Asterids	Ericales	1	3	3	
		Aquifoliales	1	1	1	
	Lamiids	Asterales	1	6	6	
		Apiales	2	3	3	
		Dipsacales	1	1	1	
		Solanales	2	3	4	
		Lamiales	3	8	9	
	Gentianales	2	2	2		
		Total	42	76	89	

**Table 3.** Comparison of floristic richness of Saqalaksar forest with Gisoum (Ravanbakhsh & Amini 2012), Sadetarik Rudbar (Haghgooy & Pourbabaei 2012) and Bibi Yanlou Astara (Salahi *et al.* 2018).

Number of plants	Saqalaksar	Sadetarik Rudbar	Bibi Yanlou	Gisoum
Number of species	94	90	92	76
Number of genera	80	81	79	66
Number of Family	46	46	48	45

The high proportion of geophytes in the study area is mainly due to the long wet period during the growing season and the relatively high annual precipitation (Danin & Orshan 1990). This is consistent with the results of floristic studies in other forest areas of Hyrcanian province (e.g. Akbarinia *et al.* 2004; Ghahreman *et al.* 2006; Razavi 2008; Bazdid Vahdati *et al.* 2014). While therophytes are second only to geophytes in the study area, a high percentage of this life form compared to geophytes indicates a relatively strong degradation in this area. Due to moisture and anthropogenic activities, they grow everywhere in the region, especially around the dam. They seem to have been adapted to dry habitats by completing their annual life cycle (Memariani *et al.* 2016). Although therophytes are found in large numbers in desert areas (Archibold 1995), a high percentage of this life form is also found in lowland areas, as most areas are damaged by anthropogenic activities and overgrazing (Naqinezhad *et al.* 2010-2011; Ghahremaninejad *et al.* 2011; Bazdid Vahdati *et al.* 2014). Regarding the relationship between

Rosaceae and their destructive factors in the region, it is noted that the edible fruits of Rosaceae can attract other factors such as humans and animals to the region and consequently cause soil erosion and displacement.



**Fig. 3.** The richest families in terms of species.

**Table 3.** Comparison of floristic richness of Saqalaksar forest with Gisoum (Ravanbakhsh & Amini 2012), Sadetarik Rudbar (Haghgooy & Pourbabaei 2012) and Bibi Yanlou Astara (Salahi *et al.* 2018).

Number of plants	Saqalaksar	Sadetarik Rudbar	Bibi Yanlou	Gisoum
Number of species	94	90	92	76
Number of genera	80	81	79	66
Number of Family	46	46	48	45

Therefore, they can create favourable conditions for the growth of therophytes in the region. Phanerophytes are the predominant life form after therophytes. The abundance of phanerophytes could be due to the Mediterranean microclimate and topographic features of the region (Duran 2002). For the establishment of phanerophytes, it is necessary to provide the resources and vegetation conditions of the region and to prevent and protect them from destruction in the region (Vaseghi *et al.* 2008). In Noor Forest Park, phanerophytes were introduced by 33 % of the largest life form spectrum in the area (Yousefvand *et al.* 2017). The percentage of life forms varies from climate to climate, and the spectrum extracted from each region reflects the climate (Raunkiaer 1934; Mueller-Dombois & Ellenberg 1974). In temperate regions, the predominant species are hemicryptophytes (Raunkiaer 1934), however, due to degradation in the region, therophytes have a high proportion, which is consistent with previous studies (Grime 2001; Bazdid vahdati *et al.* 2014). Hydrophytes are adapted to life in water and are found in areas such as ponds, rivers, streams, lakes, wetlands and other aquatic environments. They develop certain characteristics (adaptations) to ensure their survival in the aquatic environment. Of the three hydrophytes identified, *Ceratophyllum demersum*, *Lemna minor* and *Stuckenia pectinata* were fully submerged and anchored to the substrate. *L. minor*, on the other hand, is free-floating and has a strand of root (Tansley 1949; Penfound 1952; Spence 1964; Sculthorpe 1967). *Ludwigia palustris* (L.) Elliott and *Alisma gramineum* Lej. are two helophytes that occur in this area. They are aquatic macrophytes (Penfound 1952) that root in shallow water and rise above the water surface (Sender & Grabowski 2016). In general, helophytes occur in the transition zone between aquatic and terrestrial environments (Hernández & Rangel 2009; Tavakoli *et al.* 2014). For example, the amphibious nature and high morphological ductility of *Ludwigia* allow it to withstand a wide range of environmental conditions and efficiently colonize aquatic environments (especially rivers, ditches and canals, natural lakes, ponds, wetlands and artificial lakes; Thouvenot *et al.* 2013). In addition, the comparison between the life forms in the forest of Saqalaksar and other forests in Northern Iran is also shown in Fig. 5 (Gahremaninejad *et al.* 2011; Bazid Vahdati *et al.* 2014; Akhondnejad *et al.* 2016). A high proportion of geophytes can be found in Saqalaksar, Ata-Kuh (Southeastern of Lahijan, Gilan province) and Izdeh-e Noor (Noor, Mazandaran province). These concentrations appear to be the most similar to the natural structure and flora of the forests of the Hyrcanian Plain (Zohary 1973; Rastin 1983). Therophytes are present in Semeskandeh and Dasht-e Naz (South of Sari, Mazandaran province) (Gahremaninejad *et al.* 2011). The therophytes in Semeskandeh and Dasht-e Naz show the peak of occurrence (Gahremaninejad *et al.* 2011) and the high percentage of therophytes

in these two localities is probably due to some reasons, disturbance and human activities. These results are also consistent with the studies of Grime (2001).




Fig. 4. Life form spectrum of plants studied in Saqalaksar forest.

### Chorotype spectrum

In terms of geographical distribution, the flora of the studied area is mainly composed of PL elements (34 species, 36.17%), followed by ES (22 species, 23.40%), ES-M-IT (12 species, 12.76%), SCOS (8 species, 8.51%), ES-IT (7 species, 7.44%), COS (6 species, 6.37%), ES-M (4 species, 4.25%) and M elements (1 species, 1.1%) respectively (Fig. 6). Each plant species has a unique ecological range and tolerates a certain degree of ecological change. Therefore, floristic research and study of the geographical distribution of plant species in these areas serve to better understand the potential of the environment and also to manage and protect genetic resources (Naqinezhad *et al.* 2015). Based on the information available, the pluriregional element exhibited the largest geographical distribution. In the classification, geographic distribution refers to pluriregional elements that occur in more than four regions. Plants that have spread in two or more geographical areas, account for about 19% of the flora of Iran (Zohary 1973). According to Naqinezhad *et al.* (2006), water stations (e.g. the areas around the earth dam) with high humidity provide the necessary conditions for the establishment of pluriregional elements. In addition, some researchers confirmed the occurrence of pluriregional elements affected by both water resources and destructive human activities (Gahreman *et al.* 2006; Naqinezhad *et al.* 2008). In explaining the pluriregional elements, it is assumed that they mainly rely on secondary habitats with particular soil conditions such as swamps, marshes, sand dunes and rocks (Mehrabian *et al.* 2009). Euro-Siberian elements were the predominant chorotype after PL, demonstrating the association of this forest with European-Siberian forests (Gahreman *et al.* 2006; Siadati *et al.* 2010). The phytogeographic elements were compared with those of other Caspian lowland forests (Gahremaninejad *et al.* 2011; Bazdid Vahdati *et al.* 2014; Akhondnejad *et al.* 2016; Fig. 7). Two peaks can be observed in the phytochoria spectrum, one in the pluriregional and the other in the Euro-Siberian elements. Phytogeographic elements such as PL and ES exhibit high variation between sites, while ES-M-IT, ES-M, COS, SCOS, M, IT-M-PON and M-IT do not display much variation between sites. The highest proportion of pluriregional elements was found in Saqalaksar, while the lowest proportion in the forests of Izdeh-e Noor. This is due to the fact that the faces of this site are more affected by anthropogenic activities. These results are also consistent with the studies of Ghahremaninejad *et al.* (2011). In addition, Ata-Kuh has the highest proportion of Euro-Siberian elements. Our results suggest that the Ata-Kuh forest has the highest correspondence to a common Euro-Siberian forest due to the large number of Euro-Siberian elements.


### Hyrceanian endemics

These are species occur exclusively in the Hyrcanian geographical area. From a floristic point of view, the occurrence of 44% of all plant species in Iran (3234 out of 7300 species) on only 6% of Iranian surface area (Akhani *et al.* 2010) is very significant. The endemic species of each region are considered the most important species of that region, since their unique occurrence creates a rare diversity compared to other regions where they are more common, indicating the special ecological conditions and genetic diversity of the region. The Saqalaksar forest comprises about 11 species endemic/subendemic to the Hyrcanian province. Among them, *Gleditsia caspica*, *Quercus castaneifolia*, *Parrotia persica*, *Acer velutinum* comprise the best known Hyrcanian endemic trees (Akhani *et al.* 2010; Alavi *et al.* 2020), while *Ilex spinigera*, *Hedera pastuchovii* and *Ruscus hyrcanus* are evergreen species that grow in most of these forests (Rastin 1983).


### CONCLUSION

The study of the vegetation in the region shows a great ecological diversity, including aquatic and terrestrial habitats, most of which are forested areas. Most of this biodiversity is only found in a small part of the dam. This


demonstrates the importance of vegetation in marginal areas and prevents further destruction and conservation of the remaining elements.



**Fig. 5.** Variation of each life form over the sites Ata-Kuh forest (Bazdid vahdati *et al.* 2014); Semeskandeh and Dasht-e Naz forests (Gahremaninejad *et al.* 2011); Iz = Izdeh-e Noor forest (Akhondnejad *et al.* 2016). Abbreviations: (Cha = Chamaephyte, Geo = Geophyte, Hel = Helophyte, Hem = Hemicryptophyte, Hyd= Hydrophyte, Pha = Phanerophyte, Par = Parasite, Thr = Therophyte).



**Fig. 6.** Proportion of different phytochoria elements of plants in Saqalaksar forest. Abbreviations (ES= Euro-Siberian, PL= Pluriregional, M= Mediterranean, IT= Irano-Turanian, COS= Cosmopolitan, SCOS= Subcosmopolitan).



**Fig. 7.** Variation of each phytochoria over the sites. Abbreviations: Saq= Saqalaksar, Ata = Ata-kuh, Sem & Dash = Semeskandeh and Dasht-e Naz, Iz = Izdeh-e Noor, ES = Euro-Siberian, PL= Pluriregional, M = Mediterranean, IT = Irano-Turanian, COS = Cosmopolitan, SCOS = Subcosmopolitan.

As a number of species are only located in one releves, the focus can be placed on protecting and managing the vegetation in this area. In addition, unregulated tourism activities can cause serious damage to the vegetation. Therefore, the threat category of the plant species should be considered according to the IUCN categories and criteria.

**ACKNOWLEDGEMENT**

This research was supported by the project of the University of Guilan, Rasht, Iran. The authors would like to thank Dr. Asghar Zamani and Ali Sarvi in University of Guilan for help in this study.

## REFERENCES

- Aghajani, MS, Oskoueian, R, Naqinezhad, A, Asadi, M, Serpooshan, F 2023, A floristic study of Garasmasar Mountains: Mazandaran Province, Iran. *Caspian Journal of Environmental Sciences*, 21: 693-709.
- Akbarinia, M, Zare, H, Hosseini, SM & Ejtehad, H 2004, Study on vegetation structure, floristic composition and chorology of silver birch communities at Sangdeh. *Pajouhesh and Sazandegi*, 17: 84-96, [In Persian].
- Akhani, H 1998, Plant Biodiversity of Golestan National Park, Iran. *Stapfia*, 53: 1-411.
- Akhani, H, Djamali, M, Ghorbanalizadeh, A & Ramezani, E 2010, Plant biodiversity of Hyrcanian Relict Forests, N Iran: An overview of the flora, vegetation, palaeoecology and conservation. *Pakistan Journal of Botany*, 42: 231-258.
- Akhondnejad, S, Asri, Y & Khakpour Moghaddam, T 2016, Introduction of the flora, life form and chorology of the *Parrotia persica* Habitats (Case study: Izdeh-e Noor area). *Taxonomy and Biosystematics*, 8: 103-120 [In Persian].
- Alavi, SJ, Veiskarami, R, Esmailzadeh, O & Gadow, KV 2020, Analyzing the Biological and Structural Diversity of Hyrcanian Forests Dominated by *Taxus baccata* L. *Forests*, 11:701.
- APG IV 2016, An update of the angiosperm phylogeny group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society*, 181: 1-20.
- Archibold, OW 1995, Ecology of World Vegetation. Chapman & Hall press, London, 510 p.
- Asadi, H, Hosseini, SM, Esmailzadeh, O & Ahmadi, A 2011, Flora, life form and chorological study of Box tree (*Buxus hyrcanus* Pojark.) sites in Khybus protected forest, Mazandaran. *Iranian Journal of Plant Biology*, 3: 27-40 [In Persian].
- Assadi, M, Maassoumi, A, Khatamsaz, M & Mozaffarian, V 1988–2016, Flora of Iran. Research Institute of Forests and Rangelands Publication, Tehran (in Persian).
- Assadollahi, F 1980, Phytosociological and biogeographic study of Hyrcanian forests. Synthetic test and application to the Assalem region (Iran). Dissertation, Marseille University, Marseille, France, 127 p., [In French].
- Ataei, A, Kazemnezhad, F, Nimvari, ME & Sheykhholeslami, A 2021, Ecotones and forest communities along an elevation gradient in Hyrcanian forests, north of Iran. *Caspian Journal of Environmental Sciences*, 19: 317-323.
- Asri, Y 2011, Phytosociology. Payame Noor University, Tehran, Iran, 1:6-22.
- Atashgahi, Z, Ejtehad, H & Zare, H 2009, Study of floristics, life form and chorology of plants in the east of Dodangeh forests, Mazandaran province, Iran. *Iranian Journal of Biology*, 22: 193-203 [In Persian].
- Bailey, RG 2003, Ecological Climate Classification. *USDA Forest Service, Inventory & Monitoring Institute*.
- Barans, ACP 1969, Vascular flora of the college woods. The College of William and Mary, James CityCounty, Virginia, 206 pp.
- Bazdid Vahdati, F, Saeidi Mehrvarz, Sh, Naqinezhad, A & Shahi Shavvon, R 2014, Floristic characteristics of the Hyrcanian submountain forests (Case study: Ata-Kuhforest). *Caspian Journal of Environmental Sciences*, 12: 169-183.
- Bazdid vahdati, F, Saeidi Mehrvarz, Sh, Dey, D.C & Naqinezhad, A 2017, Environmental factors–ecological species group relationships in the Surash lowland-mountain forests in northern Iran. *Nordic Journal of Botany*, 35: 240-250.
- Cain, SA 1938, The species–area curve. *The American Midland Naturalist Journal*, 19: 81-573.
- Chiarucci, A & Bonini, I 2005, Quantitative floristics as a tool for the assessment of plant diversity in Tuscan forests. *Forest Ecology and Management*, 212: 160-170.
- Danin, A & Orshan, G 1990, The distribution of Raunkiaer life forms in Israel in relation to the environment. *Journal of Vegetation Science*, 1: 41-48.
- Davis, PH 1965-1985, Flora of Turkey. Edinburgh University Press, Edinburgh, pp. 1-9.
- De Martonne, E 1926, A new climatological function: The aridity index. *Meteorology*, pp. 449-458, (In French).
- Djamali, M, Akhani, H, Khoshravesh, R, Andrieu-Ponel, V, Ponel, P & Brewer, S 2011, Application of the global bioclimatic classification to Iran: implications for understanding the modern vegetation and biogeography. *Ecologia Mediterranea*, 37: 91-114.
- Djazirei, MH 1964, Contribution to the study of the Hyrcanian forest. University Faculty of Agricultural Sciences of Gembloux, Gembloux, Belgium [In French].
- Djazirei, MH 1965, Contribution to the study of primary forests of the Caspian. *Bulletin of the Agronomic Institute of Gembloux*, 33: 35-75 [In French].
- Dorostkar, H 1974, Contribution to the study of the forests of the eastern Hyrcanian district (Gorgan Range). University Faculty of Agricultural Sciences of Gembloux, Gembloux, Belgium, (In French).
- Duran, A 2002, A new species of *Scorzonera* L. (Asteraceae) from central Anatolia, Turkey. *Israel Journal of Plant Sciences*, 50: 155-159.
- Emberger, E 1932, On a climatic formula and its applications in botany. *Meteorology*, pp. 423-432.
- Fallah, F, Salimpour, F & Sharifnia F 2009, Flora, life form and chorological a part of Savadkouh, Mazandaran province. *Iranian Journal of Biological Sciences*, 4: 29-42 [In Persian].
- Frey, W & Probst, W 1986, Asynopsis of the vegetation of Iran. In: Contributions to the vegetation of southwest Asia. (Ed. Kürschner, H.), Dr. Ludwig Reichert, Wiesbaden, pp.1-43.
- Ghahreman, A & Attar, F 1998, Biodiversity of plant species in Iran. Tehran University Publications, Iran, [In Persian].
- Ghahreman, A 1979–2005, Coloured Flora of Iran. Research Institute of Forests and Rangelands Press, Tehran, Vols. 1-

- 26, [In Persian].
- Ghahreman, A, Naqinezhad, A, Hamzeh'ee, B, Attar, F & Assadi, M 2006, The flora of threatened black alder forests in the Caspian lowlands, Northern Iran. *Rostaniha*, 7: 5-30.
- Ghahremaninejad, F, Naqinezhad, A, Bahari, SH & Esmaeili, R 2011, An introduction to flora, life form, and distribution of plants in two protected lowland forests, Semeskandeh and Dasht-e Naz, Mazandaran N. Iran. *Taxonomy and Biosystematic*, 3: 53-70, [In Persian].
- Grime, JP 2001, Plant strategies, vegetation processes and ecosystem properties. John Wiley and Sons, New York. 417 p.
- Haghooy, T & Pourbabaie, H 2012, Presentation of flora, life form and chorotype of plants in Sadetarik Forest Park, Roudbar, Guilan. *Ecology of Iranian Forests*, 3: 331-340, [In Persian].
- Hamzeh'ee, B, Naqinezhad, A, Attar, F, Ghahreman, A, Assadi, M & Prieditis, N 2008, Phytosociological survey of remnant *Alnus glutinosa* ssp. *barbata* communities in the lowland Caspian forests of northern Iran. *Phytocoenologia*, 38: 117-132.
- Harding, AE, Gachon, P & Nguyen, VTV 2011, Replication of atmospheric oscillations, and their patterns, in predictors derived from Atmosphere–Ocean Global Climate Model output. *International Journal of Climatology*, 31: 1841-1847.
- Hernández-R, J & Rangel-Ch, JO 2009, Vegetation of the wetland Jaboque (Bogotá, DC). *Caldasia*, 31: 355-379.
- IPNI 2020, International Plant Names Index. Published on the Internet <http://www.ipni.org>, The Royal Botanic Gardens, Kew, Harvard University Herbaria & Libraries and Australian National Botanic Gardens (accessed 5 June 2020).
- IUCN 2020, Standards and Petitions Committee. 2019. Guidelines for Using the IUCN Red List Categories and Criteria. Version 14 (August 2019). Prepared by the Standards and Petitions Committee. Available from: <http://www.iucnredlist.org/documents/RedListGuidelines> (accessed April 24, 2020).
- Kamakhina, GL 1994, Kopetdagh-Khorassan flora: regional features of central Kopetdagh. *Biogeography and ecology of Turkmenistan*, pp. 129-148.
- Kent, M & Coker, P 1995, Vegetation Description and Analysis: a practical approach. John Wiley press, New York.
- Klein, JC 2001, The altitudinal vegetation of Central Alborz (Iran): between the Iranian-Turanian and Euro-Siberian regions. French Institute of Research in Iran, Tehran.
- Knapp, HD 2005, The global importance of the Caspian forests. In K, Nostrari, R, Marvie Mohadjer, W, Bode & HD Knapp: Protection of biological diversity and integrated management of the Caspian forests (Northern Iran). Federal Office for Nature Conservation, pp. 45-70. (In German).
- Kudryavtsev, NA, Zaitseva, LA, Savoskina, OA, Chebanenko, SI, Zavertkin, IA 2021 Herbological and agrotechnological approaches to weeding plants in modern flax growing. *Caspian Journal of Environmental Sciences*, 19: 903-908.
- Léonard, J 1988, Contribution to the study of the flora and vegetation of the deserts of Iran, Paper 8: Study of distribution areas, phytocories, chorotypes. *Bulletin du Jardin Botanique National de Belgique*, pp. 1-190.
- Mehrabian, A, Naqinezhad, A, Mahiny, A, Mostafavi, H, Liaghati, H & Kouchezkadeh, M 2009, Vegetation Mapping of the Mond Protected Area of Bushehr Province (South-west Iran). *Journal of Integrative Plant Biology*, 51: 251-260.
- Memariani, F, Zarrinpour, V & Akhiani, H 2016, A review of plant diversity, vegetation and phytogeography of the Khorasan-Kopet Dagh floristic province in the Irano-Turanian region (northeastern Iran – southern Turkmenistan). *Phytotaxa*, 249: 8-30.
- Mirhashemi, H, Moradi, F, Pourbabaie, H & Mezban, F 2021, Changes in flora, life forms and geographical distribution of herbaceous plant species along an altitudinal gradient in oak forests, Iran. *Caspian Journal of Environmental Sciences*, 19: 619-627.
- Mobayen, S & Tregubov, V 1970, Map of natural vegetation of Iran, 1:2,500,000. University of Tehran, Tehran.
- Mobayen, S 1985, Flora of Iran, vascular plants. Tehran University Press, Tehran, [In Persian].
- Mueler-Dombois, D & Ellenberg, D 1974, Aims and Methods of Vegetation Ecology. John Wiley press, New York, 547 p.
- Muratovna, ZT, Iskakovich, SK, Duisengalievna, MA, Kunsalievna, MB, Nurakhmetovna, AR & Aigul, B 2024, Flora of the gorges Merke, Sandyk, Shaisandyk in the western part of Kyrgyz Alatau, a natural border between Kyrgyzstan and Kazakhstan. *Caspian Journal of Environmental Sciences*, 22: 59-69.
- Najafi TSK, Jalili, A, Khorasani, NA, Jamzad, Z & Asri, Y 2008, Investigation on relationship between ecological factors and plant associations of geno protected area. *Iranian journal of Range and Desert Research*, 15: 179-199.
- Naqinezhad, A, Saeidi Mehrvarz, Sh, Noroozi, M & Faridi, M 2006, Contribution to the vascular and bryophyte flora as well as habitat diversity of the Boujagh National Park, N. Iran *Rostaniha*, 7: 83-105.
- Naqinezhad, A, Hamzeh'ee, B & Attar, F 2008, Vegetation-environment relationships in the alder wood communities of Caspian lowlands, N. Iran (toward an ecological classification). *Flora*, 203: 567-577.
- Naqinezhad, A, Shahsavari, A & Jahanian, T 2009, Study of plant biotypes in Babol from the perspective of urban ecology. *Journal of Ecology*, 35: 89-104, [In Persian].
- Naqinezhad, A, Hosseini, S, Rajamand, M & Saeidi Mehrvarz, Sh 2010, A floristic study on Mazibon and Sibon protected forests, Ramsar, across the altitudinal gradient (300-2300 m). *Taxonomy and Biosystematics*, 2: 93-114, [In Persian].
- Naqinezhad, A, Ghahremaninejad, F & Bahari, H 2011, *Monerma* P. Beauv. (Gramineae) a new monotypic genus record for the Flora of Iran. *The Iranian Journal of Botany (IJB)*, 17: 75-77.

- Naqinezhad, A, Zare-Maivan, H & Gholizadeh, H 2015, A floristic survey of the Hyrcanian forests in Northern Iran, using two lowland-mountain transects. *Journal of Forestry Research*, 26: 187-199.
- Pears, N 1985, Basic Biogeography. Pearson Education, New York, 368 p.
- Penfound, WT 1952, Southern swamps and marshes. The Botanical Review, Springer, 18: 413-446.
- Rastin, N 1983, Vegetation science studies in high forest remnants of the Caspian Plain. *Phytocoenology*, 11: 245-289.
- Raunkiaer, C 1934, The life forms of plants and statistical plant geography. Oxford University Press, Oxford.
- Ravanbakhsh, M & Amini, T 2012, Study on floristic composition, chorology and ecological structure of Gisoum forest reserve, Talysh, Iran. *Iranian Journal of Biology*, 25: 21-31, [In Persian].
- Razavi, SA 2008, Flora study of life forms and geographical distribution in Kouhmian region (Azadshahr-Golestan Province). *Journal of Agricultural Sciences and Natural Resources*, 15: 98-108 (In Persian).
- Razavi, SA & Abbasi, N 2009, A floristic and chorology investigation of oriental arborvitae in Sourkesh reserve (Fazel Abad-Golestan province). *Journal of Wood and Forest Science and Technology*, 16: 83-100, [In Persian].
- Rechinger, KH 1963–2015, Flora Iranica. Akademische Druck-U. Verlagsanstalt, Graz, Vols. 1-178.
- Rivas-Martinez, S, Asensi, A, Diez-Garretas, B, Molero, J & Valle, F 1997, Biogeographical synthesis of Andalusia (Southern Spain). *Journal of Biogeography*, 24: 915-928.
- Rivas-Martinez, S, Mata, D.S & Costa, M 1999, North American Boreal and Western temperate forest vegetation: (Syntaxonomical synopsis of the potential natural plant communities of North America, II). *Itinera Geobotanica*, 12: 5-316.
- Sagheb-Talebi, K, Pourhashemi, M & Sajedi, T 2014, Forests of Iran: A Treasure from the Past, a Hope for the Future. Springer.
- Salahi, T, Pourbabaei, H, Salahi, M & Karamzadeh, S 2018, Study on Floristic Composition and Chorology of Bibi yanlou's Forest Park, Astara. *Ecology of Iranian Forests*, Vol: 6, No: 11, [In Persian].
- Scharnweber, T, Rietschel, M & Manthey, M 2007, Degradation stages of the Hyrcanian forests in southern Azerbaijan. *Archiv für Naturschutz und Landschaftsforschung*, 46:133-156.
- Sculthorpe, CD 1967, The biology of aquatic vascular plants. Edward Arnold, London.
- Sender, J & Grabowski, M 2016, The relationship between land management and the nature of helophytes in small lakes (Eastern Poland). *Limnological Review*, 16: 51-62.
- Siadati, S, Moradi, H, Attar, F, Etemad, V, Hamzeh'ee, B & Naqinezhad, A 2010, Botanical diversity of Hyrcanian forests; a case study of a transect in the Kheyroud protected lowland mountain forests in northern Iran. *Phytotaxa*, 7: 1-18.
- Spence, DH 1964, The macrophytic vegetation of freshwater lochs, swamps and associated fens. In: The vegetation of Scotland. (Ed. JH, Burnett), Edinburgh pp. 306-425.
- Takhtajan, A 1986, Floristic Regions of the World. University of California Press, Berkley, 522 p.
- Tavakoli, MT, Chenari, AI, Rezaie, M, Tavakoli, A, Shahsavari, M & Mousavi, SR 2014, The importance of micronutrients in agricultural production. *Advances in Environmental Biology*, 8: 31-36.
- Thouvenot, L, Haury, J & Thiébaud, G 2013, Seasonal plasticity of *Ludwigia grandiflora* under light and water depth gradients: an outdoor mesocosm experiment. *Flora*, 208: 430-437.
- Tohidifar, M, Moser, M, Zehzad, B & Ghadirian, T 2016, Biodiversity of Caspian forests. Biodiversity of the Hyrcanian Forests: A synthesis report. DOI, 10.
- Tutin, TG, Heywood, VH, Burges, NA, Moore, DM, Valentine, DH, Walters, SM & Webb, DA 1964-1980, Flora Europaea. Cambridge University Press, Cambridge, Vols. 1-5.
- Vaseghi, P, Ejtehadi, H & Zokaei, M 2008, Flora, Life form and Chorological study of Aboveground Plant Elements in Kalat-Zirjan Gonabad, Khorasan Razavi, Iran. *Journal of Science Kharazmi University*, 8: 75-88, [In Persian].
- Yousefi, M 2009, Flora of Iran. payame noor university, Tehran.
- Yousefvand, S, Esmailzadeh, O, Jalali, SG & Asadi, H 2017, Flora, Life Form and Chorological Study of Aboveground Vegetation and Soil Seed Bank in Noor Forest Park. *Journal of plant researches*, 30: 102-114.
- Zohary, M 1973, Geobotanical foundations of the Middle East. Gustav Fischer verlag, Stuttgart, Vols. 1-2.

#### Appendix 1

Floristic list of the Saqalaksar forest.

Symbols and abbreviation used:

1. Life cycle

A = Annual, B = Biennial, P = Perennial

2. Life forms

G = Geophyte, Hel = Helophytes, Hem = Hemicyrptophyte, Hyd = Hydrophyte, Pha = Phanerophyte [PH-M = megaphanerophyte, PH-m = mesophanerophyte, PH-mi = microphanerophyte, PH-n = nanophanerophyte, PH-li = Liana], T = Therophyte.

3. Chorotype

COS = Cosmopolitan, ES= Euro-Siberian (Hyr = Hyrcanian, E-Hyr = Euxino-Hyrcanian, E = Endemic), IT = Irano-Turanian, M = Mediterranean, PL = Pluriregional, SCOS = Subcosmopolitan

Hb. No.= Herbarium number

Taxa      Duration      Life form      Chorotype      Hb. No.

Monilophytes					
Aspleniaceae					
Asplenium adianoum-nigrum L.	P		Hem	PL	8456
Asplenium scolopendrium L.	P		G	PL	8457
Dennstaedtiaceae					
Pteridium aquilinum (L.) Kuhn	P		G	COS	8458
Polypodiaceae					
Polypodium vulgare L.	P		G	PL	8459
Pteridaceae					
Pteris cretica L.	P		G	PL	8460
Angiosperms					
Eudicots					
Adoxaceae					
Sambucus ebulus L.	P		G	ES-IT[M]	8461
Apiaceae					
Centella asiatica (L.) Urb.	A		G	PL	8462
Torilis arvensis (Huds.) Link	A		T	PL	8463
Apocynaceae					
Periploca graeca L.	P		PH-li	ES-IT-M	8464
Aquifoliaceae					
Ilex spinigera (Loes.) Loes	P		PH-mi	EN(ES)(HYR)	8465
Araliaceae					
Hedera pastuchovii Woronow	P		PH-li	EN-(ES)(HYR)	8466
Asteraceae					
Artemisia annua L.	A		T	ES-M-IT	8467
Bidens tripartita L.	A		T	PL	8468
Centaurea hircanica Bornm.	A		Hem	EN-(ES)(HYR)	8469
Cirsium vulgare (Savi) Ten.	P		Hem	PL	8470
Erigeron bonariensis L.	A		T	COS	8471
Eclipta prostrata (L.) L.	A		T	PL	8472
Betulaceae					
Alnus glutinosa Subsp. barbata (C.A. Mey.) Yalt.	P		PH-M	ES(E-Hyr)	8473
Carpinus betulus L.	P		PH-M	ES	8474
Caryophyllaceae					
Stellaria media (L.) Vill.	A		T	SCOS	8475
Ceratophyllaceae					
Ceratophyllum demersum L.	P		Hyd	SCOS	8476
Convolvulaceae					
Calystegia silvatica (Kit.) Griseb.	P		G	ES-M	8477
Cuscuta campestris Yunck.	A		T	PL	8478
Euphorbiaceae					
Acalypha australis L.	A		T	PL	8479
Euphorbia amygdaloides L.	B-P		G	ES	8480
Fabaceae					
Albizia julibrissin Durazz.	P		PH-mi	PL	8481
Gleditsia caspia Desf.	P		PH-mi	EN(ES)(HYR)	8482
Lotus corniculatus L.	A		Hem	ES-M-IT	8483
Trifolium campestre Schreb.	A		T	ES-M-IT	8484
Trifolium fragiferum L.	A		G	PL	8485
Trifolium repens L.	A		G	ES-IT-M	8486
Trifolium resupinatum L.	A		G	ES-IT-M	8487
Vicia sativa L.	P		Hem	PL	8488
Fagaceae					
Quercus castaneifolia C.A.Mey.	P		PH-M	EN(ES)(HYR)	8489
Gentianaceae					
Centaurium erythraea Rafn	P		Hem	PL	8490
Geraniaceae					
Geranium molle L.	A-B		T	ES-IT	8491
Geranium purpureum Vill.	A-B		T	Es-M	8492
Hamamelidaceae					
Parrotia persica C.A.MeyP				EN(ES)(HYR)	8493
Hypericaceae					
Hypericum perforatum L.	P		Hem	SCOS	8494
Juglandaceae					
Pterocarya fraxinifolia (Poir.) Spach	P		PH-m	ES(HYR)	8495
Lamiaceae					
Clinopodium umbrosum (M.Bieb.) Kuntze			P	Hem	PL
Lycopus europaeus L.	P		G	PL	8497
Mentha aquatica L.	P		G	ES	8498
Mentha pulegium L.	P		Hem	ES	8499
Prunella vulgaris L.	P		G	PL	8500
Scutellaria tournefortii Benth.	P		G	EN-(ES)(HYR)	8501
Malvaceae					
Abutilon theophrasti Medik.	A		T	COS	8502
Moraceae					

Ficus carica L.	P	PH-mi	ES-M-IT	8503	
Onagraceae					
Ludwigia palustris (L.) Elliott	P		Hel	PL	8504
Oxalidaceae					
Oxalis corniculata L.	P	T		PL	8505
Plantaginaceae					
Plantago major L.	P	G		PL	8506
Veronica persica poir.	A	T		SCOS	8507
Polygonaceae					
Persicaria lapathifolia (L.) Delarbre	A	T		ES-M-IT	8508
Rumex sanguineus L.	P	Hem		ES	8509
Primulaceae					
Lysimachia arvensis L.	A	T		COS	8510
Cyclamen coum Mill.	P	G		ES(HYR)-IT	8511
Primula heterochroma Stapf	P			Hem	EN-(ES)(HYR) 8512
Ranunculaceae					
Ranunculus arvensis L.	A	T		ES-M-IT	8513
Ranunculus scleratus L.	A	T		PL	8514
Rosaceae					
Crataegus microphylla K.Koch	P			PH-mi	ES-IT 8515
Crataegus pentagyna Waldst. & Kit. ex Willd.				P	PH-mi ES 8516
Fragaria vesca L.	P	G		ES-IT	8517
Malus orientalis Uglitzk. ex Juz.	P			PH-mi	ES-IT 8518
Mespilus germanica L.	P	PH-mi		ES(HYR)	8519
Potentilla reptans L.	P	Hem		ES-IT	8520
Prunus cerasifera Ehrh	P	PH-mi		ES-IT	8521
Rubus caesius L.	P	PH-n		PL	8522
Rubus ulmifolius Schott	P	PH-n		ES(HYR)	8523
Rubus sanctus Schreb.	p	PH-n		PL	8524
Sapindaceae					
Acer velutinum Boiss.	p	PH-M		EN-(ES)(HYR)	8525
Simaroubaceae					
Ailanthus altissima (Mill.) Swingle	p			PH-mi	ES-M-IT 8526
Solanaceae					
Solanum americanum Mill.	A	T		SCOS	8527
Solanum sisymbriifolium Lam.	A	T		SCOS	8528
Urticaceae					
Urtica dioica L.	P	G		PL	8529
Verbenaceae					
Phyla nodiflora (L.) Greene	P	T		PL	8530
Violaceae					
Viola caspia (Rupr.) Freyn	P	G		ES(E-Hyr)	8531
Viola odorata L.	P	G		PL	8532
Monocot					
Araceae					
Lemna minor L.	P	Hyd		PL	8533
Alismataceae					
Alisma gramineum Lej.	A	Hel		PL	8534
Asparagaceae					
Ornithogalum sintenisii Freyn	P	G		EN-(ES)(HYR)	8535
Ruscus hyrcanus Woronow	P	PH-n		EN-(ES)(HYR)	8536
Cyperaceae					
Carex divulsa Stokes	P	G		ES-M-IT	8537
Carex grioletii Roem. ex Schkuhr	P	G		ES-M	8538
Carex sylvatica Huds.	P	G		SCOS	8539
Juncaceae					
Juncus acutus L.	P	G		SCOS	8540
Poaceae					
Periballia laevis (Brot.) Asch. & Graebn.				A	T M 8541
Cynodon dactylon (L.) Pers.	P			G	PL 8542
Digitaria sanguinalis (L.) Scop.	A			T	PL 8543
Echinochloa crus-galli (L.) P.Beauv.	A			T	PL 8544
Oplismenus undulatifolius (Ard.) Roem. & Schult	P			G	PL 8545
Paspalum distichum L.	P	Hem		COS	8546
Poa annua L.	A	T		PL	8547
Potamogetonaceae					
Stuckenia pectinata (L.) Börner	P			Hyd	COS 8548
Smilacaceae					
Smilax excelsa L.	P	PH-Li		ES(HYR)-M	8549

**Bibliographic information of this paper for citing:**

Ajamian, M, Beygom Faghir, M, Bazdid Vahdati, F 2024, Floristic, life form and chorological studies of the Saqalaksar forest, Rasht, Northern Iran. Caspian Journal of Environmental Sciences, 22: 277-288.