

# Protective effects of 1, 8-cineole (eucalyptol) against CCl<sub>4</sub> induced hepatic oxidative damage in rats

Mehriban S. Baratova<sup>1</sup>, Ilhom A. Ulashov<sup>2\*</sup>, Zukhra Yakhshieva<sup>3</sup>, Alimardon A. Egamberdiev<sup>4</sup>, Botirbek Xoshimov<sup>5</sup>, Umid A. Kodirov<sup>6</sup>, Ajiniyaz A. Eshmuratov<sup>7</sup>, Klara X. Avilova<sup>8</sup>, Ikromjon I. Mamadoliev<sup>9</sup>, Davlat Normurodov<sup>10</sup>

1. Candidate of Medical Sciences, Associate Professor - Head of the Department of Public Health, Healthcare Management, Bukhara State Medical Institute, Uzbekistan

2. PhD in Veterinary Sciences, Senior Researcher at the Helminthology Laboratory, Veterinary Scientific Research Institute, Taylok region, Samarkand, Uzbekistan

3. Doctor of Chemical Sciences, Professor, Jizzakh State Pedagogical University, Jizzakh, Uzbekistan

4. Assistant, Department of "Engineering Graphics and Design Theory", "Tashkent Institute of Irrigation and Agricultural Mechanization Engineers", National Research University, Tashkent, Uzbekistan

5. Doctor of Philosophy (PhD Veterinary Science), Department of Agronomy, Navoi State University of Mining and Technologies, Navoi, Uzbekistan

6. Teacher at the Department of Neurology, Faculty of Postgraduate Education, Samarkand State Medical University, Samarkand, Uzbekistan

 Lecturer, Department of Art Science, Karakalpak State University named after Berdakh, Nukus, Republic of Karakalpakstan
 Senior Teacher, Department of Pharmaceuticals and Chemistry, Faculty of Medicine, Alfraganus University, Tashkent, Uzbekistan

9. Kimyo International University in Tashkent, Tashkent, Uzbekistan

10. Professor Institute of Agrobiotechnology and Food Securite of Samarkand State University named after Sharof Rashidov, Jamboy district, Samarkand, Uzbekistan

\* Corresponding author's Email: ulashovilhom14@gmail.com

## ABSTRACT

Carbon tetrachloride ( $CCl_4$ ) is a hepatotoxin which induce oxidative stress. Therefore, supplementation with antioxidants may be beneficial in mitigating its hepatotoxic effects. 1,8-Cineole (1,3,3-trimethyl-2-oxabicyclo [2.2.2] octane), commonly referred to as euclyptol has various therapeutic properties. Hence, this research was conducted to evaluate the protective effects of eucalyptol against CCl<sub>4</sub> induced hepatic oxidative damage in rats. Thirty rats were divided randomly into the following three equal groups (n = 10): healthy control, CCl<sub>4</sub>-poisoned group (1 mL/kg BW dissolved in olive oil (1:1 volume) and eucalyptol treated group (100 mg/kg/day). After treatment period (14 days), serum sample was prepared and liver was taken out for measurement of biochemical parameters including liver enzymes such as Alanine transaminase (ALT) and Aspartate transaminase (AST) as well as oxidative stress markers including Malondialdehyde (MDA), Glutathione (GSH), Catalase (CAT), Superoxide dismutase (SOD) and Glutathione Peroxidase (GPX). The results showed that following CCl<sub>4</sub> poisoning, a significant reduction in the levels of GSH, GPX, CAT and SOD was observed in the CCl<sub>4</sub>-poisoned group compared to the control, as well as a significant elevation in the levels of AST, ALT and MDA was observed in the CCl<sub>4</sub>-poisoned group in comparison with the control (p < 0.05). Furthermore, there was a significant increase in the levels of GSH, GPX, CAT and SOD in the eucalyptol treated group compared to the CCl<sub>4</sub>-poisoned group. However, there was a significant decrease in the levels of AST and ALT in the eucalyptol-treated group compared to the CCl<sub>4</sub>-poisoned group (p < 0.05). However, there was no significant in the MDA level in the eucalyptol-treated group compared to the CCl<sub>4</sub>-poisoned one (p > 0.05). Eucalyptol may be effective in the

prevention of CCl<sub>4</sub>-induced liver damage through the reduction of oxidative stress, liver damage caused by elevated liver enzymes, and increasing antioxidant enzymes.

Keywords: Eucalyptol, 1,8-Cineole, Antioxidants, Hepatic damage, CCl4, Oxidative stress. Article type: Research Article.

#### INTRODUCTION

The liver, recognized as the largest metabolic organ in the body, plays a critical role in the biotransformation and elimination of synthetic compounds (El Rabey et al. 2021; Parpieva et al. 2023). Environmental toxins have the potential to induce liver injury through a multifaceted mechanism involving oxidative stress, inflammation, necrosis, and apoptosis (Xiong et al. 2020; Ugochukwu Madu et al. 2023; Asadi et al. 2024). Previous studies have implicated oxidative stress, inflammation and toxic compounds in the development of hepatocellular damage (Wahid et al. 2016; Kabbashi et al. 2024). Elevated oxidative stress in the liver is a key factor contributing to the development of hepatic fibrosis (Devi et al. 2021). Carbon tetrachloride (CCl4) is an industrial chemical known for its hepatotoxic properties and remains present as a component in certain insecticides (Vahidi-Eyrisofla et al. 2019; Que et al. 2022). CCl<sub>4</sub> was extensively utilized as a solvent in the chemical industry; however, its usage has been largely discontinued in most countries over the past decade due to its toxic properties (Teschke 2018; Hashemzadeh et al. 2018). Under experimental and laboratory conditions, CCl<sub>4</sub> is one of the most frequently utilized models for studying both acute and chronic liver damage (Kostic et al. 2022). Exposure of humans to CCl<sub>4</sub> has been well documented to result in liver fibrosis and hepatocellular injury (Yeh et al. 2013). The liver is the primary organ for the metabolism of CCl<sub>4</sub>, where it is converted to peroxide radicals. These highly reactive radicals induce oxidative stress, resulting in lipid peroxidation and elevation of liver enzyme levels (Moghadam et al. 2017; Di Paola et al. 2022). Induction of oxidative stress is a primary mechanism underlying CCl<sub>4</sub> hepatotoxicity. Therefore, supplementation with antioxidants could be beneficial in mitigating its hepatotoxic effects (Kabbashi et al. 2024). One of them is 1,8-cineole (1,3,3-trimethyl-2-oxabicyclo [2.2.2] octane), commonly known as eucalyptol. This compound is present in the essential oil fraction of several plant sources, including eucalyptus, rosemary, and camphor (Hoch et al. 2023). Eucalyptus essential oil has the highest concentration of eucalyptol, according to Cai et al. (2021). Owing to its pleasant taste and aroma, eucalyptol is widely used in making food, perfuming and cosmetic products (Hoch et al. 2023). The primary pharmacological properties of eucalyptol are its anti-inflammatory and antioxidant effects. In particular, the production of inflammatory cytokines and reactive oxygen species (ROS) has been shown to be inhibited by eucalyptol (Kennedy-Feitosa et al. 2019). As a result, eucalyptol has therapeutic potential for cardiovascular disease, digestive disorders, Alzheimer's disease, and respiratory disorders. In particular, eucalyptol has been used to treat bronchitis, asthma, and chronic obstructive pulmonary disease (COPD; Cai et al. 2021). Therefore, this study investigates the protective effects of eucalyptol against hepatic oxidative damage induced by CCl<sub>4</sub> in rats.

### MATERIALS AND METHODS

#### Chemicals and reagents

The materials and reagents used in this investigation included biochemical kits for alanine transaminase (ALT) and aspartate transaminase (AST) provided by Sigma-Aldrich Company, USA, and 1, 8-cineole from Sigma-Aldrich Company, USA, CCl<sub>4</sub>, Tris-HCl, thiobarbituric acid (TBA), trichloroacetic acid (TCA), tris-EDTA, dinitrothiocyanobenzene (DTNB) and phosphate-buffered saline (PBS) from Merck Company, Germany.

#### Animals and experimental design

Thirty adult male Wistar rats (two-month olds; weight 250 to 300 g) were purchased and housed in standard caging (n = 10). The rats were maintained under suitable environmental conditions, which comprised a temperature of 22 °C, a 12-hour light-dark cycle, and unrestricted access to standard laboratory food and tap water. The rats were randomly assigned to three experimental groups, with ten rats in each group:

Group 1: Healthy rats (control) were given water daily for two weeks.

**Group 2:** Designated as the poisoned group (CCl<sub>4</sub> group), was exposed to CCl<sub>4</sub> (1 mL kg<sup>-1</sup> body weight) dissolved in olive oil (1:1 volume) via intraperitoneal (IP) injection twice weekly over a period of two weeks.

**Group 3:** The rats were given  $CCl_4$  + eucalyptol by oral gavage at a dose of 100 mg kg<sup>-1</sup> per day for 14 continuous days.

Rats anesthetized with ketamine (50 mg kg<sup>-1</sup>) and xylazine (10 mg kg<sup>-1</sup>) had blood collected from the left heart ventricle and livers dissected for subsequent analyses after 14 days. Serum was subjected to biochemical analysis to determine the levels of AST and ALT. The liver was subjected to biochemical analysis to quantify the levels of malondialdehyde (MDA) and glutathione (GSH) and to assess the activities of catalase (CAT), superoxide dismutase (SOD) and glutathione peroxidase (GPx; El Rabey *et al.* 2021; Abdollahi *et al.* 2024).

## Measurements of biochemical parameters

## Determination of liver enzyme activity

Serum ALT and AST levels were quantified using biochemical enzyme kits (Sigma-Aldrich Company, USA) according to the manufacturer's instructions (Babaeenezhad *et al.* 2021).

## Homogenization of liver samples

To obtain a liver homogenate, liver samples were minced and homogenized in Tris-HCl buffer (25 mM, pH 7.5) using a homogenizer. Centrifugation at 12,000 rpm for 15 minutes at 4 °C was performed on the resulting 10% (w/v) liver homogenate (Abdollahi *et al.* 2024).

# Measurement of oxidative stress markers

## Malondialdehyde (MDA)

Malondialdehyde (MDA) levels were quantified using a modified protocol based on Esterbauer and Zollern (Esterbauer & Zollern 1989). The thiobarbituric acid (TBA) test was used to assess hepatic MDA levels. Each sample was assayed in triplicate. Absorbance was measured by spectrophotometry at 532 nm. The absorbance was quantified and reported in units of  $\mu$ mol mg<sup>-1</sup> of protein malondialdehyde (MDA; Nasiry *et al.* 2017).

## **Glutathione (GSH)**

We implemented a slight modification of the protocol described by Rahman *et al.* (Rahman *et al.* 2007) for the quantification of glutathione (GSH). Specifically, 25 mL of liver homogenate was combined with 140 mL of 0.2 M Tris-EDTA buffer (pH 8), followed by adding 30 mL of 0.1 M DTNB. Absorbance was read at 412 nm and checked against blank. Each sample was assayed in triplicate. GSH levels were finally expressed as  $\mu$ mol L<sup>-1</sup> mg<sup>-1</sup> protein GSH (Ahmadvand *et al.* 2017).

# Catalase (CAT)

Catalase (CAT) activity was determined using method modified by Aebi (Aebi 1984).First, 1 mL of 50 mM potassium phosphate buffer (pH 8) was added to 50-mL liver homogenate, followed by the addition of 50-mL  $H_2O_2$ . A spectrophotometer was used to measure absorbance at 240 nm in triplicate, with measurements taken at 0, 30, and 60 seconds against a blank. Results were expressed in units per milligram (U mg<sup>-1</sup>; Ahmadvand *et al.* 2017).

## Glutathione peroxidase (GPx)

Hepatic glutathione peroxidase (GPX) activities were determined by the method of Rotruck and colleagues (Rotruck *et al.* 1973). An ELISA reader was used to read the absorbance at 420 nm. GPX activity was obtained in units of protein (U mg<sup>-1</sup>; Babaeenezhad *et al.* 2021).

## Superoxide dismutase (SOD)

Superoxide dismutase (SOD) activity was determined by the Lund method, based on the enzyme's ability to inhibit pyrogallol oxidation. The absorbance of the samples was measured immediately and again after 60 seconds. The control was the difference in absorbance between the initial and the 60 s interval. The initial and 60-second absorbance readings were taken. The percentage inhibition of pyrogallol autoxidation was then calculated as follows (Sun *et al.* 1988):

Absorbance of the sample at zero time – absorbance of the sample after 60 sAsorbance of the control at zero time – absorbance of the control after 60 s The activity of SOD was quantified and expressed as units per milligram (U mg<sup>-1</sup>) using the following formula:

SOD enzyme activity (U mg<sup>-1</sup>) =  $\frac{Percentage of inhibition of pyrogalol autoxidation}{50\%}$ 

### Statistical analysis

One-way analysis of variance (ANOVA) was used to analyze the experimental groups after verifying the normality of the data using the Shapiro-Wilk test. Tukey's post hoc test was used to compare means between experimental groups. A 5% significant level was maintained throughout all stages of analysis ( $\alpha = 0.05$ ). SPSS version 24 software was used for statistical analysis, and GraphPad Prism version 9 for graphing.

# RESULTS

#### Effect of eucalyptol on the serum level of AST and ALT

As shown in Table 1 and Fig. 1A, the AST serum levels were significantly elevated in CCl<sub>4</sub>-treated rats versus control animals (p < 0.0001). Conversely, the serum AST levels were significantly decreased in the eucalyptol-treated group in comparison with the CCl<sub>4</sub>-group (p < 0.0001). The data presented in Table 1 and Fig. 1B clearly indicate that the serum ALT levels increased significantly in the CCl<sub>4</sub>-group in comparison with the control rats (p = 0.0479). However, no significant changes were observed in the serum ALT levels in the eucalyptol-treated group compared to the CCl<sub>4</sub> group (p = 0.8946).

 Table 1. Effects of eucalyptol on the serum levels of liver enzymes (AST and ALT) and liver levels of oxidative stress markers (MDA, GSH, GPX, CAT and SOD) in CCl4 hepatotoxicity in different groups.

Parameter	Group			<i>p</i> -value
	Healthy control	CCl <sub>4</sub> -group	Eucalyptol-treated group	
AST	$28.50\pm2.8$	$159.2 \pm 14.82$ ****	99.60 ± 5.1 ####****	< 0.0001
ALT	$33.50\pm5.13$	44.80 ± 9.53 **	41.50 ± 5.29 *	0.0036
MDA	$95.12\pm23.98$	$149.1 \pm 58.83$ *	$120.4 \pm 50.79$	0.0480
GSH	$17.28 \pm 1.69$	8.83 ± 1.12 ****	$12.29 \pm 1.35^{\texttt{#}\texttt{#}\texttt{#}\texttt{#}}$	< 0.0001
GPX	$76.91 \pm 8.21$	$46.26 \pm 7.32$ ****	$61.17 \pm 10.12^{\#}$ **	< 0.0001
CAT	$12.89 \pm 1.57$	$7.88 \pm 2.64$ ***	$9.48 \pm 2.69$ **	0.0002
SOD	$9.95 \pm 1.77$	$3.40 \pm 0.97$ ****	$6.85 \pm 1.29^{\texttt{#}\texttt{#}\texttt{#}} ~\texttt{****}$	< 0.0001

\* Significant change in comparison with the control group at p < 0.05; #Significant change in comparison with the CCl<sub>4</sub>-group at p < 0.05.

#### Effect of Eucalyptol on the level of MDA

The data indicated a statistically significant elevation in hepatic MDA levels in the CCl<sub>4</sub> group compared to the control rats (p = 0.0480). Conversely, there was no statistically significant difference in hepatic MDA levels between the eucalyptol-treated and the CCl<sub>4</sub> groups (p = 0.5482; Table 1 and Fig. 2A).

## Effect of eucalyptol on the level of GSH

Table 1 and Fig. 2B illustrates a statistically significant reduction in hepatic GSH levels in the CCl<sub>4</sub>-treated group relative to the control ones (p < 0.0001). Conversely, the eucalyptol treated-group exhibited a significant elevation in hepatic GSH levels than in the CCl<sub>4</sub>-treated group (p < 0.0001).

#### Effect of eucalyptol on the activity of GPX

Exposure to CCl<sub>4</sub> was shown to significantly decrease GPX activity in the CCl<sub>4</sub>-treated group than in the control rats (p < 0.0001). In addition, eucalyptol treatment resulted in a significant elevation in GPX activity in the eucalyptol-treated group than in the CCl<sub>4</sub>-treated one (p = 0.0021; Table 1 and Fig. 2C).

#### Effect of eucalyptol on the activity of CAT

The findings demonstrated a significant reduction in hepatic CAT activity in the CCl<sub>4</sub>-treated rats than in the control animals (p = 0.0002). Additionally, no significant change in hepatic CAT activity was found between the eucalyptol- treated and CCl<sub>4</sub>-treated groups (p = 0.4281; Table 1 and Fig. 2D).

#### Effect of eucalyptol on the activity of SOD

As shown in Table 1 and Fig. 2E, there was a meaningful decrease in hepatic SOD activity in the CCl<sub>4</sub>-treated animals than in the control rats (p < 0.0001). Furthermore, SOD activity was significantly enhanced in the eucalyptol-treated group than in the CCl<sub>4</sub>-treated rats (p < 0.0001).



Fig. 1. Effect of eucalyptol on the serum levels of liver enzymes including AST (A) and ALT (B) in CCl<sub>4</sub>-induced hepatotoxicity. Data are represented as Mean  $\pm$  SD. Comparisons between groups were made using One-Way ANOVA followed by post hoc test. \*Significant change at p < 0.05.

#### DISCUSSION

The present study was designed to evaluate the ameliorative properties of eucalyptol against liver oxidative damage caused by CCl<sub>4</sub> in rats. The present study is believed to be the first to demonstrate the ameliorative properties of eucalyptol against liver oxidative damage caused by CCl<sub>4</sub> in rats. The current study concluded that eucalyptol treatment can mitigate the harmful effects of CCl<sub>4</sub> on the rat liver. Most of the various parameters assessed in our experiment were rapidly restored in CCl4-intoxicated animals after eucalyptol treatment. Our results showed that exposure to CCl<sub>4</sub>-induced liver damage as evidenced by the elevated serum AST and ALT levels. Furthermore, exposure to CCl<sub>4</sub> was associated with an elevation in MDA, which serves as a marker of oxidative stress, and a concomitant decrease in the antioxidant parameters such as GSH, GPX, CAT, and SOD. Indeed, CCl4-induced liver damage has been similarly documented in numerous previous studies (Wahid et al. 2016; El Rabey et al. 2021). ALT and AST serve as biomarkers to assess liver damage. When the cell membrane integrity is compromised, blood levels of these enzymes increase (Abdollahi et al. 2024). The enhanced production of free radicals facilitated by CCl<sub>4</sub> may contribute to the degradation of cell membranes and subsequent liver damage (El Rabey et al. 2021). In this study, administration of eucalyptol was shown to reduce serum levels of liver enzymes, particularly AST, confirming its ameliorative effect against liver damage through its antioxidant properties. There are several pieces of evidence that the effectiveness of euclyptol in preventing the increased ALT and AST levels is dependent on dose (Abdollahi et al. 2024). It was also found that the hepatic MDA levels, an important marker of oxidative stress, decreased in treated rats after administration of eucalyptol. Several lines of evidence have shown that eucalyptol can mitigate ROS generation and improve endogenous antioxidant status. Environmental contaminants, such as CCl<sub>4</sub>, induce oxidative stress and damage biomolecules in liver cells, resulting in structural and functional changes in the cells (Abdollahi et al. 2024). Our results showed an elevation in the antioxidant parameter (GSH) after eucalyptol treatment. In addition, a meaningful recovery in the hepatic activities of antioxidant enzymes, including GPX, CAT, and SOD, was observed after eucalyptol administration. Our research aligns with prior studies, demonstrating that eucalyptol has the potential to mitigate oxidative stress

across various pathological conditions (Ciftci *et al.* 2011; Abdollahi *et al.* 2024). It can be concluded that amelioration of oxidative stress by eucalyptol can be associated with its ROS scavenging activity (Abdollahi *et al.* 2024). Our findings align with those of Ciftci *et al.*, demonstrating that the administration of eucalyptol to rats exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin enhances the activities of catalase, superoxide dismutase (SOD), and glutathione peroxidase (GPx), as well as the glutathione (GSH) levels, thereby mitigating oxidative stress in the liver (Ciftci *et al.* 2011). Through our biochemical studies, we substantiated the hypothesis regarding the hepatoprotective properties of eucalyptol.



Fig. 2. Effects of eucalyptol on the level of oxidative stress parameters including MDA (A), GSH (B), GPX (C), CAT (D) and SOD (E) in CCl<sub>4</sub>-induced hepatotoxicity. Results are represented as Mean ± SD. Comparisons between groups were made using One-Way ANOVA followed by post hoc test. \*Significant change at p < 0.05.</p>

#### CONCLUSION

Our study is the first to demonstrate the efficacy of eucalyptol in mitigating liver damage induced by carbon tetrachloride exposure. The antioxidant properties of eucalyptol significantly contribute to this hepatoprotection by ameliorating liver enzyme dysfunction, reducing oxidative stress and free radical production, as well as enhancing the endogenous antioxidant system.

#### REFERENCES

- Abdollahi, M, Asle-Rousta, M & Mahmazi, S 2024, Protective effect of 1, 8-cineole (eucalyptol) against leadinduced liver injury by ameliorating oxidative stress and inflammation and modulating TLR4/MyD88/NFκB signaling. *Iranian Journal of Basic Medical Sciences*, 27: 1293-1299.
- Aebi, H 1984, Catalase in vitro. In Methods in enzymology. Academic press. 105: 121-126.
- Ahmadvand, H, Babaeenezhad, E, Hadipour Moradi, F & Cheraghi Venool, A 2017, Effect of gallic acid on liver oxidative stress markers in renal ischemia-reperfusion injury in rats. *Annals of Research in Antioxidants*, 2: e03.
- Asadi, N, Shahriary, A, Jafarzadeh, M, Ghorbani, M & Yavari, E 2024, Effects of coenzyme Q10 and Nacetylcysteine on the expression of apoptotic biomarkers and histopathological findings in the permethrininduced hepatotoxicity in rats. *Caspian Journal of Environmental Sciences*, 22: 841-847.
- Babaeenezhad, E, Nouryazdan, N, Nasri, M, Ahmadvand, H & Moradi Sarabi, M 2021, Cinnamic acid ameliorate gentamicin-induced liver dysfunctions and nephrotoxicity in rats through induction of antioxidant activities. *Heliyon*, 7: e07465.
- Cai, ZM, Peng, JQ, Chen, Y, Tao, L, Zhang, YY, Fu, LY, Long, QD & Shen, XC 2021, 1, 8-Cineole: A review of source, biological activities, and application. *Journal of Asian Natural Products Research*, 23: 938-954.
- Ciftci, O, Ozdemir, I, Tanyildizi, S, Yildiz, S & Oguzturk, H 2011, Antioxidative effects of curcumin, β-myrcene and 1, 8-cineole against 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin-induced oxidative stress in rats liver. *Toxicology and Industrial Health*, 27: 447-453.
- Devi, AB, Rahigude, AB, Parab, PB, Engineer, AS, Dhakephalkar, PK & Apte, KG 2021. A study to evaluate the hepatoprotective activity of prebiotics, probiotics, and synbiotic in CCl<sub>4</sub> induced hepatotoxicity in rats. *Journal of Applied Pharmaceutical Science*, 11: 141-153.
- Di Paola, R, Modafferi, S, Siracusa, R, Cordaro, M, D'amico, R, Ontario, ML, Interdonato, L, Salinaro, AT, Fusco, R & Impellizzeri, D 2022, S-acetyl-glutathione attenuates carbon tetrachloride-induced liver injury by modulating oxidative imbalance and inflammation. *International Journal of Molecular Sciences*, 23: 4429.
- El Rabey, HA, Rezk, SM, Sakran, MI, Mohammed, GM, Bahattab, O, Balgoon, MJ, Elbakry, MA & Bakry, N 2021, Green coffee methanolic extract and silymarin protect against CCl<sub>4</sub>-induced hepatotoxicity in albino male rats. *BMC Complementary Medicine and Therapies*, 21: 1-11.
- Esterbauer, H & Zollern, H 1989, Methods for determination of aldehydic lipid peroxidation products. *Free Radical Biology and Medicine*, 7: 197-203.
- Hashemzadeh, M, Moghaddam, A, Karimi, I, Siavosh Haghighi, ZM & Hassanpour, S 2018, Semiprotective effects of hempseed oil on carbon tetrachloride-induced hepatotoxicity in male rats: an ultra-short toxicological intervention. *Journal of Chemical Health Risks*, 7: 319-330.
- Hoch, CC, Petry, J, Griesbaum, L, Weiser, T, Werner, K, Ploch, M, Verschoor, A, Multhoff, G, Dezfouli, AB & Wollenberg, B 2023, 1, 8-cineole (eucalyptol): A versatile phytochemical with therapeutic applications across multiple diseases. *Biomedicine & Pharmacotherapy*, 167: 115467.
- Kabbashi, AS, Eltawaty, SA, Ismail, AM, Elshikh, AA, Alrasheid, AA, Elmahi, RA, Koko, WS & Osman, EE 2024, Ethanolic extract of *Mangifera indica* protects against CCl<sub>4</sub>-induced hepatotoxicity via antioxidant capabilities in albino rats. *Journal of Toxicology*, 2024: 5539386.
- Kennedy-Feitosa, E, Cattani-Cavalieri, I, Barroso, MV, Romana-Souza, B, Brito-Gitirana, L & Valenca, SS 2019, Eucalyptol promotes lung repair in mice following cigarette smoke-induced emphysema. *Phytomedicine*, 55: 70-79.
- Kostic, T, Popović, D, Perisic, Z, Stanojevic, D, Dakic, S, Saric, S, Radojkovic, DD, Apostolovic, S, Bozinovic, N & Zdravkovic, SC 2022, The hepatoprotective effect of aminoguanidine in acute liver injury caused by CCl<sub>4</sub> in rats. *Biomedicine & Pharmacotherapy*, 156: 113918.

- Moghadam, M, Ansari, I, Roghani, M, Ghanem, A & Mehdizade, N 2017, The effect of oral administration of Hypericum perforatum on serum glucose and lipids, hepatic enzymes and lipid peroxidation in streptozotocin-induced diabetic rats. *Galen*, 6: 319-29.
- Nasiry, D, Khalatbary, AR & Ahmadvand, H 2017, Therapeutic potential of Juglans regia L. leaf extract against diabetic retinopathy in rat. *Iranian Journal of Basic Medical Sciences*, 20: 1275.
- Parpieva, M, Mirkhamidova, P, Pozilov, M & Alimova, R 2023, Effects of sophoroflavonoside and narcissin flavonoids on the amount of malondialdehyde, a product of lipid peroxidation, and the activity of the enzyme cytochrome c oxidase in rat liver mitochondria poisoned with indoxacarb pesticide. *Caspian Journal of Environmental Sciences*, 21: 997-1000.
- Que, R, Cao, M, Dai, Y, Zhou, Y, Chen, Y & Lin, L 2022, Decursin ameliorates carbon-tetrachloride-induced liver fibrosis by facilitating ferroptosis of hepatic stellate cells. *Biochemistry and Cell Biology*, 100: 378-386.
- Rahman, I, Kode, A & Biswas, SK 2007, Assay for quantitative determination of glutathione and glutathione disulfide levels using enzymatic recycling method. *Nature Protocols*, 1: 3159-3165.
- Rotruck, JT, Pope, AL, Ganther, HE Swanson, AB, Hafeman, DG & Hoekstra, WG 1973, Selenium: biochemical role as a component of glutathione peroxidase. *Science*, 179: 588-590.
- Sun, Y, Oberley, LW & Li, Y 1988, A simple method for clinical assay of superoxide dismutase. *Clinical Chemistry*, 34: 497-500.
- Teschke, R 2018, Liver injury by carbon tetrachloride intoxication in 16 patients treated with forced ventilation to accelerate toxin removal via the lungs: A clinical report. *Toxics*, 6: 25.
- Ugochukwu Madu, F, Chinwenmeri Madu, M & Aderinola Adejumo, A 2024, Hepatorenal-toxicities Accompanying Exposure to Indoor Air Pollutants in Chemical Storekeepers at an International Market. *Journal of Chemical Health Risks*, 14: 0-0.
- Vahidi-Eyrisofla, N, Hojati, V, Yazdian, MR, Zendehdel, M & Shajiee, H 2019, Effects of olive leaf extract on prevention of molecular, histopathological, and enzymatic changes in chicken carbon tetrachloride-induced liver damage. *Galen Medical Journal*, 8: e1204.
- Wahid, A, Hamed, AN, Eltahir, HM & Abouzied, MM 2016, Hepatoprotective activity of ethanolic extract of *Salix subserrata* against CCl<sub>4</sub>-induced chronic hepatotoxicity in rats. *BMC Complementary and Alternative Medicine*, 16: 1-10.
- Xiong, R, Shan, S, Wang, X, Zhang, X, Yu, H, Shi, H & Wang, X 2020, Aloperine attenuates carbon tetrachlorideinduced mouse hepatic injury via Nrf2/HO-1 pathway. *Tropical Journal of Pharmaceutical Research*, 19: 983-988.
- Yeh, YH, Hsieh, YL & Lee, YT 2013, Effects of yam peel extract against carbon tetrachloride-induced hepatotoxicity in rats. *Journal of Agricultural and Food Chemistry*, 61: 7387-7396.