



First evidence of intersex condition in Caspian kutum, *Rutilus kutum* (Kamenski, 1901) in the Southwest Caspian Sea

Babak Tizkar^{1*}, Houshang Dehghanzadeh², Ali Hallajian³, Afshar Zoughi Shalmani¹

1. Aquaculture Department, Guilan Agricultural and Natural Resources Research and Education Center Agricultural Research, Education and Extension Organization (AREEO), Rasht, Iran

2. Scientific Board Member of Animal Science Research Department, Guilan Agricultural and Natural Resources Research and Education Center, Agricultural Research, Education and Extension Organization (AREEO), Rasht, Iran

3. International Sturgeon Research Institute, Agricultural Research Education and Extension Organization (AREEO), Rasht, Iran

* Corresponding author's E-mail: btizkar@gmail.com

ABSTRACT

The present study showed intersex conditions in the wild population of Caspian kutum in the Southwest Caspian Sea. Evidence and examples of morphological and structural changes have been found in recent years among Caspian kutum, *Rutilus kutum*. The present study included sampling and examining morphological and anatomical characteristics of 350 pieces of kutum specimens (38.5 ± 1.87 cm, 755 ± 6.52 g, and +3yr) as per their gender. Out of these numbers, a single male (+3 years old and a length/weight of 32.25 cm/628.5 g respectively) showed anomalies comprising seminal and gonadal sacs with simultaneous presence of male and female gonads. Histological observations of the male gonad revealed the presence of ovotestis tissue containing oocytes and spermatozoa throughout the entire testis. Male gonad tissue exhibited somniferous tubular structures with spermatogonia, spermatocytes, and spermatids. The histological examination of the gonads also showed the predominance of oocytes at different developmental stages. The intersexuality might have been due to determining processes or a combination of natural malformation and chemical contamination.

Keywords: Kutum, Histological analysis, Intersex gonads, Oocytes.

Article type: Report.

INTRODUCTION

By a surface area of 373,300 km², the Caspian Sea is the largest landlocked water body in the world, wherein 120 species and subspecies of fish live (Pourkazemi & Razikazemi 2011). *Rutilus kutum*, known as Caspian kutum, is one of the most commercially important fish species in the Caspian Sea. The fish is found in the Azerbaijan Republic, Iranian coasts, and the northern shores of the Black Sea. (Berg 1964; Razavi-Sayyad 1990). Most of the kutum populations are distributed in the Southwestern Caspian Sea (Heidari *et al.* 2009; Pourkazemi & Razikazemi 2011). Kutum is a migratory anadromous teleost with a group-synchronous ovary, and the eggs are released once after ovulation (Sharyati 1993; Saeed 2010), on gravel and sandy substrates and aquatic weeds of rivers or lagoons (Abdoli 1999; Koochilal *et al.* 2016). The females and males reach sexual maturity between 2-3 and 3-4 years respectively (Afraei-Bandpeid *et al.* 2012). Kutum fish migrate to estuaries and coastal waters in mid-winter (Abdoli 1999). The fish migrate to the Iranian side of the Caspian Sea to spawn in the rivers that flow into the sea and in Anzali wetland from March to May (Fazli *et al.* 2013). However, if the kutum is caught in the delta, cannot spawn (Yousefian & Mosavi 2008). The biology of kutum reproduction in the Iranian side of the Caspian Sea has been studied mostly on subjects such as sex differentiation (Pourkazemi & Razikazemi 2011), oocyte development (Heidari *et al.* 2010), spawning (Dorafshan & Heyrati 2006; Heyrati *et al.* 2007; Yousefian



& Mosavi 2008; Falahatkar *et al.* 2013; Koohilaei *et al.* 2016), fertility (Farid-Pak 1968); Heidari *et al.* 2009; Masouleh *et al.* 2011; Samarin *et al.* 2011; Bandpei *et al.* 2011; Fazli *et al.* 2013), states of maturity (Sabet *et al.* 2009; Saeed 2010) and reproductive performance (Savadkouhi & Khara 2017). In addition, studies on reproductive impairments have been carried out on wild fish species in various parts of the world, such as male masculinization (Howell *et al.* 1980), interrupted or delayed testicular development (Lye *et al.* 1998), and abnormal reproductive behavior (Jones & Reynolds 1997). In wild populations of gonochoric fish, alterations may occur with the simultaneous presence of male and female reproductive stages in the same gonad at the same time which is not normal and is known as intersex (Nolan *et al.* 2001). In contrast, in protandry or protogynous fish species, the presence of the two types of gonads in the same fish is called hermaphroditism (Bahamonde *et al.* 2013) but in gonochoristic (fixed-sex) species intersex is not a normal reproduction mode or life cycle (Nolan *et al.* 2001; Tyler & Jobling 2008; Bahamonde *et al.* 2013). This means both testicular and ovarian cells appear in gonochoric fish gonads (Abdel Moneim *et al.* 2015). Widespread occurrences of intersex have been observed in freshwater and estuarine fish in various parts of the world (Jobling *et al.* 2002). There are several causes for intersex occurrences, among which temperature, pH, exogenous steroids, behavioral cues, parasites, and pollutants are considered to be involved in sex differentiation in fish (Minier *et al.* 2000; Devlin & Nagahama 2002; Hinck *et al.* 2009; Ichalal *et al.* 2016). The presence of intersex has been reported in the Cyprinidae, to which kutum belongs. The first findings reported were related to bream, *Abramis brama* in Netherlands (Sloof & Klootwijk-Vandijk 1982), Roach, *Rutilus rutilus* in British Isles (Jobling *et al.* 1998), chub, *Squalius cephalus* in England and France (Minier *et al.* 2000), barbel, *Barbus plebejus* in Italy (Viganò *et al.* 2001), common carp, *Cyprinus carpio* in Spain (Solé *et al.* 2002), spottail shiner, *Notropis hudsonius* in Canada (Aravindakshan *et al.* 2004), bighead carp, *Hypophthalmichthys nobilis* and silver carp, *H. molitrix* in USA (Papoulias *et al.* 2006). In general, intersexuality among wild fish is attributed to endocrine-disrupting substances (Nolan *et al.* 2001). However, there has been no earlier report on any cases of intersex gonads in Caspian kutum. This study aims to introduce the first gonadal intersex anomaly among Caspian kutum, *R. kutum* from the Southwest Caspian Sea.

MATERIALS AND METHODS

Sampling Area and Fish Collection

The fish specimens were captured using a beach seine net by local fishermen from the Sefidrud River estuary (37° 27'31.32"N; 49° 56'05.78" E) in April 2016. The river is approximately 670 km long and flows into the Caspian Sea (Asadollahfardi *et al.* 2018). Out of 350 fish specimens, only one intersex case was detected. Standard length (nearest 0.1 cm), body weight, and gonad weight (nearest 0.1 g) were measured. The entire gonad of intersex fish was removed, photographed, and fixed in 10% neutral buffered formalin for histological analyses.

Histological Analyses

Proximal, medium, and distal portions of the right gonads were fixed in Bouin's fixative solution, followed by tissue processing (consisting of dehydration, clearing, and paraffin embedding stages). Dehydration was conducted in ethanol 50, 70, 80, and 96% and 1-butanol, followed by clearing chloroform and paraffin embedding in a mixture of chloroform with purified paraffin at 56 °C. Then, the sample template was molded and assembled on a wooden base, sectioned longitudinally at 7 µm, mounted on slides coated with gelatin tissue (tissue to bind on the slide), and stained with hematoxylin and eosin (H&E). Next, the slides were studied through the light microscope (model Nikon E600, Japan) connected to a computer via Biocom program to study the tissues (Akhundov & Federov 1995; Hallajian 2009; Sharifpour *et al.* 2013), followed by photographing at 40X, 100X, and 200X magnifications.

RESULTS AND DISCUSSION

The mean standard length, body weight, and age of 350 pieces of Caspian kutum were 38.5 ± 1.87 cm, 755 ± 6.52 , and +3 years, respectively). Macroscopic examination of the collected fish showed characteristics of either male or female. However, in the macroscopic and histological analyses of the gonads, a single male fish (length, weight, and age 32.25 cm, 628.5 g, 3+, respectively) presented anomalies showing seminal or gonadal sac with simultaneous presence of male and female gonads. Thus, it was identified as an instance of intersex organism (length, weight and age 32.25 cm, 628.5 g, 3+ respectively), confirming the relatively low intersexuality occurrence rate (0.3%) in the Caspian kutum population, compared to many other fish species: 0.75% to 2.28% (Djoudad-Kadji *et al.* 2012; Ichalal *et al.* 2016; Hassel *et al.* 2018). The examined gonad comprised a testis

developed in the proximal region, which occupied a small portion. However, the ovarian tissue occupied the most significant portion of the gonad (Fig. 1), similar to the finding in *Trachurus trachurus* (Ichalal et al. 2016), however contrary to that found in *Coryphaena hippurus* (Retheesh et al. 2017). The testis was seen to be whitish, while the ovary was pinkish-brown in normal gonads. The presence of oocytes in the female lobe was detected in the distal portion (Fig. 1). The macroscopic and histological analyses of gonads revealed the presence of both ovarian and testicular tissues. Since the intersex specimen was obtained from the wild catch in the Sefid-Rud estuary, it was impossible to determine whether the fish was reproductively functional. Therefore, it was impossible to determine if it was an undifferentiated or differentiated gonochoristic species (Yamamoto 1969; Jafri & Ensor 1979).

The mature ovotestis was subjected to histological analysis by microscope using proximal (Fig. 2), middle (Fig. 3), and distal (Fig. 4) regions of gonads. This showed variation in the composition of male and female cells and their arrangement within the gonads. The gonads presented oocytes at various stages of development. Intersex cells in this particular kutum (testis and ovary) were identified based on shape, size, cytology criteria, and characteristics of the nucleus (Tyler & Sumpter 1996; Nolan et al. 2001). The tissues were distinct from each other since the gonad was predominantly ovarian, and the testicular tissue was always white. The male tissue was restricted to the proximal region and the gonad surrounding the ovarian sinus, called the accessory reproductive structure. In most of the histological sections, oocytes and sperm were found to exhibit normal development. Macroscopic and microscopic examinations of the ovarian portion showed the presence of oocytes at different developmental stages and the testicular portion with spermatozoa mass (Fig. 2), confirming the rare occurrence of intersex in Caspian kutum. In the testis, seminiferous tubules were evident, and cells showed in all stages of spermatogenesis (spermatozoa, spermatid, spermatocytes, and spermatogonia).

Samples with hydrated oocytes and testicular tissue containing sperm were recorded simultaneously in both regions of the gonad. A predominance of spermatogonia was found compared to spermatids and spermatocytes, with several primary and secondary stages of oocytes in testicular tissue. Seminiferous tubules containing cells in different stages of spermatogenesis were observed in the testicular portion. Oocytes at different developmental stages (1-4) were detected in the middle gonad region of intersex male. In the present study, we detected only one specimen with characteristics of intersex after examining 350 specimens, suggesting a low occurrence rate of such abnormality. Such a low occurrence of intersex in Caspian kutum may be due to the current climatic condition and/or natural anthropogenic pollution as has been described for other species (Retheesh et al. 2017). It has been found that various natural and synthetic pollutants can alter sexual differentiation in wild populations of gonochoric species by inducing sexual inversion and/or intersexuality (Colborn and Clement 1992) and also by endocrine disruption caused by water contamination (Sadovy de Mitcheson & Liu 2008). Kutum fish are known to spend part of their life cycle in some regions of the South Caspian Sea, wherein water and sediment are affected by contaminants (De Mora et al. 2004; Saeidi & Jamshidi 2010; Gharedaashi et al. 2013; Zahedi et al. 2014; Hoseini et al. 2015). The combination of heavy metals with detergents has a considerable effect on the mortality of Caspian kutum (Gholami et al. 2010). Insecticides such as fipronil applied in rice cultivation have also been found to affect alterations in the organ-somatic index and DNA structure in Caspian kutum (Ardeshir et al. 2018). It has been confirmed that kutum exposure to the butachlor (a herbicide used as weed control in rice fields in the study area) resulted in a decreased number and volume of sperm as well as an increased rate (28.6%) of abnormal spermatozoa (Lasheidani et al. 2008).

Furthermore, very low levels of Diazinon (an organophosphate pesticide) depressed spermatogenesis in testes and sperm quality of kutum (Masouleh et al. 2011). In line with the earlier findings, the impact of water pollution on the reproductive biology of kutum and the emergence of intersex cases among such species might be related to the high concentration of trace elements detected in the kutum gonads caught in the study area as compared to kutum caught from the eastern part of the South Caspian Sea (Sattari et al. 2019). In conclusion, the study showed the rare occurrence of intersex in Caspian kutum, indicating the simultaneous presence of the gonad and ovary tissue system in male kutum, as evidenced by macroscopic and histological observations. However, further research is necessary to shed more light on this subject since our study relies on the record of a single specimen.

Suggestion for further research

The causes of intersexuality might be due to determining processes or a combination of natural malformation and chemical contamination. Therefore, continuous monitoring of kutum populations in the Southwest Caspian Sea is required.




Fig. 1. Macroscopic appearance of intersex specimen. PS: proximal; MS: middle; DS: distal section; RT: right testis; LT: left testis.




Fig. 2. Histology of right side gonad from intersex Caspian kutum, *Rutilus kutum* captured in the Southwest Caspian Sea: proximal section; a) H & E 40X; b) HE 100X; c) HE 200X. Basal lamina (Bl); structures showing the lobules (Lo); lumen (Lu); Secondary oocyte (OC2); Ovarian lamellae (Ol); ovarian lumen (OLu); ovary tissue (Ot); early primary growth oocyte (Pg1); spermatocyte (SC); sperm duct (SD); spermatozoa (SG); spermatogonia (SG); spermatozoa (ST); spermatid (St); sperm sinus and sperm duct filled with spermatozoa (SZ); testicular tissue (Tt).




Fig. 3. Histology of right side gonad from intersex Caspian kutum, *Rutilus kutum* captured in the Southwest Caspian Sea medium section. a) H & E, 40X; b) H&E 100X; secondary oocyte (OC2); ovarian lumen (OLu); ovary sinus (Os); ovary tissue (Ot); early primary growth oocyte (Pg1); spermatocyte (SC); spermatogonia (SG); spermatozoa (ST); spermatid (St); sinus wall (Sw); sperm sinus and sperm duct filled with spermatozoa (SZ); testicular tissue (Tt); primary and secondary vitellogenesis stages (VtgII); quaternary vitellogenetic stage (VtgIV); yolk granules (YG).




Fig. 4. Histology of right side gonad from intersex Caspian kutum, *Rutilus kutum* captured in the Southwest Caspian Sea: distal section. a) H & E, 40X; b) H&E 100X; c) H & E 200X. Disintegration of the nucleus (N); Primary Oocyte (OC1); Secondary Oocyte (OC2); Oil droplets (OD); Oil globules (OG); Oocyte or Ovum (OO); Ovary tissue (Ot); Early primary growth oocyte (Pg1); later primary growth stage (Pg2); Spermatocyte (SC); Testicular tissue (Tt); Primary and secondary vitellogenesis stages (VtgI); Yolk granules (YG); Zona radiata (ZR).

ACKNOWLEDGEMENTS

This study was supported by the Agricultural Research, Education and Extension Organization (AREEO), Rasht, Iran.

REFERENCES

- Abdel-Moneim, A, Coulter, DP, Mahapatra, CT & Sepúlveda, MS 2015, Intersex in fishes and amphibians: population implications, prevalence, mechanisms and molecular biomarkers. *Journal of Applied Toxicology*, 35:1228-1240, <https://doi.org/10.1002/jat.3204>.
- Abdoli, A 1999, The Inland Water Fishes of Iran. *Natural and Wild Life Museum of Iran*, Tehran, Iran, 378 p. [In Persian].
- Afraei Bandpei MA, Mansor, M, Abdolmalaki, SH, Keymars, F, Mohamad-Isa, M & Janbaz, AA 2010, Age and growth of kutum (*Rutilus frisii kutum*, Kamensky, 1901) in southern Caspian Sea. *International Aquatic Research*, 2: 25- 33.
- Akhundov, MM & Fedorov, KY 1995, Effect of exogenous estradiol on ovarian development in juvenile sterlet (*Acipenser ruthenus*). *Journal of Ichthyology*, 35: 109-120.
- Aravindakshan, J, Paquet, V, Gregory, M, Dufresne, J, Fournier, M, Marcogliese, DJ & Cyr, DG 2004, Consequences of xenoestrogen exposure on male reproductive function in spot tail shiners (*Notropis hudsonius*). *Toxicological Science*, 78: 156-165, <https://doi.org/10.1093/toxsci/kfh042>.
- Ardehshir, RA, Zolgharneina, H, Movahedinia, A, Salamata, N & Zabihic, E 2018, CYP1A gene expression as a basic factor for fipronil toxicity in Caspian kutum fish. *Toxicology Reports*, 5: 113-124, <https://doi.org/10.1016/j.toxrep.2017.12.014>.
- Asadollahfardi, G, Heidarzadeh, N, Mosalli, A & Sekhavati, A 2018, Optimization of water quality monitoring stations using genetic algorithm, a case study, Sefid-Rud River, Iran. *Advances in Environment Research*, 7(2): 87-107, <https://doi.org/10.12989/aer.2018.7.2.087>.
- Bahamonde, PA, Munkittrick, KR & Martyniuk, CJ 2013, Intersex in teleost fish: are we distinguishing endocrine disruption from natural phenomena? *General and Comparative Endocrinology*, 192: 25-35, <https://doi.org/10.1016/j.ygcen.2013.04.005>.
- Bandpei, A, Mashhor, MAM, Abdolmaleki, SH, Najafpour, SH, Bani, A, Pourgholam, R, Fazli, H, Nasrolahzadeh, H & Janbaz, AA 2011, The environmental effect on spawning time, length at maturity and fecundity of Kutum (*Rutilus frisii kutum*, Kamensky, 1901) in southern part of the Caspian Sea, Iran. *Iranian Journal of Energy and Environment*, 2: 374-381, <https://doi.org/10.5829/idosi.ijee.2011.02.04.3097>.
- Berg, LS 1964, Freshwater fishes of USSR and adjacent countries. Vol. II, 4th edition. Jerusalem: Israeli Program for Scientific Translation.
- Colborn, T & Clement, C 1992, Chemically Induced Alterations in Sexual and Functional Development the Wildlife/Human Connection. vol. XXI. Princeton Scientific, Princeton, NJ, USA, 403 p.

- De Mora, S, Sheikholeslami, MR, Wyse, E, Azemard, S & Cassi, R 2004, An assessment of metal contamination in coastal sediments of the Caspian Sea. *Marine Pollution Bulletin*, 48: 61-77, [https://doi.org/10.1016/S0025-326X\(03\)00285-6](https://doi.org/10.1016/S0025-326X(03)00285-6).
- Devlin, RH & Nagahama, Y 2002, Sex determination and sex differentiation in fish: an overview of genetic, physiological, and environmental influences. *Aquaculture*, 208: 191-364.
- Djoudad-Kadji, H, Benslimane, S, Chevalier, C, Kadji, B, Exbrayat, JM & Iguer-Ouada, M 2012, First observation of intersex in barbel, *Barbus callensis* (Valenciennes, 1842) collected in Soummam River (Algeria). *Cybium*, 36: 531-538.
- Dorafshan, S & Heyrati, FP 2006, Spawning induction in Kutum (*Rutilus frisii kutum*, Kamenskii, 1901) using carp pituitary extract or GnRH analogue combined with metoclopramide. *Aquaculture Research*, 37: 751-755, <https://doi.org/10.1111/j.1365-2109.2006.01488.x>.
- Falahatkar, B, Poursaeid, S, Langroudi, HE, Efatpanah, I, Meknatkhah, B & Rahmati, M 2013, Spawning induction in Kutum, *Rutilus frisii kutum*, Kamenski, 1901), with different hormones: Analysis of hormone profiles and induced spawning success. *Fisheries & Aquatic Life*, 21(4): 271-281.
- Farid-Pak, F 1968, Fertility of the Kutum (*Rutilus frisikutum* Kamenski, 1901). *Problems of Ichthyology*, 8: 61-68.
- Fazli, H, Bandpei, MAA, Pourgholam, R & Roohi, A 2013, Long-term changes in fecundity of the Kutum, *Rutilus frisii kutum* Kamensky, 1901, in the Caspian Sea (Osteichthyes: Cyprinidae). *Zoology in the Middle East*, 59: 51- 58.
- Gharedaashi, E, Nekoubin, H, Imanpoor, MR & Taghizadeh, V 2013, Effect of copper sulfate on the survival and growth performance of Caspian Sea kutum, *Rutilus frisii kutum*. *Springer plus*, 2: 498.
- Gholami, M, Fatemi, SMR, Falahi, M, Esmaili, A & Mashinchian, A 2010, Effects of heavy metals (copper and cadmium) and detergent (LAS) on white fish fry *Rutilus frisii kutum*. *Research Journal of Environmental Toxicology*, 4: 231-236, doi: 10.3923/rjet.2010.231.236.
- Hallajian, A 2009, Histological techniques methods. International Sturgeon Research Institute, 212 p.
- Hassell, KL, Rawson CA & Gagnon, MM 2018, First report of intersex in a lutjanid: The goldband snapper *Pristipomoides multidens*. *Journal of Fish Biology*, 93: 428-431.
- Heidari, B, Roozati, SA & Yavari, L 2010, Changes in plasma levels of steroid hormones during oocyte development of Caspian Kutum (*Rutilus frisii kutum*, Kamensky, 1901). *Animal Reproduction*, 7(4): 373-381.
- Heidari, B, Shabanipour, N, Savari, A, Yavari, V & Hosseini, N 2009, The oocyte development of Kutum, *Rutilus frisii kutum*, K. with special emphasis on the zona radiata structure. *Animal Reproduction*, 6: 465-472.
- Heyrati, FP, Mostafavi, H, Toloei, H & Dorafshan, S 2007, Induced spawning of kutum, *Rutilus frisii kutum* (Kamenskii, 1901) using (D-Ala⁶, Pro⁹-NEt) GnRH_a combined with domperidone. *Aquaculture*, 265: 288-293.
- Hinc, KJE, Blazer, VS, Schmitt, CJ, Papoulias, DM & Tillitt, DE 2009. Widespread occurrence of intersex in black basses (*Micropterus* spp.) from US rivers. *Aquatics Toxicology*, 95: 60-70.
- Hoseini, H, Tahami, MS & Hesni, MA 2015, Bioaccumulation of lead and cadmium in liver and muscle tissues of Kutum fish (Teleostei: Cyprinidae) in the southern Caspian Sea. *Iranian Journal of Ichthyology*, 2: 165-171.
- Howell, WM, Black, DA & Bortone, SA 1980, Abnormal expression of secondary sex characters in a population of mosquitofish, *Gambusia affinis holbrooki*: evidence for environmentally-induced masculinization. *Copeia*, 1980: 676-681. <https://doi.org/10.2307/1444443>.
- Ichalal, K, Ramdane, Z, Iguer-Ouada, M & Kacher, M 2016, First observation of intersex in *Trachurus trachurus* (Carangidae) from the Eastern Coast of Algeria: are nematodes the causative factor? *Cybium*, 40: 225- 233.
- Jafri, SIH & Ensor, DM 1979, Occurrence of an intersex condition in the roach, *Rutilus rutilus* (L), *Journal of Fish Biology*, 15: 547-549.
- Jobling, S, Nolan, M, Tyler, CR, Brighty, G & Sumpter, JP 1998, Widespread sexual disruption in wild fish. *Environmental Science & Technology*, 32: 2498-2506.
- Jobling, S, Coey, S & Whitmore, J.G 2002, Wild intersex roach (*Rutilus rutilus*) have reduced fertility. *Biology of Reproduction*, 67: 515-524, DOI: 10.1095/biolreprod67.2.515.

- Jones, JC & Reynolds, JD 1997, Effects of pollution on reproductive behavior of fishes. *Reviews in Fish Biology and Fisheries*, 7: 463-491.
- Koohilai, S, Oryan, S, Sahafi, HH, Mostafavi, PG & Behzadi, S 2016, Ovulation and spawning induction in Caspian Kutum (*Rutilus frisii kutum*) by administration of GnRHa and catecholaminergic pharmaceutical compounds with Ovaprim. *Aquaculture Research*, 48: 1469-1477.
- Lasheidani, MF, Balouchi, SN, Keyvan, A, Jamili, S & Falakrou, K 2008, Effects of butachlor on density, volume and number of abnormal sperms in Caspian Kutum (*Rutilus frisii Kutum*, Kamenskii, 1901). *Research Journal of Environmental Sciences*, 2: 474-482.
- Lye, CM, Frid, CLJ & Gill, ME 1998, Seasonal reproductive health of flounder *Platichthys flesus* exposed to sewage effluent. *Marine Ecology Progress Series*, 170: 249-260.
- Masouleh, FF, Amiri, BM, Mirvaghefi, AR & Nemtollahi, MA 2011, In vitro effects of diazinon on male reproductive tissue and sperm motility of Caspian kutum (*Rutilus frisii kutum*). *Research Journal of Environmental Toxicology*, 5: 108-116.
- Minier, C, Caltot, G, Leboulanger, F & Hill, EM 2000, An investigation of the incidence of intersex fish in Seine Maritime and Sussex region, *Analisis*, 28 (9): 801-806, doi:10.1051/analisis:2000280801.
- Nolan, M, Jobling, S, Brighty, G, Sumpter, JP & Tyler, CR 2001, A histological description of intersexuality in the roach (*Rutilus rutilus*). *Journal of Fish Biology*, 58: 160-176.
- Papoulias, DM, Chapman, D & Tillitt, DE 2006, Reproductive condition and occurrence of intersex in bighead carp and silver carp in the Missouri River. *Hydrobiologia*, 571: 355-360.
- Pourkazemi, M & Razikazemi, S 2011, Failure of PCR-RAPD technique to differentiate sex in Mahisefied (*Rutilus frisii kutum*) from the South Caspian Sea. *Caspian Journal of Environmental Sciences*, 9: 235-242.
- Razavi-Sayyad, BA 1990, Evaluation and management of economic fish stock of the Caspian Sea. Fishery Research organization, Guilan Province, Bandar Anzali, 86 p. [In Persian].
- Retheesh, TB, Roul, SQ, Prakasan, D, Beni, N, Thangaraja, R & Abdussamad, EM 2017, First Record of abnormal hermaphroditism in the common dolphin fish. *Coryphaena hippurus* (Linnaeus, 1758). *Thalassas*, 33: 173-177.
- Sabet, SS, Imanpoor, MR & Fatideh, BA 2009, Relation between gonadal hormones and sexual maturity of female Kutum (*Rutilus frisii kutum*) during spawning season. *Journal of Cell and Molecular Research*, 1: 96-105, <https://doi.org/10.22067/jcmr.v1i2.1344>.
- Sadovy de Mitcheson, Y & Liu, M 2008, Functional hermaphroditism in teleosts. *Fish and Fisheries*, 9: 1-43, <https://doi.org/10.1111/j.1467-2979.2007.00266>.
- Saeed, SS, Reza, IM, Bagher, AF & Saeed, G 2010. Histological study of ovarian development and sexual maturity of kutum (*Rutilus frisii kutum* Kamenskii, 1901). *World Applied Sciences Journal*, 8: 1343-1350.
- Saeidi, M, Abessi, O & Jamshidi, A 2010, Assessment of heavy metal and oil pollution of sediments of south eastern Caspian Sea using indices. *Journal of Environmental Studies*, 36(53): 21-38.
- Samarin, AM, Mojazi-Amiri, B, Soltani, M, Nazari, RM, Kamali, A & Naghavi, MR 2011, Effects of post-ovulatory oocyte ageing and temperature on egg quality in kutum (*Rutilus frisii kutum*). *World Applied Sciences Journal*, 15: 14-18.
- Sattari, M, Namin, JI, Bibak, M, Vajargah, MF, Bakhshalizadeh, S & Faggio, C 2019, Determination of trace element accumulation in gonads of *Rutilus kutum* (Kamensky, 1901) from the South Caspian Sea traceelement contaminations in gonads. Proceeding of The National Academy of Sciences, India, Section. B, Biological Sciences, <https://doi.org/10.1007/s40011-019-01150-5>.
- Savadkouhi, EB & Khara, H 2017, Effect of age on reproductive performance of Kutum, *Rutilus frisii* (Nordmann, 1840) in Shirood River, southern coast of the Caspian Sea. *Caspian Journal of Environmental Sciences*, 15: 205-212.
- Sharifpour, I, Hallajian, A & Kazemi, R 2013, Histological laboratorial methods of aquatic animal. Iranian fisheries research organization, 347 p.
- Sharyati, A 1993, Fishes of the Caspian Sea Region. Iranian Fisheries Company, Iran, 171 p. (In Persian).
- Sloof, F and Klootwijk-Vandijk, E 1982, Hermaphroditism in the bream, *Abramis brama* (L.). *Journal of Fish Disease*, 5: 79-81.
- Solé, M, Barceló, D & Porte, C 2002, Seasonal variation of plasmatic and hepatic vitellogenin and EROD activity in carp, *Cyprinus carpio*, in relation to sewage treatment plants. *Aquatic Toxicology*, 60: 233-248.

- Tyler, CR & Jobling, S 2008, Roach, sex, and gender-bending chemicals: The feminization of wild fish in English rivers. *Bioscience*, 58: 1051-1059, <https://doi.org/10.1641/B581108>.
- Tyler, CR & Sumpter, JP 1996, Oocyte growth and development in teleosts. *Reviews in Fish Biology and Fisheries*, 6: 287-318.
- Viganò, L, Arillo, A, Bottero, S, Massari, A & Mandich, A 2001, First observation of intersex cyprinids in the Po River (Italy). *Science of The Total Environment*, 269: 189-194, [https://doi.org/10.1016/S0048-9697\(00\)00821-4](https://doi.org/10.1016/S0048-9697(00)00821-4).
- Vrecl, M & Jenčič, V 2016, Occurrence of intersex in wild freshwater fish in Slovenian rivers: a histological evaluation. *Archives of Industrial Hygiene and Toxicology*, 67: 216-222, DOI: 10.1515/aiht-2016-67-2730.
- Yamamoto, T 1969, *Fish Physiology*, 3: 117-175, [https://doi.org/10.1016/S1546-5098\(08\)60113-2](https://doi.org/10.1016/S1546-5098(08)60113-2).
- Yousefian, M & Mosavi, H 2008, Spawning of south Caspian kutum (*Rutilus frisii kutum*) in most migratory rivers of south Caspian Sea. *Asian Journal of Animal and Veterinary Advances*, 3: 437-442, DOI: 10.3923/ajava.2008.437.442.
- Zahedi, S, Vaezzade, H, Rafati, M & Dangesaraki, MZ 2014, Acute toxicity and accumulation of iron, manganese and aluminum in Caspian kutum fish (*Rutilus frisii kutum*). *Iranian Journal Toxicology*, 8(24): 1028-1033, <http://ijt.arakmu.ac.ir/article-1-299>.

Bibliographic information of this paper for citing:

Tizkar, B, Dehghanzadeh, H, Hallajian, A, Zoughi Shalmani, A 2026, First evidence of intersex condition in Caspian kutum, *Rutilus kutum* (Kamenski, 1901) in the Southwest Caspian Sea. *Caspian Journal of Environmental Sciences*, 24: 633-640.
